

Some counterexamples in two-weight norm inequalities for Calderón-Zygmund operators

Michel Alexis

Joint work with José Luis Luna-Garcia, Eric Sawyer and Ignacio Uriarte-Tuero

Bonn

Friday, November 3

My favorite operator: the Hilbert transform on \mathbb{R}

My favorite operator: the Hilbert transform on \mathbb{R}


$$Hf(x) \stackrel{\text{def}}{=} \int \frac{f(t)}{x-t} dt.$$

My favorite operator: the Hilbert transform on \mathbb{R}

$$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt.$$

My favorite operator: the Hilbert transform on \mathbb{R}

$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt$. Because the kernel $\frac{1}{x-t}$ is not integrable, cancellation as $\epsilon \rightarrow 0$ is essential to make sense of this!





My favorite operator: the Hilbert transform on \mathbb{R}

$$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt. \text{ Because the}$$

kernel $\frac{1}{x-t}$ is not integrable, cancellation as $\epsilon \rightarrow 0$ is essential to make sense of this!

How does H act on functions? Cancellation transforms points with discontinuities into singularities.





My favorite operator: the Hilbert transform on \mathbb{R}

$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt$. Because the kernel $\frac{1}{x-t}$ is not integrable, cancellation as $\epsilon \rightarrow 0$ is essential to make sense of this!

How does H act on functions? Cancellation transforms points with discontinuities into singularities.

But turns out $H : L^2(dx) \rightarrow L^2(dx)$, i.e.,

$$\|Hf\|_{L^2(dx)} \lesssim \|f\|_{L^2(dx)} \text{ for all } f.$$



My favorite operator: the Hilbert transform on \mathbb{R}


$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt$. Because the kernel $\frac{1}{x-t}$ is not integrable, cancellation as $\epsilon \rightarrow 0$ is essential to make sense of this!

How does H act on functions? Cancellation transforms points with discontinuities into singularities.

But turns out $H : L^2(dx) \rightarrow L^2(dx)$, i.e.,

$$\|Hf\|_{L^2(dx)} \lesssim \|f\|_{L^2(dx)} \text{ for all } f.$$

The Hilbert transform is an example of a **Calderón-Zygmund Operator (CZO)**.



Calderón-Zygmund operators

Definition (CZO)

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$.

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

$$Tf(x) \equiv \int K(x, y) f(y) dy ,$$

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where K is smooth and satisfies the decay conditions

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}} \text{ for all } j \geq 0,$$

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where K is smooth and satisfies the decay conditions

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}} \text{ for all } j \geq 0,$$

and there exists a unit vector \mathbf{v} s.t. $|K(x, y)| \approx \frac{1}{|x - y|^n}$ when $x - y \parallel \mathbf{v}$.

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where K is smooth and satisfies the decay conditions

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}} \text{ for all } j \geq 0,$$

and there exists a unit vector \mathbf{v} s.t. $|K(x, y)| \approx \frac{1}{|x - y|^n}$ when $x - y \parallel \mathbf{v}$.

Examples include

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where K is smooth and satisfies the decay conditions

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}} \text{ for all } j \geq 0,$$

and there exists a unit vector \mathbf{v} s.t. $|K(x, y)| \approx \frac{1}{|x - y|^n}$ when $x - y \parallel \mathbf{v}$.

Examples include

- Hilbert transform H on \mathbb{R} , with $K(x, y) = \frac{1}{x - y}$.

Calderón-Zygmund operators

Definition (CZO)

Assume $T : L^2(dx) \rightarrow L^2(dx)$. Then T is a (smooth, non-degenerate) Calderón-Zygmund operator on \mathbb{R}^n if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where K is smooth and satisfies the decay conditions

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}} \text{ for all } j \geq 0,$$

and there exists a unit vector \mathbf{v} s.t. $|K(x, y)| \approx \frac{1}{|x - y|^n}$ when $x - y \parallel \mathbf{v}$.

Examples include

- Hilbert transform H on \mathbb{R} , with $K(x, y) = \frac{1}{x - y}$.
- Riesz transform R_j on \mathbb{R}^n , with $K(x, y) = \frac{x_j - y_j}{|x - y|^{n+1}}$.

L^p boundedness and relevance

L^p boundedness and relevance

Note that because a CZO

$$T : L^2(dx) \rightarrow L^2(dx),$$

L^p boundedness and relevance

Note that because a CZO

$$T : L^2(dx) \rightarrow L^2(dx) ,$$

then by CZ theory

$$T : L^p(dx) \rightarrow L^p(dx) \text{ for all } 1 < p < \infty .$$

L^p boundedness and relevance

Note that because a CZO

$$T : L^2(dx) \rightarrow L^2(dx) ,$$

then by CZ theory

$$T : L^p(dx) \rightarrow L^p(dx) \text{ for all } 1 < p < \infty .$$

Q: Why should one care about CZOs?

L^p boundedness and relevance

Note that because a CZO

$$T : L^2(dx) \rightarrow L^2(dx),$$

then by CZ theory

$$T : L^p(dx) \rightarrow L^p(dx) \text{ for all } 1 < p < \infty.$$

Q: Why should one care about CZOs?

- For Hardy functions $f \in H^p(\mathbb{C}^+)$, which are analytic on the upper half-plane, f has boundary values $u + iHu$ on $\mathbb{R} + 0i \subset \mathbb{C}$.

L^p boundedness and relevance

Note that because a CZO

$$T : L^2(dx) \rightarrow L^2(dx),$$

then by CZ theory

$$T : L^p(dx) \rightarrow L^p(dx) \text{ for all } 1 < p < \infty.$$

Q: Why should one care about CZOs?

- For Hardy functions $f \in H^p(\mathbb{C}^+)$, which are analytic on the upper half-plane, f has boundary values $u + iHu$ on $\mathbb{R} + 0i \subset \mathbb{C}$.
- The Riesz transforms pop-up in all sorts of PDEs and even some GMT problems.

When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure μ ?

When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure μ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

Suppose $\mu = w(x)dx$. Then $H : L^2(\mu) \rightarrow L^2(\mu)$ iff w satisfies the A_2 condition

$$A_2(w, w^{-1})^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q w \, dx \right) \left(\frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure μ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

Suppose $\mu = w(x)dx$. Then $H : L^2(\mu) \rightarrow L^2(\mu)$ iff w satisfies the A_2 condition

$$A_2(w, w^{-1})^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q w \, dx \right) \left(\frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any CZO.

When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure μ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

Suppose $\mu = w(x)dx$. Then $H : L^2(\mu) \rightarrow L^2(\mu)$ iff w satisfies the A_2 condition

$$A_2(w, w^{-1})^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q w \, dx \right) \left(\frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any CZO.

This is a great theorem! Boundedness of any CZO boils down to an easy-to-verify condition about the weight w !! No need to delicately estimate any nasty cancellation!

When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure μ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

Suppose $\mu = w(x)dx$. Then $H : L^2(\mu) \rightarrow L^2(\mu)$ iff w satisfies the A_2 condition

$$A_2(w, w^{-1})^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q w \, dx \right) \left(\frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any CZO.

This is a great theorem! Boundedness of any CZO boils down to an easy-to-verify condition about the weight w !! No need to delicately estimate any nasty cancellation!

Example: if $w = |x|^\alpha$ on \mathbb{R} , then $A_2(w, w^{-1}) < \infty$ iff $-1 < \alpha < 1$.

When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure μ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

Suppose $\mu = w(x)dx$. Then $H : L^2(\mu) \rightarrow L^2(\mu)$ iff w satisfies the A_2 condition

$$A_2(w, w^{-1})^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q w \, dx \right) \left(\frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any CZO.

This is a great theorem! Boundedness of any CZO boils down to an easy-to-verify condition about the weight w !! No need to delicately estimate any nasty cancellation!

Example: if $w = |x|^\alpha$ on \mathbb{R} , then $A_2(w, w^{-1}) < \infty$ iff $-1 < \alpha < 1$.

Can extend this characterisation to L^p by replacing the A_2 condition by an appropriate A_p condition.

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\nu(x).$$

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\nu(x).$$

The above is ill-posed: how should it be posed correctly?

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\nu(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 \nu dx.$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f\sigma)(x)|^2 d\omega(x) \leq C \int |f(x)\sigma|^2 \nu dx.$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f\sigma)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 \sigma^2 \nu dx.$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H(f\sigma)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.
- The (now correct) problem is about showing $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$, where $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{x-y}$

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.
- The (now correct) problem is about showing $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$, where $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{x-y}$
- One does same rescaling for other CZOs

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.
- The (now correct) problem is about showing $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$, where $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{x-y}$
- One does same rescaling for other CZOs

However $A_2(\sigma, \omega)^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q d\sigma(y) \right) \left(\frac{1}{|Q|} \int_Q d\omega(y) \right) < \infty$ isn't a sufficient condition for boundedness!

The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral $Hf(x) = \int \frac{f(y)}{x-y} dy$ should be taken with respect to the same measure that f is measurable with respect to, not w.r.t. dy
- Let's assume $\nu = \nu(x)dx$ is absolutely continuous,
- and replace f by $f\sigma$, where $\sigma \stackrel{\text{def}}{=} \nu^{-1}$.
- The (now correct) problem is about showing $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$, where $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{x-y}$
- One does same rescaling for other CZOs

However $A_2(\sigma, \omega)^2 \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left(\frac{1}{|Q|} \int_Q d\sigma(y) \right) \left(\frac{1}{|Q|} \int_Q d\omega(y) \right) < \infty$ isn't a sufficient condition for boundedness! No easy characterization in terms of weights alone!

Sawyer Testing Conditions for Hilbert transform

Sawyer Testing Conditions for Hilbert transform

Theorem (Lacey-Sawyer-Shen-Uriarte-Tuero; Lacey, 2012)

Assume ω and σ don't have common point masses and a stronger A_2 condition holds where the averages are replaced by Poisson averages. Then

$$\|H_\sigma f(x)\|_{L^2(\omega)} \lesssim \|f\|_{L^2(\sigma)} \text{ for all } f$$

if and only if the testing conditions hold, i.e.,

$$\begin{cases} \|\mathbf{1}_Q H_\sigma \mathbf{1}_Q(x)\|_{L^2(\omega)} \lesssim \|\mathbf{1}_Q\|_{L^2(\sigma)} \\ \|\mathbf{1}_Q H_\omega \mathbf{1}_Q(x)\|_{L^2(\sigma)} \lesssim \|\mathbf{1}_Q\|_{L^2(\omega)} \end{cases} \quad \text{for all cubes } Q.$$

Sawyer Testing Conditions for Hilbert transform

Theorem (Lacey-Sawyer-Shen-Uriarte-Tuero; Lacey, 2012)

Assume ω and σ don't have common point masses and a stronger A_2 condition holds where the averages are replaced by Poisson averages. Then

$$\|H_\sigma f(x)\|_{L^2(\omega)} \lesssim \|f\|_{L^2(\sigma)} \text{ for all } f$$

if and only if the testing conditions hold, i.e.,

$$\begin{cases} \|\mathbf{1}_Q H_\sigma \mathbf{1}_Q(x)\|_{L^2(\omega)} \lesssim \|\mathbf{1}_Q\|_{L^2(\sigma)} \\ \|\mathbf{1}_Q H_\omega \mathbf{1}_Q(x)\|_{L^2(\sigma)} \lesssim \|\mathbf{1}_Q\|_{L^2(\omega)} \end{cases} \quad \text{for all cubes } Q.$$

Morally, to check the norm inequality, it suffices to *test* the norm inequality over indicators of cubes.

Sawyer Testing Conditions for CZOs

For CZOs, have a similar result if the measures are “regular enough.”

Sawyer Testing Conditions for CZOs

For CZOs, have a similar result if the measures are “regular enough.”

Definition (Doubling)

A measure μ is doubling if there exists a constant C such that $\mu(2B) \leq C\mu(B)$.

Sawyer Testing Conditions for CZOs

For CZOs, have a similar result if the measures are “regular enough.”

Definition (Doubling)

A measure μ is doubling if there exists a constant C such that $\mu(2B) \leq C\mu(B)$.

Doubling measures can't locally change too much (e.g. no delta masses).

Sawyer Testing Conditions for CZOs

For CZOs, have a similar result if the measures are “regular enough.”

Definition (Doubling)

A measure μ is doubling if there exists a constant C such that $\mu(2B) \leq C\mu(B)$.

Doubling measures can't locally change too much (e.g. no delta masses).

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

Assume the measures σ, ω are doubling. If T is a CZO, then

$$T_\sigma : L^2(\sigma) \rightarrow L^2(\omega) \Leftrightarrow A_2(\sigma, \omega) < \infty \text{ and the testing conditions hold.}$$

Sawyer Testing Conditions for CZOs

For CZOs, have a similar result if the measures are “regular enough.”

Definition (Doubling)

A measure μ is doubling if there exists a constant C such that $\mu(2B) \leq C\mu(B)$.

Doubling measures can't locally change too much (e.g. no delta masses).

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

Assume the measures σ, ω are doubling. If T is a CZO, then

$$T_\sigma : L^2(\sigma) \rightarrow L^2(\omega) \Leftrightarrow A_2(\sigma, \omega) < \infty \text{ and the testing conditions hold.}$$

Theorem (Sawyer-Wick, 2022)

If the measures σ, ω are doubling, then a more technical (but similar in spirit) L^p testing theorem holds.

One-weight vs two-weight problems: A_2 vs Testing

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check:

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas A_2 is easy to check, since everything's positive and the averaging integrals are simple.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas A_2 is easy to check, since everything's positive and the averaging integrals are simple. Furthermore, A_2 robust under dilations, translations, and biLipschitz change of variable.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas A_2 is easy to check, since everything's positive and the averaging integrals are simple. Furthermore, A_2 robust under dilations, translations, and biLipschitz change of variable.
- A_2 is *stable* under biLipschitz change of variable.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas A_2 is easy to check, since everything's positive and the averaging integrals are simple. Furthermore, A_2 robust under dilations, translations, and biLipschitz change of variable.
- A_2 is *stable* under biLipschitz change of variable. So one-weight norm inequalities are too.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas A_2 is easy to check, since everything's positive and the averaging integrals are simple. Furthermore, A_2 robust under dilations, translations, and biLipschitz change of variable.
- A_2 is *stable* under biLipschitz change of variable. So one-weight norm inequalities are too. Not the case for two-weight norm inequalities.

One-weight vs two-weight problems: A_2 vs Testing

- In the one-weight case, A_2 holds iff your operator is bounded.
- In the two-weight case, A_2 cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas A_2 is easy to check, since everything's positive and the averaging integrals are simple. Furthermore, A_2 robust under dilations, translations, and biLipschitz change of variable.
- A_2 is *stable* under biLipschitz change of variable. So one-weight norm inequalities are too. Not the case for two-weight norm inequalities.

Theorem (Lacey-Sawyer-Urriarte-Tuero, 2012)

There exist measures σ, ω and a biLipchitz map φ such that the norm inequality for H holds w.r.t. (σ, ω) , but does not w.r.t. the pushforwards $(\varphi_\sigma, \varphi_*\omega)$.*

Two-weight norms are biLipschitz unstable: 1st example

Two-weight norms are biLipschitz unstable: 1st example


Recall the middle-third Cantor set E and Cantor measure σ .

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .




Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .




Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .



Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .



σ assigns equal weight to these intervals

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .



Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .


Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.

— — — — — — — —

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .


Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .


Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .


Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.




ω = sum of delta masses at these zeros of $H\sigma(x)$.

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.




ω = sum of delta masses at these zeros of $H\sigma(x)$.

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.




ω = sum of delta masses at these zeros of $H\sigma(x)$.

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x) = 0$. Then one can show $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Now, shift the point masses to blow-up points of $H\sigma(x)$ via a biLipschitz map.


ω = sum of delta masses at these zeros of $H\sigma(x)$.

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x) = 0$. Then one can show $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Now, shift the point masses to blow-up points of $H\sigma(x)$ via a biLipschitz map.


ω = sum of delta masses at these **blow-up points** of $H\sigma(x)$.

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x) = 0$. Then one can show $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Now, shift the point masses to blow-up points of $H\sigma(x)$ via a biLipschitz map.


ω = sum of delta masses at these blow-up points of $H\sigma(x)$.

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x) = \infty$. Then one can show $H_\sigma : L^2(\sigma) \not\rightarrow L^2(\omega)$

Two-weight norms are biLipschitz unstable: 1st example

Recall the middle-third Cantor set E and Cantor measure σ .

Let's graph $H(\sigma)(x) = \int \frac{d\sigma(y)}{x-y}$.



Now, shift the point masses to blow-up points of $H\sigma(x)$ via a biLipschitz map.

ω = sum of delta masses at these blow-up points of $H\sigma(x)$.

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x) = \infty$. Then one can show $H_\sigma : L^2(\sigma) \nrightarrow L^2(\omega)$

Morally, this is why you need testing: one can only characterize the norm inequality with some biLipschitz unstable condition.

Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)


Consider the Riesz transform R_2 on \mathbb{R}^2 . For every angle $\theta > 0$, there exist doubling measures σ, ω such that the norm inequality for R_2 holds w.r.t. (σ, ω) , but does not w.r.t. the rotated measures $(\sigma_\theta, \omega_\theta)$.

Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Consider the Riesz transform R_2 on \mathbb{R}^2 . For every angle $\theta > 0$, there exist doubling measures σ, ω such that the norm inequality for R_2 holds w.r.t. (σ, ω) , but does not w.r.t. the rotated measures $(\sigma_\theta, \omega_\theta)$.

We construct measures that look like sheet metal!




Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Consider the Riesz transform R_2 on \mathbb{R}^2 . For every angle $\theta > 0$, there exist doubling measures σ, ω such that the norm inequality for R_2 holds w.r.t. (σ, ω) , but does not w.r.t. the rotated measures $(\sigma_\theta, \omega_\theta)$.

We construct measures that look like sheet metal!




Start with R_2 : it points vertically along the level curves of the sheet metal.

Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Consider the Riesz transform R_2 on \mathbb{R}^2 . For every angle $\theta > 0$, there exist doubling measures σ, ω such that the norm inequality for R_2 holds w.r.t. (σ, ω) , but does not w.r.t. the rotated measures $(\sigma_\theta, \omega_\theta)$.

We construct measures that look like sheet metal!



Start with R_2 : it points vertically along the level curves of the sheet metal.
Rotate R_2 slightly: it sees all the ripples all of a sudden.

Rotational instability on doubling measures: the key ideas

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) for which norm inequality holds with respect to R_2 , but fails with respect to R_1 .

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) for which norm inequality holds with respect to R_2 , but fails with respect to R_1 .



Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but the norm inequality fails for H .

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) for which norm inequality holds with respect to R_2 , but fails with respect to R_1 .



Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but the norm inequality fails for H .



Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) for which norm inequality holds with respect to R_2 , but fails with respect to R_1 .



Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but the norm inequality fails for H .



Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)} . \text{ Actually done by Nazarov!}$$

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) for which norm inequality holds with respect to R_2 , but fails with respect to R_1 .



Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but the norm inequality fails for H .

↑↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)} . \text{ Actually done by Nazarov!}$$

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) for which norm inequality holds with respect to R_2 , but fails with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but the norm inequality fails for H .


↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)} . \text{ Actually done by Nazarov!}$$


The dyadic problem

The dyadic problem




- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.

The dyadic problem




- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.

The dyadic problem




- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k}), j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.
- $A_2^{\mathcal{D}}(\sigma, \omega)^2 \equiv \sup_{I \in \mathcal{D}} (E_I \sigma) (E_I \omega)$.

The dyadic problem



- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.
- $A_2^{\mathcal{D}}(\sigma, \omega)^2 \equiv \sup_{I \in \mathcal{D}} (E_I \sigma)(E_I \omega)$.
- The dyadic square function $Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}$ measures the oscillation/discontinuities of f .


The dyadic problem



- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.
- $A_2^{\mathcal{D}}(\sigma, \omega)^2 \equiv \sup_{I \in \mathcal{D}} (E_I \sigma)(E_I \omega)$.
- The dyadic square function $Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J|} \mathbf{1}_J(x)$ measures the oscillation/discontinuities of f .

Nazarov found dyadically doubling measures (σ, ω) on $[0, 1]$ which satisfy $A_2^{\mathcal{D}}(\sigma, \omega)$, but fail the testing condition for the dyadic square function.

The dyadic problem




- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.
- $A_2^{\mathcal{D}}(\sigma, \omega)^2 \equiv \sup_{I \in \mathcal{D}} (E_I \sigma) (E_I \omega)$.
- The dyadic square function $Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J|} \mathbf{1}_J(x)$ measures the oscillation/discontinuities of f .

Nazarov found dyadically doubling measures (σ, ω) on $[0, 1]$ which satisfy $A_2^{\mathcal{D}}(\sigma, \omega)$, but fail the testing condition for the dyadic square function.

How?

The dyadic problem




- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.
- $A_2^{\mathcal{D}}(\sigma, \omega)^2 \equiv \sup_{I \in \mathcal{D}} (E_I \sigma)(E_I \omega)$.
- The dyadic square function $Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}$ measures the oscillation/discontinuities of f .

Nazarov found dyadically doubling measures (σ, ω) on $[0, 1]$ which satisfy $A_2^{\mathcal{D}}(\sigma, \omega)$, but fail the testing condition for the dyadic square function.

How? Nazarov showed existence via a non-constructive Bellman proof.

The dyadic problem



- the dyadic grid is given by the intervals $[j2^{-k}, (j+1)2^{-k})$, $j, k \in \mathbb{Z}$.
- μ is dyadically doubling if $\mu(\pi I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.
- $A_2^{\mathcal{D}}(\sigma, \omega)^2 \equiv \sup_{I \in \mathcal{D}} (E_I \sigma)(E_I \omega)$.
- The dyadic square function $Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}$ measures the oscillation/discontinuities of f .

Nazarov found dyadically doubling measures (σ, ω) on $[0, 1]$ which satisfy $A_2^{\mathcal{D}}(\sigma, \omega)$, but fail the testing condition for the dyadic square function.

How? Nazarov showed existence via a non-constructive Bellman proof. Kakaroumpas-Treil got explicit weights for an essentially equivalent pb.

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) which are well-behaved with respect to R_2 , but badly behaved with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but fail the testing condition for H .

↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) which are well-behaved with respect to R_2 , but badly behaved with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but fail the testing condition for H .

↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

A transference principle: from dyadic square function to H

A transference principle: from dyadic square function to H




A transference principle: from dyadic square function to H




σ, ω are dyadic step functions s.t.
testing fails for S .

A transference principle: from dyadic square function to H




σ, ω are dyadic step functions s.t.
testing fails for S .


$\sigma = E_{[0,1]} \sigma + \sum_{I \in \mathcal{D}} c_I h_I$, where h_I
is a Haar function



A transference principle: from dyadic square function to H




\Rightarrow




σ, ω are dyadic step functions s.t.
testing fails for S .


$\sigma = E_{[0,1]} \sigma + \sum_{I \in \mathcal{D}} c_I h_I$, where h_I
is a Haar function



A transference principle: from dyadic square function to H




\Rightarrow




σ, ω are dyadic step functions s.t. testing fails for S . \Rightarrow

Spacing out Haar frequencies of σ and ω makes testing fail for H


$\sigma = E_{[0,1]} \sigma + \sum_{I \in \mathcal{D}} c_I h_I$, where h_I is a Haar function



A transference principle: from dyadic square function to H



\Rightarrow




σ, ω are dyadic step functions s.t. testing fails for S . \Rightarrow


Spacing out Haar frequencies of σ and ω makes testing fail for H

$\sigma = E_{[0,1]} \sigma + \sum_{I \in \mathcal{D}} c_I h_I$, where h_I is a Haar function \Rightarrow

$\sigma = E_{[0,1]} \sigma + \sum_{I \in \mathcal{S}} c_I s'_I$, where \mathcal{S} is a sparser set of cubes



\Rightarrow



How the s_k 's give us estimates for the Hilbert transform

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly,

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly,

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.
By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\omega^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly. Real part gives $(Hs_k)^2 \rightarrow (s_k)^2 = \mathbf{1}_{[0,1]}^2$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} \mathbf{1}_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly. Real part gives $(Hs_k)^2 \rightarrow (s_k)^2 = \mathbf{1}_{[0,1]}^2$ weakly.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} \mathbf{1}_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly. Real part gives $(Hs_k)^2 \rightarrow (s_k)^2 = \mathbf{1}_{[0,1]}^2$ weakly.

Looking at $(s_k + iHs_k)^3 \rightarrow 0$ weakly yields last term $\rightarrow 0$.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x)$ equals

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} \mathbf{1}_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly. Real part gives $(Hs_k)^2 \rightarrow (s_k)^2 = \mathbf{1}_{[0,1]}^2$ weakly.

Looking at $(s_k + iHs_k)^3 \rightarrow 0$ weakly yields last term $\rightarrow 0$.

How the s_k 's give us estimates for the Hilbert transform

Simple example: $\sigma = A\mathbf{1}_{[0,1]} + c_\sigma s_k^{[0,1]}$ and $\omega = B\mathbf{1}_{[0,1]} + c_\omega s_k^{[0,1]}$.

Then $\int_{[0,1]} |H\mathbf{1}_{[0,1]}\sigma(x)|^2 d\omega(x) \rightarrow \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\sigma^2 B$.

$$\begin{aligned} & \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 B dx + c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A)^2 s_k^{[0,1]} dx \\ & + 2c_\sigma \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) B dx + 2c_\sigma c_\omega \int_{[0,1]} (H\mathbf{1}_{[0,1]}A) (Hs_k^{[0,1]}) s_k^{[0,1]} dx \\ & + c_\sigma^2 \int_{[0,1]} \mathbf{1}_{[0,1]} (Hs_k^{[0,1]})^2 B dx + c_\sigma^2 \int_{[0,1]} (Hs_k^{[0,1]})^2 s_k^{[0,1]} dx \end{aligned}$$

Since $s_k \equiv s_k^{[0,1]} \rightarrow 0$ weakly, then $s_k + iHs_k \rightarrow 0$ weakly.

By complex function theory, $(s_k + iHs_k)^2 \rightarrow 0$ weakly. Imaginary part gives $(Hs_k) s_k \rightarrow 0$ weakly. Real part gives $(Hs_k)^2 \rightarrow (s_k)^2 = \mathbf{1}_{[0,1]}^2$ weakly.

Looking at $(s_k + iHs_k)^3 \rightarrow 0$ weakly yields last term $\rightarrow 0$.

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) which are well-behaved with respect to R_2 , but badly behaved with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but fail the testing condition for H .

↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J^-} f - E_{J^+} f)^2 |J| \mathbf{1}_J(x)}.$$

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) which are well-behaved with respect to R_2 , but badly behaved with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but fail the testing condition for H .

↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J^-} f - E_{J^+} f)^2 |J| \mathbf{1}_J(x)}.$$

From \mathbb{R} to \mathbb{R}^2

From \mathbb{R} to \mathbb{R}^2

Tensor the $1D$ weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

From \mathbb{R} to \mathbb{R}^2

Tensor the $1D$ weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

Decompose $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$ into base frequency elements.

From \mathbb{R} to \mathbb{R}^2

Tensor the 1D weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

Decompose $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$ into base frequency elements.

Must now consider R_1 acting on the horizontal oscillating functions

$$s_k^{Q, \text{horizontal}} \equiv \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) (x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \text{ where cube } Q \equiv Q_1 \times Q_2.$$

From \mathbb{R} to \mathbb{R}^2

Tensor the 1D weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

Decompose $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$ into base frequency elements.

Must now consider R_1 acting on the horizontal oscillating functions

$$s_k^{Q, \text{horizontal}} \equiv \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) (x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \text{ where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ① $R_1 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow C_n \left(Hs_k^{Q_1} \right) \otimes \mathbf{1}_{Q_2}$ in $L^p(dx)$ for all $1 < p < \infty$.
- ② $R_2 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow 0$ in $L^p(dx)$ for all $1 < p < \infty$.

From \mathbb{R} to \mathbb{R}^2

Tensor the 1D weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

Decompose $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$ into base frequency elements.

Must now consider R_1 acting on the horizontal oscillating functions

$$s_k^{Q, \text{horizontal}} \equiv \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) (x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \text{ where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ① $R_1 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow C_n \left(Hs_k^{Q_1} \right) \otimes \mathbf{1}_{Q_2}$ in $L^p(dx)$ for all $1 < p < \infty$.
- ② $R_2 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow 0$ in $L^p(dx)$ for all $1 < p < \infty$.

Proofs of both statements use the alternating series test.

From \mathbb{R} to \mathbb{R}^2

Tensor the 1D weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

Decompose $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$ into base frequency elements.

Must now consider R_1 acting on the horizontal oscillating functions

$$s_k^{Q, \text{horizontal}} \equiv \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right)(x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \text{ where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ① $R_1 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow C_n \left(Hs_k^{Q_1} \right) \otimes \mathbf{1}_{Q_2}$ in $L^p(dx)$ for all $1 < p < \infty$.
- ② $R_2 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow 0$ in $L^p(dx)$ for all $1 < p < \infty$.

Proofs of both statements use the alternating series test.

By (1), the testing condition for R_1 fails, like for the Hilbert transform.

From \mathbb{R} to \mathbb{R}^2

Tensor the 1D weights of Nazarov with $\mathbf{1}_{[0,1]}(x_2)$ to get “sheet metal” weights σ, ω .

Decompose $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$ into base frequency elements.

Must now consider R_1 acting on the horizontal oscillating functions

$$s_k^{Q, \text{horizontal}} \equiv \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right)(x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \text{ where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ① $R_1 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow C_n \left(Hs_k^{Q_1} \right) \otimes \mathbf{1}_{Q_2}$ in $L^p(dx)$ for all $1 < p < \infty$.
- ② $R_2 \left(s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow 0$ in $L^p(dx)$ for all $1 < p < \infty$.

Proofs of both statements use the alternating series test.

By (1), the testing condition for R_1 fails, like for the Hilbert transform.

By (2), R_2 doesn't interact with the horizontal oscillation, so testing for R_2 holds!

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) which are well-behaved with respect to R_2 , but badly behaved with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but fail the testing condition for H .

↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J^-} f - E_{J^+} f)^2 |J| \mathbf{1}_J(x)}.$$

Rotational instability on doubling measures: the key ideas

We really just need to find doubling measures (σ, ω) which are well-behaved with respect to R_2 , but badly behaved with respect to R_1 .

↑ (done by us)

Find doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy $A_2(\sigma, \omega) < \infty$, but fail the testing condition for H .

↑ (done by Nazarov)

Find dyadically doubling measures (σ, ω) on $[0, 1] \subset \mathbb{R}$ which satisfy the dyadic $A_2^{\mathcal{D}}(\sigma, \omega) < \infty$, but fail the testing condition for the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

The L^p case, $1 < p < \infty$

The L^p case, $1 < p < \infty$

Conjecture (A simple conjecture)

$T_\sigma : L^p(\sigma) \rightarrow L^p(\omega) \Leftrightarrow A_p(\sigma, \omega) < \infty$ and the L^p -testing conditions hold.

The L^p case, $1 < p < \infty$

Conjecture (A simple conjecture)

$T_\sigma : L^p(\sigma) \rightarrow L^p(\omega) \Leftrightarrow A_p(\sigma, \omega) < \infty$ and the L^p -testing conditions hold.

Here the scalar A_p constant is $A_p(\sigma, \omega) \equiv \sup_Q (E_Q \sigma)^{\frac{1}{p}} (E_Q \omega)^{\frac{1}{p'}}.$

The L^p case, $1 < p < \infty$

Conjecture (A simple conjecture)

$T_\sigma : L^p(\sigma) \rightarrow L^p(\omega) \Leftrightarrow A_p(\sigma, \omega) < \infty$ and the L^p -testing conditions hold.

Here the scalar A_p constant is $A_p(\sigma, \omega) \equiv \sup_Q (E_Q \sigma)^{\frac{1}{p}} (E_Q \omega)^{\frac{1}{p'}}.$

Theorem (Sawyer-Wick 2022 + A.-Luna-Sawyer-Uriarte-Tueoro 2023)

Let σ, ω be doubling. If T is a CZO, then

$T_\sigma : L^p(\sigma) \rightarrow L^p(\omega) \Leftrightarrow A_p^{\ell^2}(\sigma, \omega) < \infty$ and the L^p -testing conditions hold.

The L^p case, $1 < p < \infty$

Conjecture (A simple conjecture)

$T_\sigma : L^p(\sigma) \rightarrow L^p(\omega) \Leftrightarrow A_p(\sigma, \omega) < \infty$ and the L^p -testing conditions hold.

Here the scalar A_p constant is $A_p(\sigma, \omega) \equiv \sup_Q (E_Q \sigma)^{\frac{1}{p}} (E_Q \omega)^{\frac{1}{p'}}.$

Theorem (Sawyer-Wick 2022 + A.-Luna-Sawyer-Uriarte-Tueoro 2023)

Let σ, ω be doubling. If T is a CZO, then

$T_\sigma : L^p(\sigma) \rightarrow L^p(\omega) \Leftrightarrow A_p^{\ell^2}(\sigma, \omega)$ and the L^p -testing conditions hold.

Here the quadratic, or vector-valued $A_p^{\ell^2}(\sigma, \omega)$ condition is

$$\left\| \left\{ \sum_Q a_Q^2 (E_Q \sigma)^2 \mathbf{1}_Q \right\}^{\frac{1}{2}} \right\|_{L^p(\omega)} \lesssim \left\| \left\{ \sum_Q a_Q^2 \mathbf{1}_Q \right\}^{\frac{1}{2}} \right\|_{L^p(\sigma)},$$

which must hold over all sequences of cubes $\{Q\}$, sequences of reals $\{a_Q\}$.

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

No!

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?


Theorem (A.-Luna-Sawyer-Uriarte-Tuero 2023)

Let $p \neq 2$. There exist doubling measures σ, ω on \mathbb{R}^2 for which the scalar A_p and testing conditions for R_2 hold, and yet $(R_2)_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$.

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

Theorem (A.-Luna-Sawyer-Uriarte-Tuero 2023)


Let $p \neq 2$. There exist doubling measures σ, ω on \mathbb{R}^2 for which the scalar A_p and testing conditions for R_2 hold, and yet $(R_2)_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$.



Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

Theorem (A.-Luna-Sawyer-Uriarte-Tuero 2023)

Let $p \neq 2$. There exist doubling measures σ, ω on \mathbb{R}^2 for which the scalar A_p and testing conditions for R_2 hold, and yet $(R_2)_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$.




$A_p^{\ell^2}(\sigma, \omega)$ fails “along” \mathbf{e}_1

Choose σ, ω dyadically doubling on \mathbb{R} which fail the $A_p^{\ell^2}(\sigma, \omega)$ condition.

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

Theorem (A.-Luna-Sawyer-Uriarte-Tuero 2023)

Let $p \neq 2$. There exist doubling measures σ, ω on \mathbb{R}^2 for which the scalar A_p and testing conditions for R_2 hold, and yet $(R_2)_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$.




$A_p^{\ell^2}(\sigma, \omega)$ fails “along” \mathbf{e}_1

Choose σ, ω dyadically doubling on \mathbb{R} which fail the $A_p^{\ell^2}(\sigma, \omega)$ condition. Apply Nazarov’s remodeling, then tensor with 1 in the \mathbf{e}_2 direction to get σ, ω doubling on \mathbb{R}^2 .

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

Theorem (A.-Luna-Sawyer-Uriarte-Tuero 2023)

Let $p \neq 2$. There exist doubling measures σ, ω on \mathbb{R}^2 for which the scalar A_p and testing conditions for R_2 hold, and yet $(R_2)_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$.




Choose σ, ω dyadically doubling on \mathbb{R} which fail the $A_p^{\ell^2}(\sigma, \omega)$ condition. Apply Nazarov's remodeling, then tensor with 1 in the \mathbf{e}_2 direction to get σ, ω doubling on \mathbb{R}^2 .

Can we weaken $A_p^{\ell^2}(\sigma, \omega)$ to $A_p(\sigma, \omega)$ in the $T1$ theorem?

Theorem (A.-Luna-Sawyer-Uriarte-Tuero 2023)

Let $p \neq 2$. There exist doubling measures σ, ω on \mathbb{R}^2 for which the scalar A_p and testing conditions for R_2 hold, and yet $(R_2)_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$.

(Testing condition for R_2) $\approx A_p(\sigma, \omega)$



$A_p^{\ell^2}(\sigma, \omega)$ fails "along" \mathbf{e}_1

Choose σ, ω dyadically doubling on \mathbb{R} which fail the $A_p^{\ell^2}(\sigma, \omega)$ condition. Apply Nazarov's remodeling, then tensor with 1 in the \mathbf{e}_2 direction to get σ, ω doubling on \mathbb{R}^2 .

Thank you for Listening!