

# BiLipschitz (in)stability of weighted norm inequalities for singular integral operators

Michel Alexis

Joint work with José Luis Luna-Garcia, Eric Sawyer and Ignacio Uriarte-Tuero

McMaster University

February 21, 2023

# My favorite operator: the Hilbert transform on $\mathbb{R}$

# My favorite operator: the Hilbert transform on $\mathbb{R}$

$$Hf(x) \stackrel{\text{def}}{=} \int \frac{f(t)}{x-t} dt.$$

# My favorite operator: the Hilbert transform on $\mathbb{R}$

$$Hf(x) \stackrel{\text{def}}{=} \int \frac{f(t)}{x-t} dt. \text{ Is } H$$

well-defined?

# My favorite operator: the Hilbert transform on $\mathbb{R}$


$$Hf(x) \stackrel{\text{def}}{=} \int \frac{f(t)}{x-t} dt. \text{ Is } H$$

well-defined? Note, kernel  $\frac{1}{x-t}$  is not integrable:

# My favorite operator: the Hilbert transform on $\mathbb{R}$

$$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt. \text{ Is } H$$

well-defined? Note, kernel  $\frac{1}{x-t}$  is not integrable: cancellation as  $\epsilon \rightarrow 0$  is essential to make sense of this!





# My favorite operator: the Hilbert transform on $\mathbb{R}$

$$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt. \text{ Is } H$$

well-defined? Note, kernel  $\frac{1}{x-t}$  is not integrable: cancellation as  $\epsilon \rightarrow 0$  is essential to make sense of this!

How does  $H$  act on functions? Cancellation transforms points of discontinuity into singularities.





# My favorite operator: the Hilbert transform on $\mathbb{R}$

$$Hf(x) \stackrel{\text{def}}{=} \lim_{\epsilon \rightarrow 0} \int_{\{|x-t|>\epsilon\}} \frac{f(t)}{x-t} dt. \text{ Is } H$$

well-defined? Note, kernel  $\frac{1}{x-t}$  is not integrable: cancellation as  $\epsilon \rightarrow 0$  is essential to make sense of this!

How does  $H$  act on functions? Cancellation transforms points of discontinuity into singularities.

The Hilbert transform is an example of a **Calderón-Zygmund Operator** (CZO).



# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition.

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition.

Examples include

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition.

Examples include

- Hilbert transform  $H$  on  $\mathbb{R}$ , with  $K(x, y) = \frac{1}{x-y}$ .

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition.

Examples include

- Hilbert transform  $H$  on  $\mathbb{R}$ , with  $K(x, y) = \frac{1}{x-y}$ .
- Riesz transform  $R_j$  on  $\mathbb{R}^n$ , with  $K(x, y) = \frac{x_j - y_j}{|x-y|^{n+1}}$ .

# Calderón-Zygmund operators

## Definition (CZOs)

$T$  is a (smooth) Calderón-Zygmund operator if

$$Tf(x) \equiv \int K(x, y) f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition.

Examples include

- Hilbert transform  $H$  on  $\mathbb{R}$ , with  $K(x, y) = \frac{1}{x-y}$ .
- Riesz transform  $R_j$  on  $\mathbb{R}^n$ , with  $K(x, y) = \frac{x_j - y_j}{|x - y|^{n+1}}$ .
- Beurling transform on  $\mathbb{C}$ , with  $K(z, w) = \frac{1}{(z-w)^2}$ .

# Why should one care about CZOs?

Say  $f$  is a real-valued function in  $\mathbb{R}$ .

# Why should one care about CZOs?

Say  $f$  is a real-valued function in  $\mathbb{R}$ .

Q: Is  $f$  the real part of some analytic function on  $\mathbb{C}^+$ ?

# Why should one care about CZOs?

Say  $f$  is a real-valued function in  $\mathbb{R}$ .

Q: Is  $f$  the real part of some analytic function on  $\mathbb{C}^+$ ?

A: Yes, if  $f$  has sufficient decay!

# Why should one care about CZOs?

Say  $f$  is a real-valued function in  $\mathbb{R}$ .

Q: Is  $f$  the real part of some analytic function on  $\mathbb{C}^+$ ?

A: Yes, if  $f$  has sufficient decay!

For any point  $z \in \mathbb{C}^+$ , define  $F(z) \stackrel{\text{def}}{=} \int_{\mathbb{R}} \frac{f(t)}{t-z} dt$ .


# Why should one care about CZOs?

Say  $f$  is a real-valued function in  $\mathbb{R}$ .

Q: Is  $f$  the real part of some analytic function on  $\mathbb{C}^+$ ?

A: Yes, if  $f$  has sufficient decay!

For any point  $z \in \mathbb{C}^+$ , define  $F(z) \stackrel{\text{def}}{=} \int_{\mathbb{R}} \frac{f(t)}{t-z} dt$ . Then  $F$  is indeed holomorphic in variable  $z$  as it varies in  $\mathbb{C}^+$ .




# Why should one care about CZOs?

Say  $f$  is a real-valued function in  $\mathbb{R}$ .

Q: Is  $f$  the real part of some analytic function on  $\mathbb{C}^+$ ?

A: Yes, if  $f$  has sufficient decay!

For any point  $z \in \mathbb{C}^+$ , define  $F(z) \stackrel{\text{def}}{=} \int_{\mathbb{R}} \frac{f(t)}{t-z} dt$ . Then  $F$  is indeed holomorphic in variable  $z$  as it varies in  $\mathbb{C}^+$ .



One can check that for  $z = x + iy$ , as  $y \rightarrow 0^+$ , we get

$$\lim_{y \rightarrow 0^+} F(x + iy) = f(x) + iHf(x),$$

# How does $H$ affect $L^2(dx)$ ?

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

## Theorem

$H$  is a bounded from  $L^2(dx)$  to itself, i.e. there exists some constant  $C > 0$  such that

$$\|Hf(x)\|_{L^2(dx)} \leq C \|f\|_{L^2(dx)}.$$

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

## Theorem

$H$  is a bounded from  $L^2(dx)$  to itself, i.e. there exists some constant  $C > 0$  such that

$$\|Hf(x)\|_{L^2(dx)} \leq C \|f\|_{L^2(dx)}.$$

In this talk, we'll only be concerned with “nice” CZOs,

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

## Theorem

$H$  is a bounded from  $L^2(dx)$  to itself, i.e. there exists some constant  $C > 0$  such that

$$\|Hf(x)\|_{L^2(dx)} \leq C \|f\|_{L^2(dx)}.$$

In this talk, we'll only be concerned with “nice” CZOs, i.e. CZOs  $T$  which are

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

## Theorem

$H$  is a bounded from  $L^2(dx)$  to itself, i.e. there exists some constant  $C > 0$  such that

$$\|Hf(x)\|_{L^2(dx)} \leq C \|f\|_{L^2(dx)}.$$

In this talk, we'll only be concerned with “nice” CZOs, i.e. CZOs  $T$  which are

- bounded on  $L^2(dx)$ .

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

## Theorem

$H$  is a bounded from  $L^2(dx)$  to itself, i.e. there exists some constant  $C > 0$  such that

$$\|Hf(x)\|_{L^2(dx)} \leq C \|f\|_{L^2(dx)}.$$

In this talk, we'll only be concerned with “nice” CZOs, i.e. CZOs  $T$  which are

- bounded on  $L^2(dx)$ .
- non-degenerate kernels  $K$ , i.e.  $|K(x, y)| \approx \frac{1}{|x-y|^n}$ .

# How does $H$ affect $L^2(dx)$ ?

Recall  $L^2(dx)$  is the vector space of functions for which

$$\|f\|_{L^2(dx)} \stackrel{\text{def}}{=} \left( \int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < \infty.$$

## Theorem

$H$  is a bounded from  $L^2(dx)$  to itself, i.e. there exists some constant  $C > 0$  such that

$$\|Hf(x)\|_{L^2(dx)} \leq C \|f\|_{L^2(dx)}.$$

In this talk, we'll only be concerned with “nice” CZOs, i.e. CZOs  $T$  which are

- bounded on  $L^2(dx)$ .
- non-degenerate kernels  $K$ , i.e.  $|K(x, y)| \approx \frac{1}{|x-y|^n}$ .

Hilbert transform, Riesz transforms and Beurling transform are all “nice.”

When does  $H : L^2(\mu) \rightarrow L^2(\mu)$  for a general measure  $\mu$ ?

# When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure $\mu$ ?

Q: For which measures do we have  $\|Hf\|_{L^2(\mu)} \leq C\|f\|_{L^2(\mu)}$ ?

# When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure $\mu$ ?

Q: For which measures do we have  $\|Hf\|_{L^2(\mu)} \leq C \|f\|_{L^2(\mu)}$ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

*The above inequality holds if and only if  $\mu = w(x) dx$  is an absolutely continuous measure and has finite  $A_2$  characteristic,*

$$[w]_{A_2} \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \, dx \right) \left( \frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

# When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure $\mu$ ?

Q: For which measures do we have  $\|Hf\|_{L^2(\mu)} \leq C\|f\|_{L^2(\mu)}$ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

*The above inequality holds if and only if  $\mu = w(x) dx$  is an absolutely continuous measure and has finite  $A_2$  characteristic,*

$$[w]_{A_2} \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \, dx \right) \left( \frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any nice CZO.

# When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure $\mu$ ?

Q: For which measures do we have  $\|Hf\|_{L^2(\mu)} \leq C \|f\|_{L^2(\mu)}$ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

*The above inequality holds if and only if  $\mu = w(x) dx$  is an absolutely continuous measure and has finite  $A_2$  characteristic,*

$$[w]_{A_2} \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \, dx \right) \left( \frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any nice CZO.

This is a great theorem! Boundedness of any nice CZO boils down to an easy-to-verify condition about the weight  $w$ !! Don't need to delicately estimate any nasty cancellation!

# When does $H : L^2(\mu) \rightarrow L^2(\mu)$ for a general measure $\mu$ ?

Q: For which measures do we have  $\|Hf\|_{L^2(\mu)} \leq C \|f\|_{L^2(\mu)}$ ?

Theorem (Hunt-Muckenhoupt-Wheeden, 1973)

*The above inequality holds if and only if  $\mu = w(x) dx$  is an absolutely continuous measure and has finite  $A_2$  characteristic,*

$$[w]_{A_2} \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \, dx \right) \left( \frac{1}{|Q|} \int_Q w^{-1} \, dx \right) < \infty.$$

Same is true for any nice CZO.

This is a great theorem! Boundedness of any nice CZO boils down to an easy-to-verify condition about the weight  $w$ !! Don't need to delicately estimate any nasty cancellation!

Calculus Exercise: show that if  $w = |x|^\alpha$  on  $\mathbb{R}$ , then  $A_2(w, w^{-1}) < \infty$  if and only if  $-1 < \alpha < 1$ .

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\nu(x).$$

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\nu(x).$$

The above is ill-posed: how should it be posed correctly?

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\nu(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 \nu dx.$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f\sigma)(x)|^2 d\omega(x) \leq C \int |f(x)\sigma|^2 \nu dx.$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f\sigma)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 \sigma^2 \nu dx.$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H(f\sigma)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .
- The (now correct) problem is about showing  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$ ,  
where  $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{|x-y|}$

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .
- The (now correct) problem is about showing  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$ , where  $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{|x-y|}$
- One does same rescaling for other nice CZOs

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .
- The (now correct) problem is about showing  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$ , where  $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{|x-y|}$
- One does same rescaling for other nice CZOs

However  $A_2(\sigma, \omega) \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left( \frac{1}{|Q|} \int_Q d\sigma(y) \right) \left( \frac{1}{|Q|} \int_Q d\omega(y) \right) < \infty$  isn't a sufficient condition for boundedness!

# The two-weight problem: when is $H : L^2(\nu) \rightarrow L^2(\omega)$ ?

$$\int |H_\sigma(f)(x)|^2 d\omega(x) \leq C \int |f(x)|^2 d\sigma(x).$$

The above is ill-posed: how should it be posed correctly?

- the integral  $Hf(x) = \int \frac{f(y)}{|x-y|} dy$  should be taken with respect to the same measure that  $f$  is measurable with respect to, not w.r.t.  $dy$
- Let's assume  $\nu = \nu(x)dx$  is absolutely continuous,
- and replace  $f$  by  $f\sigma$ , where  $\sigma \stackrel{\text{def}}{=} \nu^{-1}$ .
- The (now correct) problem is about showing  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$ , where  $H_\sigma f(x) \stackrel{\text{def}}{=} \int \frac{f(y) d\sigma(y)}{|x-y|}$
- One does same rescaling for other nice CZOs

However  $A_2(\sigma, \omega) \stackrel{\text{def}}{=} \sup_{Q \subset \mathbb{R}} \left( \frac{1}{|Q|} \int_Q d\sigma(y) \right) \left( \frac{1}{|Q|} \int_Q d\omega(y) \right) < \infty$  isn't a sufficient condition for boundedness! No easy characterization in terms of weights alone!

# Sawyer Testing Conditions for Hilbert transform

# Sawyer Testing Conditions for Hilbert transform

Theorem (Lacey-Sawyer-Shen-Uriarte-Tuero; Lacey, 2012)

Assume  $\omega$  and  $\sigma$  don't have common point masses and a stronger  $A_2$  condition holds where the averages are replaced by Poisson averages. Then

$$\|H_\sigma f(x)\|_{L^2(\omega)} \leq C \|f\|_{L^2(\sigma)} \text{ for all } f$$

if and only if

$$\begin{cases} \|\mathbf{1}_Q H_\sigma \mathbf{1}_Q(x)\|_{L^2(\omega)} \leq C \|\mathbf{1}_Q\|_{L^2(\sigma)} \\ \|\mathbf{1}_Q H_\omega \mathbf{1}_Q(x)\|_{L^2(\sigma)} \leq C \|\mathbf{1}_Q\|_{L^2(\omega)} \end{cases} \quad \text{for all cubes } Q.$$

# Sawyer Testing Conditions for Hilbert transform

Theorem (Lacey-Sawyer-Shen-Uriarte-Tuero; Lacey, 2012)

Assume  $\omega$  and  $\sigma$  don't have common point masses and a stronger  $A_2$  condition holds where the averages are replaced by Poisson averages. Then

$$\|H_\sigma f(x)\|_{L^2(\omega)} \leq C \|f\|_{L^2(\sigma)} \text{ for all } f$$

if and only if

$$\begin{cases} \|\mathbf{1}_Q H_\sigma \mathbf{1}_Q(x)\|_{L^2(\omega)} \leq C \|\mathbf{1}_Q\|_{L^2(\sigma)} & \text{for all cubes } Q. \\ \|\mathbf{1}_Q H_\omega \mathbf{1}_Q(x)\|_{L^2(\sigma)} \leq C \|\mathbf{1}_Q\|_{L^2(\omega)} \end{cases}$$

i.e. morally, to check the norm inequality, it suffices to test the norm inequality over indicators of cubes, i.e. check the “testing condition.”

# Sawyer Testing Conditions for nice CZOs

For general nice CZOs, have a similar statement if the measures are “regular enough.”

# Sawyer Testing Conditions for nice CZOs

For general nice CZOs, have a similar statement if the measures are “regular enough.”

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

# Sawyer Testing Conditions for nice CZOs

For general nice CZOs, have a similar statement if the measures are “regular enough.”

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

Morally, if  $\mu$  is doubling, it means that  $\mu$  can't locally change too much (e.g. no delta masses).

# Sawyer Testing Conditions for nice CZOs

For general nice CZOs, have a similar statement if the measures are “regular enough.”

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

Morally, if  $\mu$  is doubling, it means that  $\mu$  can't locally change too much (e.g. no delta masses).

## Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

*A similar testing theorem holds for all nice CZOs so long as the measures  $\sigma, \omega$  are doubling.*

# Sawyer Testing Conditions for nice CZOs

For general nice CZOs, have a similar statement if the measures are “regular enough.”

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

Morally, if  $\mu$  is doubling, it means that  $\mu$  can't locally change too much (e.g. no delta masses).

## Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

*A similar testing theorem holds for all nice CZOs so long as the measures  $\sigma, \omega$  are doubling.*

## Theorem (Sawyer-Wick, 2022)

*A more technical (but similar in spirit)  $L^p$  testing theorem holds for all nice CZOs so long as the measures  $\sigma, \omega$  are doubling.*

# One-weight vs two-weight problems: $A_2$ vs Testing

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check:

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas  $A_2$  is easy to check, since everything's positive and the averaging integrals are simple.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas  $A_2$  is easy to check, since everything's positive and the averaging integrals are simple. Furthermore,  $A_2$  robust under dilations, translations, and biLipschitz change of variable.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas  $A_2$  is easy to check, since everything's positive and the averaging integrals are simple. Furthermore,  $A_2$  robust under dilations, translations, and biLipschitz change of variable.
- $A_2$  is *stable* under biLipschitz change of variable.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas  $A_2$  is easy to check, since everything's positive and the averaging integrals are simple. Furthermore,  $A_2$  robust under dilations, translations, and biLipschitz change of variable.
- $A_2$  is *stable* under biLipschitz change of variable. So one-weight norm inequalities are too.

## One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas  $A_2$  is easy to check, since everything's positive and the averaging integrals are simple. Furthermore,  $A_2$  robust under dilations, translations, and biLipschitz change of variable.
- $A_2$  is *stable* under biLipschitz change of variable. So one-weight norm inequalities are too. Not the case for two-weight norm inequalities.

# One-weight vs two-weight problems: $A_2$ vs Testing

- In the one-weight case,  $A_2$  holds iff your operator is bounded.
- In the two-weight case,  $A_2$  cannot *characterize* boundedness of your operator, only testing can.
- However, testing can be hard to check: need to delicately estimate cancellation while operator acts on exotic measures.
- Whereas  $A_2$  is easy to check, since everything's positive and the averaging integrals are simple. Furthermore,  $A_2$  robust under dilations, translations, and biLipschitz change of variable.
- $A_2$  is *stable* under biLipschitz change of variable. So one-weight norm inequalities are too. Not the case for two-weight norm inequalities.

## Theorem (Lacey-Sawyer-Uriarte-Tuero, 2012)

$\exists$  measures  $\sigma, \omega$  and a biLipchitz map  $\varphi$  such that the norm inequality for  $H$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the pushforwards  $(\varphi_*\sigma, \varphi_*\omega)$ .

# Two-weight norm inequalities are unstable: a first example

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .


## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

---


## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .




## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .




## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .



## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .



$\sigma$  assigns equal weight to these intervals

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

— — — — — — — —

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .


Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .

— — — — — — — —

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .


Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .


Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



# Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .


Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .




$\omega = \text{sum of delta masses at these zeros of } H\sigma(x)$ .

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .




$\omega = \text{sum of delta masses at these zeros of } H\sigma(x)$ .

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .




$\omega = \text{sum of delta masses at these zeros of } H\sigma(x)$ .

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^p d\omega(x) = 0$ . Then one can show  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



Now, shift the point masses to blow-up points of  $H\sigma(x)$  via a biLipschitz map.


$\omega = \text{sum of delta masses at these zeros of } H\sigma(x)$ .

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^p d\omega(x) = 0$ . Then one can show  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



Now, shift the point masses to blow-up points of  $H\sigma(x)$  via a biLipschitz map.


$\omega = \text{sum of delta masses at these blow-up points of } H\sigma(x)$ .

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^p d\omega(x) = 0$ . Then one can show  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



Now, shift the point masses to blow-up points of  $H\sigma(x)$  via a biLipschitz map.


$\omega = \text{sum of delta masses at these blow-up points of } H\sigma(x)$ .

$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^p d\omega(x) = \infty$ . Then one can show  $H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$

## Two-weight norm inequalities are unstable: a first example

Recall the middle-third Cantor set  $E$  and Cantor measure  $\sigma$ .

Let's graph  $H(\sigma)(x) = \int \frac{d\sigma(y)}{|x-y|}$ .



Now, shift the point masses to blow-up points of  $H\sigma(x)$  via a biLipschitz map.

$\omega = \text{sum of delta masses at these blow-up points of } H\sigma(x)$ .

$$\int |H(\mathbf{1}_{[0,1]}\sigma)(x)|^p d\omega(x) = \infty. \text{ Then one can show } H_\sigma : L^2(\sigma) \rightarrow L^2(\omega)$$

Morally, this is why you need testing: one can only characterize the norm inequality with some biLipschitz unstable condition.

# How far does the biLipschitz instability go?

# How far does the biLipschitz instability go?

- **Devil's Advocate:** biLipschitz instability in previous example feels a little manufactured.

# How far does the biLipschitz instability go?

- **Devil's Advocate:** biLipschitz instability in previous example feels a little manufactured.
- Maybe there's a simple condition which would rule this sort of example out? What if we restricted ourselves to a class of more regular measures?

# How far does the biLipschitz instability go?

- **Devil's Advocate:** biLipschitz instability in previous example feels a little manufactured.
- Maybe there's a simple condition which would rule this sort of example out? What if we restricted ourselves to a class of more regular measures?
- A: Ok we can do that with strong assumptions!

# How far does the biLipschitz instability go?

- **Devil's Advocate:** biLipschitz instability in previous example feels a little manufactured.
- Maybe there's a simple condition which would rule this sort of example out? What if we restricted ourselves to a class of more regular measures?
- A: Ok we can do that with strong assumptions!



# How far does the biLipschitz instability go?

- **Devil's Advocate:** biLipschitz instability in previous example feels a little manufactured.
- Maybe there's a simple condition which would rule this sort of example out? What if we restricted ourselves to a class of more regular measures?
- A: Ok we can do that with strong assumptions!



## Theorem (Hytönen-Lacey, 2012)

Let  $T$  be a nice CZO. Then the norm inequality for  $T_\sigma$  is biLipschitz stable for pairs of  $A_\infty$  measures  $(\sigma, \omega)$ .


# How far does the biLipschitz instability go?

- **Devil's Advocate:** biLipschitz instability in previous example feels a little manufactured.
- Maybe there's a simple condition which would rule this sort of example out? What if we restricted ourselves to a class of more regular measures?
- A: Ok we can do that with strong assumptions!



## Theorem (Hytönen-Lacey, 2012)

Let  $T$  be a nice CZO. Then the norm inequality for  $T_\sigma$  is biLipschitz stable for pairs of  $A_\infty$  measures  $(\sigma, \omega)$ .



# What are $A_\infty$ weights?

# What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

# What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ .

## What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

## What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

# What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

- $A_p$  measures are absolutely continuous.

## What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

- $A_p$  measures are absolutely continuous.
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.

# What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

- $A_p$  measures are absolutely continuous.
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- If  $w \in A_p$ , then it satisfies a Reverse Hölder inequality, i.e.

$$\left( \frac{1}{|Q|} \int_Q w^{1+\epsilon} \right)^{\frac{1}{1+\epsilon}} \leq C \frac{1}{|Q|} \int_Q w.$$

# What are $A_\infty$ weights?

We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

- $A_p$  measures are absolutely continuous.
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- If  $w \in A_p$ , then it satisfies a Reverse Hölder

inequality, i.e.  $\left( \frac{1}{|Q|} \int_Q w^{1+\epsilon} \right)^{\frac{1}{1+\epsilon}} \leq C \frac{1}{|Q|} \int_Q w$ .

- If  $w \in A_p$ , then  $w$  is a doubling weight.

# What are $A_\infty$ weights?

We say  $w \in A_p$  if


$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{ cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

- $A_p$  measures are absolutely continuous.
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- If  $w \in A_p$ , then it satisfies a Reverse Hölder inequality, i.e.

$$\left( \frac{1}{|Q|} \int_Q w^{1+\epsilon} \right)^{\frac{1}{1+\epsilon}} \leq C \frac{1}{|Q|} \int_Q w.$$

- If  $w \in A_p$ , then  $w$  is a doubling weight.



# What are $A_\infty$ weights?

We say  $w \in A_p$  if



$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{Q: \text{cube in } \mathbb{R}} \left( \frac{1}{|Q|} \int_Q w \right) \left( \frac{1}{|Q|} \int_Q w^{-p'/p} \right)^{p/p'} < \infty.$$

Set  $A_\infty \equiv \bigcup_{p>1} A_p$ . This is an incredibly regular class of weights:

- $A_p$  measures are absolutely continuous.
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- If  $w \in A_p$ , then it satisfies a Reverse Hölder inequality, i.e.

$$\left( \frac{1}{|Q|} \int_Q w^{1+\epsilon} \right)^{\frac{1}{1+\epsilon}} \leq C \frac{1}{|Q|} \int_Q w.$$

- If  $w \in A_p$ , then  $w$  is a doubling weight.



# Weakening regularity while maintaining biLipschitz stability?

# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.

# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.



# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.
- Is there a wider, more ubiquitous class of regular measures which are biLipschitz stable, e.g., doubling measures?



# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.
- Is there a wider, more ubiquitous class of regular measures which are biLipschitz stable, e.g., doubling measures?



## Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .\*

# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.
- Is there a wider, more ubiquitous class of regular measures which are biLipschitz stable, e.g., doubling measures?




## Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .\*

\* We proved a broad quantitative version of this statement corresponding to our notion of stability.

# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.
- Is there a wider, more ubiquitous class of regular measures which are biLipschitz stable, e.g., doubling measures?




Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .\*

\* We proved a broad quantitative version of this statement corresponding to our notion of stability.

# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.
- Is there a wider, more ubiquitous class of regular measures which are biLipschitz stable, e.g., doubling measures?




Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .\*

\* We proved a broad quantitative version of this statement corresponding to our notion of stability. The set of biLipschitz stable measures is already a lot smaller than I'd have first thought.

# Weakening regularity while maintaining biLipschitz stability?

- **Devil's Advocate:** Ok fine, for very regular measures, we get biLipschitz stability.
- Is there a wider, more ubiquitous class of regular measures which are biLipschitz stable, e.g., doubling measures?



Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .\*

\* We proved a broad quantitative version of this statement corresponding to our notion of stability. The set of biLipschitz stable measures is already a lot smaller than I'd have first thought. Finding the precise dividing line between biLipschitz stable and unstable measures is an open problem!

# Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .

# Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .

We construct measures that look like sheet metal!




# Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .

We construct measures that look like sheet metal!




Start with  $R_2$ : it points vertically along the level curves of the sheet metal.

# Rotational instability on doubling measures: big picture

Theorem (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

Let  $R_j$  be a Riesz transform. For every  $\epsilon > 0$ , there exists an angle  $\theta \in (0, \epsilon)$  and doubling measures  $\sigma, \omega$  such that the norm inequality for  $R_j$  holds w.r.t.  $(\sigma, \omega)$ , but does not w.r.t. the rotated measures  $(\sigma_\theta, \omega_\theta)$ .

We construct measures that look like sheet metal!



Start with  $R_2$ : it points vertically along the level curves of the sheet metal.  
Rotate  $R_2$  slightly: it sees all the ripples all of a sudden.

# Rotational instability on doubling measures: the key ideas

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.
  - ② Break the resulting weights into their key frequency components, and space them out via a rearrangement/transplantation argument.

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.
  - ② Break the resulting weights into their key frequency components, and space them out via a rearrangement/transplantation argument. This localizes the action of the Hilbert transform on each “piece” of the weights.

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.
  - ② Break the resulting weights into their key frequency components, and space them out via a rearrangement/transplantation argument. This localizes the action of the Hilbert transform on each “piece” of the weights.
  - ③ Use complex analytic properties of the Hilbert transform to get appropriate convergence of Hilbert transform acting on each “piece” of the weights to recover ill-behaved data from dyadic operator.

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.
  - ② Break the resulting weights into their key frequency components, and space them out via a rearrangement/transplantation argument. This localizes the action of the Hilbert transform on each “piece” of the weights.
  - ③ Use complex analytic properties of the Hilbert transform to get appropriate convergence of Hilbert transform acting on each “piece” of the weights to recover ill-behaved data from dyadic operator.
- ② Try to bootstrap from dimension 1 to higher dimensions!

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.
  - ② Break the resulting weights into their key frequency components, and space them out via a rearrangement/transplantation argument. This localizes the action of the Hilbert transform on each “piece” of the weights.
  - ③ Use complex analytic properties of the Hilbert transform to get appropriate convergence of Hilbert transform acting on each “piece” of the weights to recover ill-behaved data from dyadic operator.
- ② Try to bootstrap from dimension 1 to higher dimensions!
  - ① Tensor the weights on  $[0, 1]$  with  $\mathbf{1}_{[0,1]}(x_2)$  to get weights which are constant in one direction, i.e. “sheet metal” weights.

# Rotational instability on doubling measures: the key ideas

- ① First create doubling weights on  $\mathbb{R}$  for which the Hilbert transform  $H$  is badly behaved, following F. Nazarov.
  - ① Construct weights  $\sigma, \omega$  on  $[0, 1] \subset \mathbb{R}$  which are ill-behaved with respect to a discrete *dyadic operator*, which serves as a threshold for bad behavior of CZOs.
  - ② Break the resulting weights into their key frequency components, and space them out via a rearrangement/transplantation argument. This localizes the action of the Hilbert transform on each “piece” of the weights.
  - ③ Use complex analytic properties of the Hilbert transform to get appropriate convergence of Hilbert transform acting on each “piece” of the weights to recover ill-behaved data from dyadic operator.
- ② Try to bootstrap from dimension 1 to higher dimensions!
  - ① Tensor the weights on  $[0, 1]$  with  $\mathbf{1}_{[0,1]}(x_2)$  to get weights which are constant in one direction, i.e. “sheet metal” weights.
  - ② Show these weights are ill-behaved w.r.t. the Riesz transform  $R_1$  by reducing to  $H$ , but well-behaved w.r.t. the Riesz transform  $R_2$ .

# Step 1: Dyadic model construction on $[0, 1] \subset \mathbb{R}$

## Step 1: Dyadic model construction on $[0, 1] \subset \mathbb{R}$

Define the dyadic grid  $\mathcal{D} = \{[j2^{-k}, (j+1)2^{-k})\}_{j,k \in \mathbb{Z}}$   
and consider the dyadic square function


$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

# Step 1: Dyadic model construction on $[0, 1] \subset \mathbb{R}$

Define the dyadic grid  $\mathcal{D} = \{[j2^{-k}, (j+1)2^{-k})\}_{j,k \in \mathbb{Z}}$  and consider the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

We first construct weights to get its norm to blow-up.




## Step 1: Dyadic model construction on $[0, 1] \subset \mathbb{R}$

Define the dyadic grid  $\mathcal{D} = \{[j2^{-k}, (j+1)2^{-k})\}_{j,k \in \mathbb{Z}}$  and consider the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

We first construct weights to get its norm to blow-up.

Why? Because the Hilbert transform is an average of dyadic operators!




## Step 1: Dyadic model construction on $[0, 1] \subset \mathbb{R}$

Define the dyadic grid  $\mathcal{D} = \{[j2^{-k}, (j+1)2^{-k})\}_{j,k \in \mathbb{Z}}$  and consider the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

We first construct weights to get its norm to blow-up.

Why? Because the Hilbert transform is an average of dyadic operators! A Bellman function argument by F. Nazarov shows that the norm of  $S$  blows up for a pair of dyadically doubling weights  $(\sigma, \omega)$  on  $[0, 1]$  which satisfy the  $A_2$  condition.




## Step 1: Dyadic model construction on $[0, 1] \subset \mathbb{R}$

Define the dyadic grid  $\mathcal{D} = \{[j2^{-k}, (j+1)2^{-k})\}_{j,k \in \mathbb{Z}}$  and consider the dyadic square function

$$Sf(x) \equiv \sqrt{\sum_{J \in \mathcal{D}} (E_{J_-} f - E_{J_+} f)^2 |J| \mathbf{1}_J(x)}.$$

We first construct weights to get its norm to blow-up.



Why? Because the Hilbert transform is an average of dyadic operators! A Bellman function argument by F. Nazarov shows that the norm of  $S$  blows up for a pair of dyadically doubling weights  $(\sigma, \omega)$  on  $[0, 1]$  which satisfy the  $A_2$  condition.

(Note: virtually only way I know to get doubling weight counter-examples is via this method.)


Step 2: rearrange our weights by transplanting averages!

## Step 2: rearrange our weights by transplanting averages!

Colors = average of weight on interval

Initial weight  $W_0$

Transplanted weight  $W_1$




## Step 2: rearrange our weights by transplanting averages!

Colors = average of weight on interval

Initial weight  $W_0$


Transplanted weight  $W_1$



Why do this?

## Step 2: rearrange our weights by transplanting averages!

Colors = average of weight on interval




Why do this? The weight  $W_0$  on the left can be written as

$$E_{[0,1]} W_0 + \sum_{I \in \mathcal{D}} c_I h_I, \text{ where } h_I = \begin{cases} -1 & \text{on } I_- \\ 1 & \text{on } I_+ \end{cases}$$

is a Haar wavelet.

## Step 2: rearrange our weights by transplanting averages!

Colors = average of weight on interval



Why do this? The weight  $W_0$  on the left can be written as


$$E_{[0,1]} W_0 + \sum_{I \in \mathcal{D}} c_I h_I, \text{ where } h_I = \begin{cases} -1 & \text{on } I_- \\ 1 & \text{on } I_+ \end{cases}$$

is a Haar wavelet.

By re-arranging the weight  $W_0$  into  $W_1$ , we preserve the basic data of  $W_0$  (has same averages on cubes),

## Step 2: rearrange our weights by transplanting averages!

Colors = average of weight on interval



Why do this? The weight  $W_0$  on the left can be written as

$$E_{[0,1]} W_0 + \sum_{I \in \mathcal{D}} c_I h_I, \text{ where } h_I = \begin{cases} -1 & \text{on } I_- \\ 1 & \text{on } I_+ \end{cases}$$

is a Haar wavelet.


By re-arranging the weight  $W_0$  into  $W_1$ , we preserve the basic data of  $W_0$  (has same averages on cubes), but shift all the Haar wavelets to frequencies of drastically different scales, ensuring they are well-localized and don't "interfere" with each other.

# The oscillatory functions

# The oscillatory functions

By relocating Haar data and increasing its frequency, to get  $W_1 = E_{[0,1]} W_1 + \sum_{I \in S} c_I s_k^I$ ,

where  $s_k^I$  is an alternating function of frequency  $2^k$  on  $I$ , and  $S$  is a set of intervals at sparse scales.




# The oscillatory functions

By relocating Haar data and increasing its frequency, to get  $W_1 = E_{[0,1]} W_1 + \sum_{I \in S} c_I s_k^I$ ,

where  $s_k^I$  is an alternating function of frequency  $2^k$  on  $I$ , and  $S$  is a set of intervals at sparse scales.

We want to show  $\int_{[0,1]} |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  blows-up for our weights  $\sigma, \omega$  constructed.



# The oscillatory functions


By relocating Haar data and increasing its frequency, to get  $W_1 = E_{[0,1]} W_1 + \sum_{I \in S} c_I s_k^I$ ,

where  $s_k^I$  is an alternating function of frequency  $2^k$  on  $I$ , and  $S$  is a set of intervals at sparse scales.

We want to show  $\int_{[0,1]} |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  blows-up for our weights  $\sigma, \omega$  constructed. Foiling yields this looks like

$$\sum_{I_1, I_2, I_3 \in S} c_{I_1 I_2 I_3} \int \left( H s_{k(I_1)}^{I_1} H s_{k(I_2)}^{I_2} \right) s_{k(I_3)}^{I_3} + \text{lower order terms ,}$$

where  $c_{I_1, I_2, I_3}$  represents data from the initial weights that will hopefully make the sum diverge.



# The oscillatory functions

By relocating Haar data and increasing its frequency, to get  $W_1 = E_{[0,1]} W_1 + \sum_{I \in S} c_I s_k^I$ ,


where  $s_k^I$  is an alternating function of frequency  $2^k$  on  $I$ , and  $S$  is a set of intervals at sparse scales.

We want to show  $\int_{[0,1]} |H(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  blows-up for our weights  $\sigma, \omega$  constructed. Foiling yields this looks like

$$\sum_{I_1, I_2, I_3 \in S} c_{I_1 I_2 I_3} \int \left( H s_{k(I_1)}^{I_1} H s_{k(I_2)}^{I_2} \right) s_{k(I_3)}^{I_3} + \text{lower order terms ,}$$

where  $c_{I_1, I_2, I_3}$  represents data from the initial weights that will hopefully make the sum diverge.

One has to then estimate expressions like  $\int (H s_k^I) s_k^I$ ,  $\int (H s_k^I)^2$ , etc ... as  $k \rightarrow \infty$ .



# One such estimate using complex analysis with $H$

## One such estimate using complex analysis with $H$

One such estimate boils down to showing  $(Hs_k^l)^2 \rightarrow \mathbf{1}_I$ , weakly, which we only know how to do using complex analysis:

## One such estimate using complex analysis with $H$

One such estimate boils down to showing  $(Hs_k^I)^2 \rightarrow \mathbf{1}_I$  weakly, which we only know how to do using complex analysis:

- $s_k^I \rightarrow 0$  weakly as  $k \rightarrow \infty$ .

## One such estimate using complex analysis with $H$

One such estimate boils down to showing  $(Hs_k^I)^2 \rightarrow \mathbf{1}_I$  weakly, which we only know how to do using complex analysis:

- $s_k^I \rightarrow 0$  weakly as  $k \rightarrow \infty$ .
- Then  $s_k^I + iHs_k^I \rightarrow 0$  weakly. And so does  $(s_k^I + iHs_k^I)^2 \rightarrow 0$ .

## One such estimate using complex analysis with $H$

One such estimate boils down to showing  $(Hs_k^I)^2 \rightarrow \mathbf{1}_I$  weakly, which we only know how to do using complex analysis:

- $s_k^I \rightarrow 0$  weakly as  $k \rightarrow \infty$ .
- Then  $s_k^I + iHs_k^I \rightarrow 0$  weakly. And so does  $(s_k^I + iHs_k^I)^2 \rightarrow 0$ .
- This last function has real part  $(s_k^I)^2 - (Hs_k^I)^2 = \mathbf{1}_I - (Hs_k^I)^2$ .

## One such estimate using complex analysis with $H$

One such estimate boils down to showing  $(Hs_k^I)^2 \rightarrow \mathbf{1}_I$  weakly, which we only know how to do using complex analysis:

- $s_k^I \rightarrow 0$  weakly as  $k \rightarrow \infty$ .
- Then  $s_k^I + iHs_k^I \rightarrow 0$  weakly. And so does  $(s_k^I + iHs_k^I)^2 \rightarrow 0$ .
- This last function has real part  $(s_k^I)^2 - (Hs_k^I)^2 = \mathbf{1}_I - (Hs_k^I)^2$ .
- Since the real part is going to 0, then  $(Hs_k^I)^2 \rightarrow \mathbf{1}_I$ .

# Tensoring weights, and the needed reductions

## Tensoring weights, and the needed reductions

Now tensor 1D weights with  $\mathbf{1}_{[0,1]}(x_2)$  to get “sheet metal” weights  $\sigma, \omega$ .

## Tensoring weights, and the needed reductions

Now tensor 1D weights with  $\mathbf{1}_{[0,1]}(x_2)$  to get “sheet metal” weights  $\sigma, \omega$ .

Decompose  $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  into base frequency elements.

## Tensoring weights, and the needed reductions

Now tensor 1D weights with  $\mathbf{1}_{[0,1]}(x_2)$  to get “sheet metal” weights  $\sigma, \omega$ .

Decompose  $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  into base frequency elements.

Must now consider  $R_1$  acting on the horizontal oscillating functions

$$s_k^Q(x) \equiv \left( s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right)(x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \quad \text{where cube } Q \equiv Q_1 \times Q_2.$$

## Tensoring weights, and the needed reductions

Now tensor 1D weights with  $\mathbf{1}_{[0,1]}(x_2)$  to get “sheet metal” weights  $\sigma, \omega$ .

Decompose  $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  into base frequency elements.

Must now consider  $R_1$  acting on the horizontal oscillating functions

$$s_k^Q(x) \equiv \left( s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right)(x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \quad \text{where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ①  $R_1 \left( s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow \left( Hs_k^{Q_1} \right) \otimes \mathbf{1}_{Q_2}$  weakly.
- ②  $R_2 \left( s_k^{Q_1} \otimes \mathbf{1}_{Q_2} \right) \rightarrow 0$  in  $L^p(\mathbb{R}^2)$ .

## Tensoring weights, and the needed reductions

Now tensor 1D weights with  $\mathbf{1}_{[0,1]}(x_2)$  to get “sheet metal” weights  $\sigma, \omega$ .

Decompose  $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  into base frequency elements.

Must now consider  $R_1$  acting on the horizontal oscillating functions

$$s_k^Q(x) \equiv (s_k^{Q_1} \otimes \mathbf{1}_{Q_2})(x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \quad \text{where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ①  $R_1(s_k^{Q_1} \otimes \mathbf{1}_{Q_2}) \rightarrow (Hs_k^{Q_1}) \otimes \mathbf{1}_{Q_2}$  weakly.
- ②  $R_2(s_k^{Q_1} \otimes \mathbf{1}_{Q_2}) \rightarrow 0$  in  $L^p(\mathbb{R}^2)$ .

By the first property, we can apply the same techniques as for the Hilbert transform and get norm for  $R_1$  to blow-up.

## Tensoring weights, and the needed reductions

Now tensor 1D weights with  $\mathbf{1}_{[0,1]}(x_2)$  to get “sheet metal” weights  $\sigma, \omega$ .

Decompose  $\int_{[0,1]} |R_1(\mathbf{1}_{[0,1]}\sigma)(x)|^2 d\omega(x)$  into base frequency elements.

Must now consider  $R_1$  acting on the horizontal oscillating functions

$$s_k^Q(x) \equiv (s_k^{Q_1} \otimes \mathbf{1}_{Q_2})(x) = s_k^{Q_1}(x_1) \mathbf{1}_{Q_2}(x_2), \quad \text{where cube } Q \equiv Q_1 \times Q_2.$$

Proposition (A.-Luna-Garcia-Sawyer-Uriarte-Tuero, 2022)

- ①  $R_1(s_k^{Q_1} \otimes \mathbf{1}_{Q_2}) \rightarrow (Hs_k^{Q_1}) \otimes \mathbf{1}_{Q_2}$  weakly.
- ②  $R_2(s_k^{Q_1} \otimes \mathbf{1}_{Q_2}) \rightarrow 0$  in  $L^p(\mathbb{R}^2)$ .

By the first property, we can apply the same techniques as for the Hilbert transform and get norm for  $R_1$  to blow-up.

If we rotate our measures 90 degrees, our oscillatory functions are killed off by  $R_2$ , and so the norm of  $R_2$  is under control!

*Thank you for Listening!*