

# The Steklov Problem for OPUC and Krein Systems

Michel Alexis

McMaster University

JMM, January 6 2023


# Orthogonal Polynomials on the Unit Circle (OPUC)

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .


# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



# Orthogonal Polynomials on the Unit Circle (OPUC)


Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




$$1, z, z^2, z^3, \dots$$

Can write  $z = e^{i\theta}$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




Can write  $z = e^{i\theta}$ .

$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .

$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$




$$w(\theta) \frac{d\theta}{2\pi}$$



# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .


$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

↓ Gram-Schmidt

$$\downarrow \\ w(\theta) \frac{d\theta}{2\pi}$$

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




Can write  $z = e^{i\theta}$ .

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




Can write  $z = e^{i\theta}$ .

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

- $\varphi_n(z)$  is orthonormal polynomial of degree  $n$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

- $\varphi_n(z)$  is orthonormal polynomial of degree  $n$ .
- $\{\varphi_n(z)\}$  are called Orthogonal Polynomials on the Unit Circle.

# How do OPUC behave?

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ .



# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ . Makes sense since  $\int |\varphi_n|^2 w = 1$ , and  $\varphi_n$  oscillates more and more rapidly.

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ . Makes sense since  $\int |\varphi_n|^2 w = 1$ , and  $\varphi_n$  oscillates more and more rapidly.


We expect this behavior pointwise as well.

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ . Makes sense since  $\int |\varphi_n|^2 w = 1$ , and  $\varphi_n$  oscillates more and more rapidly.

We expect this behavior pointwise as well.





# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ . Makes sense since  $\int |\varphi_n|^2 w = 1$ , and  $\varphi_n$  oscillates more and more rapidly.

We expect this behavior pointwise as well.



$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)


*If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .*

$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.




$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.



But  $\|\varphi_n\|_{L^2(w)} = 1$  by definition.


Maybe we're asking for too much regularity?

$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.



But  $\|\varphi_n\|_{L^2(w)} = 1$  by definition.

Maybe we're asking for too much regularity?

## Problem (Steklov problem)

Does there exist  $p > 2$  such that  $\sup_n \|\varphi_n\|_{L^p(w)} < \infty$ ?




$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.



But  $\|\varphi_n\|_{L^2(w)} = 1$  by definition.

Maybe we're asking for too much regularity?

## Problem (Steklov problem)

Does there exist  $p > 2$  such that  $\sup_n \|\varphi_n\|_{L^p(w)} < \infty$ ?

**Remark:** If  $\int_{\mathbb{T}} \log w > -\infty$ , then  $|\varphi_n(z)| \sim |\Phi_n(z)|$ , where  $\Phi_n(z)$  are the monic orthogonal polynomials of degree  $n$ .

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq \Lambda$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq \lambda 1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

Proof:

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq \chi$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq \chi 1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n = z^n + \mathcal{P}_{[0,n-1]} \Phi_n & (\text{monic}) \end{cases}$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq \chi 1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]} \Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]} w \Phi_n \quad (\text{orthogonal}) \end{cases}$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n & (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n & (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$



Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq \chi 1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n & (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n & (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

$$\|\Phi_n\|_p \leq \|(I - \mathcal{P}_{[0,n-1]}(1-w))^{-1}\|_{p,p} \|z^n\|_p$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

$$\|\Phi_n\|_p \leq \|(I - \mathcal{P}_{[0,n-1]}(1-w))^{-1}\|_{p,p} \|z^n\|_p \leq \left\| \sum_{k=0}^{\infty} (\mathcal{P}_{[0,n-1]}(1-w))^k \right\|_{p,p}$$

Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2


Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,





Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,






Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,






Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,



Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2




- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,



- Given  $\|\mathcal{P}_{[0,n]}\|_{2,2} = 1$ , one can show  $\|\mathcal{P}_{[0,n]}\|_{p,p} \leq 1 + O(|p - 2|)$  uniformly in  $n$ .

Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,



- Given  $\|\mathcal{P}_{[0,n]}\|_{2,2} = 1$ , one can show  $\|\mathcal{P}_{[0,n]}\|_{p,p} \leq 1 + O(|p - 2|)$  uniformly in  $n$ .
- Choose  $p$  close enough to 2 so that  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$

□

# $A_p$ weights

- We say  $w \in A_2$  if

$$[w]_{A_2} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-1} \right) < \infty .$$

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts about  $A_p$  weights:



# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts about  $A_p$  weights:
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts about  $A_p$  weights:
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- $[w]_{A_p} \geq 1$ ; the closer  $[w]_{A_p}$  is to 1 the “flatter”  $w$  is.

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts about  $A_p$  weights:
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- $[w]_{A_p} \geq 1$ ; the closer  $[w]_{A_p}$  is to 1 the “flatter”  $w$  is.
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$


# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts about  $A_p$  weights:

- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- $[w]_{A_p} \geq 1$ ; the closer  $[w]_{A_p}$  is to 1 the “flatter”  $w$  is.
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$



# $A_p$ weights


- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts about  $A_p$  weights:

- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- $[w]_{A_p} \geq 1$ ; the closer  $[w]_{A_p}$  is to 1 the “flatter”  $w$  is.
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$

$A_p$  plays nicely with the Hilbert transform  $\mathcal{H}$ .



# $A_p$ weights


- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty.$$

- Facts about  $A_p$  weights:

- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- $[w]_{A_p} \geq 1$ ; the closer  $[w]_{A_p}$  is to 1 the “flatter”  $w$  is.
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$

$A_p$  plays nicely with the Hilbert transform  $\mathcal{H}$ .



## Theorem (Hunt-Muckenhoupt-Wheeden)

If  $w \in A_p$ , then  $\|w^{1/p} \mathcal{H} w^{-1/p}\|_{p,p} = \|\mathcal{H}\|_{L^p(w) \rightarrow L^p(w)} < \infty$ .

# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

*If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p = p([w]_{A_2}) > 2$ .*

# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p = p([w]_{A_2}) > 2$ .

## Proof.

- Some algebra similar to before yields

$$(I - Q_{w,p})w^{1/p}\Phi_n = w^{1/p}z^n,$$

where  $Q_{w,p} = w^{1/p}\mathcal{P}_{[0,n-1]}w^{-1/p} - w^{-1/p'}\mathcal{P}_{[0,n-1]}w^{1/p'}$ .



# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p = p([w]_{A_2}) > 2$ .

## Proof.

- Some algebra similar to before yields

$$(I - Q_{w,p})w^{1/p}\Phi_n = w^{1/p}Z^n,$$

where  $Q_{w,p} = w^{1/p}\mathcal{P}_{[0,n-1]}w^{-1/p} - w^{-1/p'}\mathcal{P}_{[0,n-1]}w^{1/p'}$ .

- We can then invert  $I - Q_{w,p}$  when  $|\frac{1}{2} - \frac{1}{p}| < \epsilon([w]_{A_2})$  using spectral theory and analytic interpolation.



# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$   
OPUC

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

# Krein Systems, the continuous analogue of OPUC

$$\begin{array}{l} (d\mu, \mathbb{T}) \\ \text{OPUC} \\ \Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k \\ \left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2 \end{array}$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

Krein Systems

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$



# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

$$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$$

# Krein Systems, the continuous analogue of OPUC

$$(d\mu, \mathbb{T})$$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$$

$$(d\sigma, \mathbb{R})$$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

$$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$ OPUC	$(d\sigma, \mathbb{R})$ Krein Systems
$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$	$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$
$\left\  \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\ _{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty}  a_j ^2$	$\left\  \int_0^{\infty} f(s) P(s, \lambda) ds \right\ _{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty}  f(s) ^2$
$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$	$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

**Notation:**

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$ OPUC	$(d\sigma, \mathbb{R})$ Krein Systems
$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$	$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$
$\left\  \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\ _{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty}  a_j ^2$	$\left\  \int_0^{\infty} f(s) P(s, \lambda) ds \right\ _{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty}  f(s) ^2$
$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$	$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

## Notation:

- $r > 0$  is an index (analogous to  $n$  in  $\varphi_n$ ).

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$ OPUC	$(d\sigma, \mathbb{R})$ Krein Systems
$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$	$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$
$\left\  \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\ _{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty}  a_j ^2$	$\left\  \int_0^{\infty} f(s) P(s, \lambda) ds \right\ _{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty}  f(s) ^2$
$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$	$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

## Notation:

- $r > 0$  is an index (analogous to  $n$  in  $\varphi_n$ ).
- $\lambda$  is the variable denoting points in the space  $\mathbb{R}$ .

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$ OPUC	$(d\sigma, \mathbb{R})$ Krein Systems
$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$	$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$
$\left\  \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\ _{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty}  a_j ^2$	$\left\  \int_0^{\infty} f(s) P(s, \lambda) ds \right\ _{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty}  f(s) ^2$
$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$	$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

## Notation:

- $r > 0$  is an index (analogous to  $n$  in  $\varphi_n$ ).
- $\lambda$  is the variable denoting points in the space  $\mathbb{R}$ .

We will be interested in the case when  $d\sigma = w(\lambda) \frac{d\lambda}{2\pi}$ , where  $w - 1 \in L^1(\mathbb{R}) + L^2(\mathbb{R})$  to ensure  $P(r, \lambda)$  exists.



# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda)\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ .

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

*Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which*

$$\sup_{r \geq 0} \|P(r, \lambda)\|_{L^p(w)} < \infty$$

*whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ .*

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

*Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which*

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

*whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ .*

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $\max\{2 - \epsilon, p_0\} \leq p \leq 2 + \epsilon$ .

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $\max\{2 - \epsilon, p_0\} \leq p \leq 2 + \epsilon$ .

The proof is similar to the previous one, using the useful explicit identity

$$(I - Q_{w,p})X_p = w^{-1/p'} \mathcal{P}_{[0,r]} w^{1/p'} (w^{1/p} - w^{-1/p'}) e^{i\lambda r},$$

where  $X_p \stackrel{\text{def}}{=} w^{1/p} (P(r, \lambda) - e^{i\lambda r})$ .

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $\max\{2 - \epsilon, p_0\} \leq p \leq 2 + \epsilon$ .

The proof is similar to the previous one, using the useful explicit identity

$$X_p = (I - Q_{w,p})^{-1} w^{-1/p'} \mathcal{P}_{[0,r]} w^{1/p'} (w^{1/p} - w^{-1/p'}) e^{i\lambda r},$$

where  $X_p \stackrel{\text{def}}{=} w^{1/p} (P(r, \lambda) - e^{i\lambda r})$ .

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $\max\{2 - \epsilon, p_0\} \leq p \leq 2 + \epsilon$ .

The proof is similar to the previous one, using the useful explicit identity

$$X_p = (I - Q_{w,p})^{-1} w^{-1/p'} \mathcal{P}_{[0,r]}(w - 1) e^{i\lambda r},$$

where  $X_p \stackrel{\text{def}}{=} w^{1/p} (P(r, \lambda) - e^{i\lambda r})$ .

# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $\max\{2 - \epsilon, p_0\} \leq p \leq 2 + \epsilon$ .

The proof is similar to the previous one, using the **useful** explicit identity

$$X_p = (I - Q_{w,p})^{-1} w^{-1/p'} \mathcal{P}_{[0,r]}(w - 1) e^{i\lambda r},$$

where  $X_p \stackrel{\text{def}}{=} w^{1/p} (P(r, \lambda) - e^{i\lambda r})$ .



# Steklov for Krein systems and an explicit formula

## Theorem (A. '21)

Suppose  $w - 1 \in L^{p_0}(\mathbb{R})$  for  $1 < p_0 \leq 2$  and  $w \in A_2(\mathbb{R})$ . Then there exists  $\epsilon = \epsilon([w]_{A_2}) > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $\max\{2 - \epsilon, p_0\} \leq p \leq 2 + \epsilon$ .

The proof is similar to the previous one, using the **useful** explicit identity

$$X_p = (I - Q_{w,p})^{-1} w^{-1/p'} \mathcal{P}_{[0,r]}(w - 1) e^{i\lambda r},$$

where  $X_p \stackrel{\text{def}}{=} w^{1/p} (P(r, \lambda) - e^{i\lambda r})$ .

## Corollary

Above theorem is sharp: one cannot take  $p < p_0$ .

*Thank you for Listening!*