

# The Steklov Problem for OPUC and Krein Systems

Michel Alexis

UW-Madison

Defense, May 21st 2021


# Orthogonal Polynomials on the Unit Circle (OPUC)

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .


# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



# Orthogonal Polynomials on the Unit Circle (OPUC)


Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




$$1, z, z^2, z^3, \dots$$

Can write  $z = e^{i\theta}$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

Can write  $z = e^{i\theta}$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .

$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$




$$w(\theta) \frac{d\theta}{2\pi}$$



# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .


$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

↓ Gram-Schmidt

$$\downarrow \\ w(\theta) \frac{d\theta}{2\pi}$$

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .


$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

↓ Gram-Schmidt

$$\varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots \quad \perp \text{ w.r.t. } w(\theta) \frac{d\theta}{2\pi}$$

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .




Can write  $z = e^{i\theta}$ .

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

- $\varphi_n(z)$  is orthonormal polynomial of degree  $n$ .

# Orthogonal Polynomials on the Unit Circle (OPUC)

Setting:  $\mathbb{T} \subset \mathbb{C}$ .



Can write  $z = e^{i\theta}$ .

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

- $\varphi_n(z)$  is orthonormal polynomial of degree  $n$ .
- $\{\varphi_n(z)\}$  are called Orthogonal Polynomials on the Unit Circle.

# How do OPUC behave?

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ .



# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ .


We expect this behavior pointwise as well.

# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ .

We expect this behavior pointwise as well.





# How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$  in the weak-\* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$  in  $L^1$  and weak-\* sense if  $\int_{\mathbb{T}} \log w > -\infty$ .

Trend seems to be  $|\varphi_n| \rightarrow w^{-1/2}$ .

We expect this behavior pointwise as well.



$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)


*If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .*

$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.




$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.



But  $\|\varphi_n\|_{L^2(w)} = 1$  by definition.


Maybe we're asking for too much regularity?

$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.



But  $\|\varphi_n\|_{L^2(w)} = 1$  by definition.

Maybe we're asking for too much regularity?

## Problem (Steklov problem)


Does there exist  $p > 2$  such that  $\sup_n \|\varphi_n\|_{L^p(w)} < \infty$ ?

$$|\varphi_n| \rightarrow w^{-1/2}$$

## Conjecture (Steklov's Conjecture)

If  $w \geq \delta > 0$ , then  $\{\varphi_n\}$  are bounded above, i.e.  $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$ .

**False!** Can create weights  $w$  which oscillate rapidly, forcing  $\{\varphi_n\}$  to blow-up.



But  $\|\varphi_n\|_{L^2(w)} = 1$  by definition.

Maybe we're asking for too much regularity?

## Problem (Steklov problem)

Does there exist  $p > 2$  such that  $\sup_n \|\varphi_n\|_{L^p(w)} < \infty$ ?

**Remark:** If  $\int_{\mathbb{T}} \log w > -\infty$ , then  $|\varphi_n(z)| \sim |\Phi_n(z)|$ , where  $\Phi_n(z)$  are the monic orthogonal polynomials of degree  $n$ .



Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq \Lambda$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq \lambda 1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

Proof:

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq \chi 1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}.$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

*Suppose  $\epsilon \leq w \leq X1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .*

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n = z^n + \mathcal{P}_{[0,n-1]} \Phi_n & (\text{monic}) \end{cases}$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq \chi$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n & (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n & (\text{orthogonal}) \end{cases}$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n & (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n & (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .



Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1 + \epsilon)(1 - \epsilon) \leq 1 - \epsilon^2 < 1$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

$$\|\Phi_n\|_p \leq \|(I - \mathcal{P}_{[0,n-1]}(1-w))^{-1}\|_{p,p} \|z^n\|_p$$

Does there exist  $p > 2$  such that  $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$ ?

### Theorem (Nazarov, 2016)

Suppose  $\epsilon \leq w \leq K1$ . Then there exists  $p > 2$  for which  $\sup_n \|\Phi_n\|_{L^p} < \infty$ .

**Proof:**  $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$ .

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather  $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$ .

If we can show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2, then done!

- Indeed,  $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

$$\|\Phi_n\|_p \leq \|(I - \mathcal{P}_{[0,n-1]}(1-w))^{-1}\|_{p,p} \|z^n\|_p \leq \left\| \sum_{k=0}^{\infty} (\mathcal{P}_{[0,n-1]}(1-w))^k \right\|_{p,p}$$


Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,

Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2



- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,








Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,






Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,



Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2




- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,



- Given  $\|\mathcal{P}_{[0,n]}\|_{2,2} = 1$ , one can show  $\|\mathcal{P}_{[0,n]}\|_{p,p} \leq 1 + O(|p - 2|)$  uniformly in  $n$ .

Show  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$  for  $p$  close to 2

- Have uniform control of  $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$  since  $\mathcal{P}_{[0,n-1]}$  is a linear combination of Hilbert transforms,



- Given  $\|\mathcal{P}_{[0,n]}\|_{2,2} = 1$ , one can show  $\|\mathcal{P}_{[0,n]}\|_{p,p} \leq 1 + O(|p - 2|)$  uniformly in  $n$ .
- Choose  $p$  close enough to 2 so that  $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$

□

# $A_p$ weights

- We say  $w \in A_2$  if

$$[w]_{A_2} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-1} \right) < \infty .$$

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts/notions about  $A_p$ :

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts/notions about  $A_p$ :
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.



# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts/notions about  $A_p$ :
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- if  $w \in A_p$  then  $\log w \in \text{BMO}$ .

# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty .$$

- Facts/notions about  $A_p$ :
- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- if  $w \in A_p$  then  $\log w \in \text{BMO}$ .
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$


# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty.$$

- Facts/notions about  $A_p$ :

- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- if  $w \in A_p$  then  $\log w \in \text{BMO}$ .
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$




# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty.$$

- Facts/notions about  $A_p$ :

- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- if  $w \in A_p$  then  $\log w \in \text{BMO}$ .
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$



## Theorem (Hunt-Muckenhoupt-Wheeden)

If  $w \in A_p$ , then  $\|w^{1/p} \mathcal{H} w^{-1/p}\|_{p,p} = \|\mathcal{H}\|_{L^p(w) \rightarrow L^p(w)} < \infty$ .


# $A_p$ weights

- We say  $w \in A_p$  if

$$[w]_{A_p} \stackrel{\text{def}}{=} \sup_{I: \text{ arc in } \mathbb{T}} \left( \frac{1}{|I|} \int_I w \right) \left( \frac{1}{|I|} \int_I w^{-p'/p} \right)^{p/p'} < \infty.$$

- Facts/notions about  $A_p$ :

- $w \in A_p$  means  $w$  cannot be too singular/oscillatory.
- if  $w \in A_p$  then  $\log w \in \text{BMO}$ .
- Power weights  $|\theta|^\alpha \in A_p$  if  $-1 < \alpha < p - 1$



## Theorem (Hunt-Muckenhoupt-Wheeden)

If  $w \in A_p$ , then  $\|w^{1/p} \mathcal{H} w^{-1/p}\|_{p,p} = \|\mathcal{H}\|_{L^p(w) \rightarrow L^p(w)} < \infty$ .

$A_p$  plays nicely with the Hilbert transform  $\mathcal{H}$ , so makes sense to adapt previous proof to  $A_p$  weights.

# The Steklov problem for $A_2$ weights

Theorem (A.-Aptekarev-Denisov, '20)

*If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .*

# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .

## Proof.

$$\bullet \begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]} & \Phi_n \text{ (monic)} \\ 0 &= \mathcal{P}_{[0,n-1]} w & \Phi_n \text{ (orthogonal)} \end{cases}$$



# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .

## Proof.

$$\bullet \begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0, n-1]} & \Phi_n \text{ (monic)} \\ 0 &= \mathcal{P}_{[0, n-1]} w & \Phi_n \text{ (orthogonal)} \end{cases}$$





# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .

## Proof.

$$\bullet \begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} w^{1/p} \Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]} w \Phi_n \quad (\text{orthogonal}) \end{cases}$$



# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .

## Proof.

$$\bullet \begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0, n-1]} w^{-1/p} w^{1/p} \Phi_n \quad (\text{monic}) \\ 0 &= w^{-1/p'} \mathcal{P}_{[0, n-1]} w^{1/p'} w^{1/p} \Phi_n \quad (\text{orthogonal}) \end{cases}$$



# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p}\Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .

## Proof.

- $$\begin{cases} w^{1/p}\Phi_n &= w^{1/p}z^n + w^{1/p}\mathcal{P}_{[0,n-1]}w^{-1/p}w^{1/p}\Phi_n \quad (\text{monic}) \\ 0 &= w^{-1/p'}\mathcal{P}_{[0,n-1]}w^{1/p'}w^{1/p}\Phi_n \quad (\text{orthogonal}) \end{cases}$$
- Subtract bottom from top and re-arrange to get

$$(I - Q_{w,p})w^{1/p}\Phi_n = w^{1/p}z^n,$$

$$\text{where } Q_{w,p} = w^{1/p}\mathcal{P}_{[0,n-1]}w^{-1/p} - w^{-1/p'}\mathcal{P}_{[0,n-1]}w^{1/p'}.$$



# The Steklov problem for $A_2$ weights

## Theorem (A.-Aptekarev-Denisov, '20)

If  $w \in A_2$ , then  $\sup_n \|w^{1/p}\Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$  for some  $p > 2$ .

## Proof.

- $$\begin{cases} w^{1/p}\Phi_n &= w^{1/p}z^n + w^{1/p}\mathcal{P}_{[0,n-1]}w^{-1/p}w^{1/p}\Phi_n \quad (\text{monic}) \\ 0 &= w^{-1/p'}\mathcal{P}_{[0,n-1]}w^{1/p'}w^{1/p}\Phi_n \quad (\text{orthogonal}) \end{cases}$$
- Subtract bottom from top and re-arrange to get

$$(I - Q_{w,p})w^{1/p}\Phi_n = w^{1/p}z^n,$$

where  $Q_{w,p} = w^{1/p}\mathcal{P}_{[0,n-1]}w^{-1/p} - w^{-1/p'}\mathcal{P}_{[0,n-1]}w^{1/p'}$ .

- And as it turns out we can invert  $I - Q_{w,p}$  for some  $p$  near 2 using spectral theory and analytic interpolation. (Deferred till end of presentation, if time permits.)



# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$   
OPUC

# Krein Systems, the continuous analogue of OPUC

$$\begin{array}{l} (d\mu, \mathbb{T}) \\ \text{OPUC} \\ \Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k \end{array}$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$



# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

Krein Systems

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$

OPUC

$$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$$

$$\left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2$$

$$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$$

$(d\sigma, \mathbb{R})$

Krein Systems

$$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$$

$$\left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2$$

$$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$$

# Krein Systems, the continuous analogue of OPUC

$$\begin{array}{l} (d\mu, \mathbb{T}) \\ \text{OPUC} \\ \Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k \\ \left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2 \\ d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n \end{array}$$

$$\begin{array}{l} (d\sigma, \mathbb{R}) \\ \text{Krein Systems} \\ P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds \\ \left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2 d\sigma \\ d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r} \end{array}$$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

# Krein Systems, the continuous analogue of OPUC

$$\begin{array}{c} (d\mu, \mathbb{T}) \\ \text{OPUC} \\ \Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k \\ \left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2 \\ d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n \end{array}$$

$$\begin{array}{c} (d\sigma, \mathbb{R}) \\ \text{Krein Systems} \\ P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds \\ \left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2 d\sigma \\ d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r} \end{array}$$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

**Notation:**



# Krein Systems, the continuous analogue of OPUC

$$\begin{array}{c} (d\mu, \mathbb{T}) \\ \text{OPUC} \\ \Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k \\ \left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2 \\ d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n \end{array}$$

$$\begin{array}{c} (d\sigma, \mathbb{R}) \\ \text{Krein Systems} \\ P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds \\ \left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2 d\sigma \\ d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r} \end{array}$$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

## Notation:

- $r > 0$  is an index (analogous to  $n$  in  $\varphi_n$ ).

# Krein Systems, the continuous analogue of OPUC

$$\begin{array}{c} (d\mu, \mathbb{T}) \\ \text{OPUC} \\ \Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k \\ \left\| \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\|_{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty} |a_j|^2 \\ d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n \end{array}$$

$$\begin{array}{c} (d\sigma, \mathbb{R}) \\ \text{Krein Systems} \\ P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds \\ \left\| \int_0^{\infty} f(s) P(s, \lambda) ds \right\|_{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty} |f(s)|^2 ds \\ d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r} \end{array}$$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

## Notation:

- $r > 0$  is an index (analogous to  $n$  in  $\varphi_n$ ).
- $\lambda$  is the variable denoting points in the space  $\mathbb{R}$ .

# Krein Systems, the continuous analogue of OPUC

$(d\mu, \mathbb{T})$ OPUC	$(d\sigma, \mathbb{R})$ Krein Systems
$\Phi_n(z) = z^n + \sum_{k=0}^{n-1} b_{n,k} z^k$	$P(r, \lambda) = e^{i\lambda r} + \int_0^r b_r(s) e^{i\lambda s} ds$
$\left\  \sum_{j=0}^{\infty} a_j \varphi_j(z) \right\ _{L^2_{d\mu}(\mathbb{T})}^2 = \sum_{j=0}^{\infty}  a_j ^2$	$\left\  \int_0^{\infty} f(s) P(s, \lambda) ds \right\ _{L^2_{d\sigma(\lambda)}(\mathbb{R})}^2 = \int_0^{\infty}  f(s) ^2$
$d\mu = \frac{d\theta}{2\pi} \rightarrow \varphi_n(z) = z^n$	$d\sigma = \frac{d\lambda}{2\pi} \rightarrow P(r, \lambda) = e^{i\lambda r}$

Krein systems are the continuous analogue of OPUC, and so results for one can usually be carried over to the other.

## Notation:

- $r > 0$  is an index (analogous to  $n$  in  $\varphi_n$ ).
- $\lambda$  is the variable denoting points in the space  $\mathbb{R}$ .

We will be interested in the case when  $d\sigma = w(\lambda) \frac{d\lambda}{2\pi}$ , where  $w - 1 \in L^1(\mathbb{R}) + L^2(\mathbb{R})$  to ensure  $P(r, \lambda)$  exists.

# A Steklov problem for Krein Systems

## Problem (Steklov problem for Krein Systems)

*Does there exist  $p > 2$  for which  $\{P(r, \lambda)\}_{r \geq 0}$  is bounded in  $L^p_{loc}(w, \mathbb{R})$ ?*

# A Steklov problem for Krein Systems

## Problem (Steklov problem for Krein Systems)

*Does there exist  $p > 2$  for which  $\{P(r, \lambda)\}_{r \geq 0}$  is bounded in  $L^p_{loc}(w, \mathbb{R})$ ?*

More precisely, what conditions on  $w$  ensure that for each compact  $\Delta \subset \mathbb{R}$ , we have

$$\sup_{r \geq 0} \|P(r, \lambda)\|_{L^p(w, \Delta)} < \infty ?$$

# A Steklov problem for Krein Systems

## Problem (Steklov problem for Krein Systems)

*Does there exist  $p > 2$  for which  $\{P(r, \lambda)\}_{r \geq 0}$  is bounded in  $L^p_{loc}(w, \mathbb{R})$ ?*

More precisely, what conditions on  $w$  ensure that for each compact  $\Delta \subset \mathbb{R}$ , we have

$$\sup_{r \geq 0} \|P(r, \lambda)\|_{L^p(w, \Delta)} < \infty ?$$

Note this will follow if we can show

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w, \mathbb{R})} < \infty$$

for some  $p > 2$ .

# A Steklov problem for Krein Systems

## Problem (Steklov problem for Krein Systems)

*Does there exist  $p > 2$  for which  $\{P(r, \lambda)\}_{r \geq 0}$  is bounded in  $L^p_{loc}(w, \mathbb{R})$ ?*

More precisely, what conditions on  $w$  ensure that for each compact  $\Delta \subset \mathbb{R}$ , we have

$$\sup_{r \geq 0} \|P(r, \lambda)\|_{L^p(w, \Delta)} < \infty ?$$

Note this will follow if we can show

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w, \mathbb{R})} < \infty$$

for some  $p > 2$ .

**Remark:** In fact, this is nontrivial for  $p = 2$  and even  $p < 2$ .

# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $w - 1 \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 \leq p \leq 2 + \epsilon$ .

## Proof.

{





# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

{



# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

$$\begin{cases} P(r, \lambda) = e^{i\lambda r} + \mathcal{P}_{[0, r]} P(r, \lambda) \\ \end{cases}$$



# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

$$\begin{cases} P(r, \lambda) = e^{i\lambda r} + \mathcal{P}_{[0,r]}(P(r, \lambda) - e^{i\lambda r}) \\ \end{cases}$$



# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

$$\begin{cases} P(r, \lambda) &= e^{i\lambda r} + \mathcal{P}_{[0,r]}(P(r, \lambda) - e^{i\lambda r}) \\ 0 &= \mathcal{P}_{[0,r]} w P(r, \lambda) \end{cases}$$



# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

$$\begin{cases} P(r, \lambda) &= e^{i\lambda r} + \mathcal{P}_{[0,r]}(P(r, \lambda) - e^{i\lambda r}) \\ 0 &= \mathcal{P}_{[0,r]}(wP(r, \lambda) - e^{i\lambda r}) \end{cases}$$



# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

$$\begin{cases} P(r, \lambda) &= e^{i\lambda r} + \mathcal{P}_{[0,r]}(P(r, \lambda) - e^{i\lambda r}) \\ 0 &= \mathcal{P}_{[0,r]}(wP(r, \lambda) - e^{i\lambda r}) \end{cases}$$

Some algebra and the same process as before shows that

$$(I - Q_{w,p})X = -w^{-1/p'} \mathcal{P}_{[0,r]} w^{1/p'} (w^{1/p} - w^{-1/p'}) e^{i\lambda r}$$

where  $X = w^{1/p}(P(r, \lambda) - e^{i\lambda r})$ .



# Generalizing the OPUC result to Krein systems

## Theorem

Suppose  $w \in A_2(\mathbb{R})$  and  $\langle \lambda \rangle (w - 1) \in L^1(\mathbb{R})$ . Then there exists  $\epsilon > 0$  for which

$$\sup_{r \geq 0} \|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ . (Recall  $\langle \lambda \rangle \stackrel{\text{def}}{=} (1 + \lambda^2)^{1/2}$ .)

## Proof.

$$\begin{cases} P(r, \lambda) &= e^{i\lambda r} + \mathcal{P}_{[0,r]}(P(r, \lambda) - e^{i\lambda r}) \\ 0 &= \mathcal{P}_{[0,r]}(wP(r, \lambda) - e^{i\lambda r}) \end{cases}$$

Some algebra and the same process as before shows that

$$(I - Q_{w,p})X = -w^{-1/p'} \mathcal{P}_{[0,r]} w^{1/p'} (w^{1/p} - w^{-1/p'}) e^{i\lambda r}$$

where  $X = w^{1/p}(P(r, \lambda) - e^{i\lambda r})$ . Since  $I - Q_{w,p}$  is invertible, we're done. □

# An orthogonality property unique to $\mathbb{R}$

Recall that for OPUC we have

$$\mathcal{P}_{[0,n-1]} w \Phi_n = 0.$$



# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} wP(r, \lambda) = 0,$$

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} wP(r, \lambda) = 0,$$

i.e.

$$\left\langle wP(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \right\rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} wP(r, \lambda) = 0,$$

i.e.

$$\left\langle wP(r, \lambda), \int_0^r f'(s) e^{i\lambda s} ds \right\rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} wP(r, \lambda) = 0,$$

i.e.

$$\langle \lambda wP(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} \lambda wP(r, \lambda) = 0,$$

i.e.

$$\langle \lambda wP(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} \lambda w P(r, \lambda) = 0,$$

i.e.

$$\langle \lambda w P(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

**Idea:** let's try to bound

$$\lambda(P(r, \lambda) - e^{i\lambda r})$$

in  $L^p(w)$ .

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} \lambda w P(r, \lambda) = 0,$$

i.e.

$$\langle \lambda w P(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

**Idea:** let's try to bound

$$\lambda(P(r, \lambda) - e^{i\lambda r})$$

in  $L^p(w)$ .

$\lambda(P(r, \lambda) - e^{i\lambda r})$  isn't in  $L^p(w)$ , need to subtract off a constant-like term

# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} \lambda w P(r, \lambda) = 0,$$

i.e.

$$\langle \lambda w P(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

**Idea:** let's bound

$$R(r, \lambda) \stackrel{\text{def}}{=} \lambda(P(r, \lambda) - e^{i\lambda r}) - e^{i\lambda r} \alpha_\infty(r) - \alpha_2(r)$$

in  $L^p(w)$ .



# An orthogonality property unique to $\mathbb{R}$

Ignoring issues of integrability, we morally have

$$\mathcal{P}_{[0,r]} \lambda w P(r, \lambda) = 0,$$

i.e.

$$\langle \lambda w P(r, \lambda), \int_0^r f(s) e^{i\lambda s} ds \rangle_{d\lambda} = 0,$$

for all  $f \in C_c^\infty(0, r)$ .

**Idea:** let's bound

$$R(r, \lambda) \stackrel{\text{def}}{=} \lambda(P(r, \lambda) - e^{i\lambda r}) - e^{i\lambda r} \alpha_\infty(r) - \alpha_2(r)$$

in  $L^p(w)$ .

Under certain conditions on  $w$  (see next slide),  $\alpha_2(r) \in L^2_{dr}(\mathbb{R}^+)$ ,  $\alpha_\infty(r) \in L^\infty_{dr}(\mathbb{R}^+)$  and  $\lim_{r \rightarrow \infty} \alpha_\infty(r)$  exists.

## A remainder estimate

Estimating  $\|R(r, \lambda)\|_{L^p(W)}$  quantifies the decay of  $P(r, \lambda) - e^{i\lambda r}$  and provides an asymptotic expansion of sorts, in both  $\lambda$  and  $r$ .

## A remainder estimate

Estimating  $\|R(r, \lambda)\|_{L^p(w)}$  quantifies the decay of  $P(r, \lambda) - e^{i\lambda r}$  and provides an asymptotic expansion of sorts, in both  $\lambda$  and  $r$ .

### Theorem

Suppose  $w \in A_2(\mathbb{R})$ ,  $\langle \lambda \rangle^q (w - 1) \in L^1(\mathbb{R})$  for some  $q > 2$ , and

$$\exp \left( \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(w(t))}{t - \lambda} dt \right) - 1 = \int_0^{\infty} h(x) e^{i\lambda x} dx = \hat{h}(\lambda) \in H^2(\mathbb{C}^+).$$

Then there exists  $\epsilon > 0$  for which

$$\| \|R(r, \lambda)\|_{L^p(w)} \|_{L^\infty(dr, \mathbb{R}^+) + L^2(dr, \mathbb{R}^+)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ .

## A remainder estimate

Estimating  $\|R(r, \lambda)\|_{L^p(w)}$  quantifies the decay of  $P(r, \lambda) - e^{i\lambda r}$  and provides an asymptotic expansion of sorts, in both  $\lambda$  and  $r$ .

### Theorem

Suppose  $w \in A_2(\mathbb{R})$ ,  $\langle \lambda \rangle^q (w - 1) \in L^1(\mathbb{R})$  for some  $q > 2$ , and

$$\exp \left( \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\log(w(t))}{t - \lambda} dt \right) - 1 = \int_0^{\infty} h(x) e^{i\lambda x} dx = \hat{h}(\lambda) \in H^2(\mathbb{C}^+).$$

Then there exists  $\epsilon > 0$  for which

$$\| \|R(r, \lambda)\|_{L^p(w)} \|_{L^\infty(dr, \mathbb{R}^+) + L^2(dr, \mathbb{R}^+)} < \infty$$

whenever  $2 - \epsilon \leq p \leq 2 + \epsilon$ .

**Examples:** If  $-1 < \beta < 1$ , then  $w(\lambda) = |\lambda|^\beta$  in  $[-1, 1]$ , and equals 1 outside  $[-1, 1]$  satisfies all the conditions of the above.

# Parting questions/thoughts

## Parting questions/thoughts

- Idea: we saw that morally,  $\mathcal{P}_{[0,r]} \lambda^k w P(r, \lambda) = 0$ . So as far as the algebra is concerned, we might as well pretend  $P(r, \lambda)$  is generated by the weight  $\tilde{w} \stackrel{\text{def}}{=} q(\lambda)w$  for some polynomial  $q(\lambda) \geq 0$ , which we'll require to be in  $A_2(\mathbb{R})$ .

# Parting questions/thoughts

- Idea: we saw that morally,  $\mathcal{P}_{[0,r]} \lambda^k w P(r, \lambda) = 0$ . So as far as the algebra is concerned, we might as well pretend  $P(r, \lambda)$  is generated by the weight  $\tilde{w} \stackrel{\text{def}}{=} q(\lambda)w$  for some polynomial  $q(\lambda) \geq 0$ , which we'll require to be in  $A_2(\mathbb{R})$ .

Issue:  $w$  needs to be centered around 1, which would mean  $\tilde{w}$  would blow-up, meaning the  $A_2$  condition wouldn't be verified.

# Parting questions/thoughts

- Idea: we saw that morally,  $\mathcal{P}_{[0,r]} \lambda^k w P(r, \lambda) = 0$ . So as far as the algebra is concerned, we might as well pretend  $P(r, \lambda)$  is generated by the weight  $\tilde{w} \stackrel{\text{def}}{=} q(\lambda)w$  for some polynomial  $q(\lambda) \geq 0$ , which we'll require to be in  $A_2(\mathbb{R})$ .

Issue:  $w$  needs to be centered around 1, which would mean  $\tilde{w}$  would blow-up, meaning the  $A_2$  condition wouldn't be verified.

Possible solution: work with de Branges canonical systems, where  $w$  might not need to be centered around 1?



# Parting questions/thoughts

- Idea: we saw that morally,  $\mathcal{P}_{[0,r]} \lambda^k w P(r, \lambda) = 0$ . So as far as the algebra is concerned, we might as well pretend  $P(r, \lambda)$  is generated by the weight  $\tilde{w} \stackrel{\text{def}}{=} q(\lambda)w$  for some polynomial  $q(\lambda) \geq 0$ , which we'll require to be in  $A_2(\mathbb{R})$ .

Issue:  $w$  needs to be centered around 1, which would mean  $\tilde{w}$  would blow-up, meaning the  $A_2$  condition wouldn't be verified.

Possible solution: work with de Branges canonical systems, where  $w$  might not need to be centered around 1?

- Estimating  $\|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)}$  is kind of overkill. Seems much more natural to try and estimate

$$\|P(r, \lambda)\|_{L^p\left(\frac{w(\lambda)d\lambda}{1+\lambda^2}\right)}.$$

# Parting questions/thoughts

- Idea: we saw that morally,  $\mathcal{P}_{[0,r]} \lambda^k w P(r, \lambda) = 0$ . So as far as the algebra is concerned, we might as well pretend  $P(r, \lambda)$  is generated by the weight  $\tilde{w} \stackrel{\text{def}}{=} q(\lambda)w$  for some polynomial  $q(\lambda) \geq 0$ , which we'll require to be in  $A_2(\mathbb{R})$ .

Issue:  $w$  needs to be centered around 1, which would mean  $\tilde{w}$  would blow-up, meaning the  $A_2$  condition wouldn't be verified.

Possible solution: work with de Branges canonical systems, where  $w$  might not need to be centered around 1?

- Estimating  $\|P(r, \lambda) - e^{i\lambda r}\|_{L^p(w)}$  is kind of overkill. Seems much more natural to try and estimate

$$\|P(r, \lambda)\|_{L^p\left(\frac{w(\lambda)d\lambda}{1+\lambda^2}\right)}.$$

I have no clue how to come up with such an estimate beyond the content of this presentation.

*Thank you for Listening!*

*Thank you for Listening!*  
Just kidding!

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,2} = w^{1/2}\mathcal{P}_{[0,n-1]}w^{-1/2} - w^{-1/2}\mathcal{P}_{[0,n-1]}w^{1/2}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
  - $Q_{w,2}$  is antisymmetric, i.e.  $Q_{w,2}^* = -Q_{w,2}$ . Thus the spectrum of  $Q_{w,2}$  is pure imaginary and so  $I - \kappa Q_{w,2}$  must be invertible.



# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,2} = w^{1/2} \mathcal{P}_{[0,n-1]} w^{-1/2} - w^{-1/2} \mathcal{P}_{[0,n-1]} w^{1/2}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
  - $Q_{w,2}$  is antisymmetric, i.e.  $Q_{w,2}^* = -Q_{w,2}$ . Thus the spectrum of  $Q_{w,2}$  is pure imaginary and so  $I - \kappa Q_{w,2}$  must be invertible.
  - In fact, an inner-product computation shows  $\|(I - \kappa Q_{w,2})f\|_2^2 \geq \|f\|_2^2$ .

# Inverting $I - Q_{w,p}$


A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .



Algorithm:

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

- By geometric sum,  
 $\|(I - \frac{1}{N}Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{0}{N}Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

- By geometric sum,  
 $\|(I - \frac{1}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{0}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.
- Analytically interpolate with good  $p = 2$  estimate to get  
 $\|(I - \frac{1}{N} Q_{w,p})^{-1}\|_{p,p} \leq 5$  for  $2 \leq p \leq 2 + \epsilon_1$ , where  $\epsilon_1 < \epsilon_0$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .



Algorithm:


- By geometric sum,  
 $\|(I - \frac{1}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{0}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.
- Analytically interpolate with good  $p = 2$  estimate to get  
 $\|(I - \frac{1}{N} Q_{w,p})^{-1}\|_{p,p} \leq 5$  for  $2 \leq p \leq 2 + \epsilon_1$ , where  $\epsilon_1 < \epsilon_0$ .



# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

- By geometric sum,  
 $\|(I - \frac{2}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{1}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

- By geometric sum,  
 $\|(I - \frac{2}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{1}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.
- Analytically interpolate with good  $p = 2$  estimate to get  
 $\|(I - \frac{2}{N} Q_{w,p})^{-1}\|_{p,p} \leq 5$  for  $2 \leq p \leq 2 + \epsilon_2$ , where  $\epsilon_2 < \epsilon_1$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

- By geometric sum,  
 $\|(I - \frac{2}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{1}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.
- Analytically interpolate with good  $p = 2$  estimate to get  
 $\|(I - \frac{2}{N} Q_{w,p})^{-1}\|_{p,p} \leq 5$  for  $2 \leq p \leq 2 + \epsilon_2$ , where  $\epsilon_2 < \epsilon_1$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

- By geometric sum,  
 $\|(I - \frac{3}{N}Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{2}{N}Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .



Algorithm:

- By geometric sum,  
 $\|(I - \frac{3}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{2}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.
- Analytically interpolate with good  $p = 2$  estimate to get  
 $\|(I - \frac{3}{N} Q_{w,p})^{-1}\|_{p,p} \leq 5$  for  $2 \leq p \leq 2 + \epsilon_3$ , where  $\epsilon_3 < \epsilon_2$ .

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .



Algorithm:


- By geometric sum,  
 $\|(I - \frac{3}{N} Q_{w,p})^{-1}\|_{p,p} = \|(I - \frac{2}{N} Q_{w,p} - \frac{Q_{w,p}}{N})^{-1}\|_{p,p} \leq 10^{10}$  for  $p$  green.
- Analytically interpolate with good  $p = 2$  estimate to get  
 $\|(I - \frac{3}{N} Q_{w,p})^{-1}\|_{p,p} \leq 5$  for  $2 \leq p \leq 2 + \epsilon_3$ , where  $\epsilon_3 < \epsilon_2$ .



# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm: Repeat  $N$  times ...

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .




Algorithm:

# Inverting $I - Q_{w,p}$

A few facts:

- Recall  $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$ .
- $\|Q_{w,p}\|_{p,p} \leq C([w]_{A_2})$  for all  $2 \leq p \leq 2 + \epsilon_0$ .
- $\|(I - \kappa Q_{w,2})^{-1}\|_{2,2} \leq 1$  for all  $\kappa \in \mathbb{R}$ .
- Choose  $N$  large enough so that  $\|\frac{Q_{w,p}}{N}\|_{p,p} \leq \frac{1}{10}$ .



Algorithm:

Done!

*Thank you for Listening (for real this time)!*