

# How to represent a function in a quantum computer?

Michel Alexis

University of Bonn

June 25, ACOTCA

# How to represent a function in a quantum computer?

# How to represent a function in a quantum computer?

**More precisely:**

# How to represent a function in a quantum computer?

**More precisely:** the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors*  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  in  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and an input variable  $x \in [0, 1]$  and acts on a qubit (i.e., a unit vector in  $\mathbb{C}^2$ ) by the infinite  $2 \times 2$  matrix product  $U_{\infty}(x, \Psi) :=$

# How to represent a function in a quantum computer?

**More precisely:** the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors*  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  in  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and an input variable  $x \in [0, 1]$  and acts on a qubit (i.e., a unit vector in  $\mathbb{C}^2$ ) by the infinite  $2 \times 2$  matrix product  $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

# How to represent a function in a quantum computer?

**More precisely:** the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors*  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  in  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and an input variable  $x \in [0, 1]$  and acts on a qubit (i.e., a unit vector in  $\mathbb{C}^2$ ) by the infinite  $2 \times 2$  matrix product  $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where  $\theta = \arccos(x)$ ,

# How to represent a function in a quantum computer?

**More precisely:** the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors*  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  in  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and an input variable  $x \in [0, 1]$  and acts on a qubit (i.e., a unit vector in  $\mathbb{C}^2$ ) by the infinite  $2 \times 2$  matrix product  $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where  $\theta = \arccos(x)$ ,

and  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  are the Pauli matrices.

# How to represent a function in a quantum computer?

**More precisely:** the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors*  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  in  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and an input variable  $x \in [0, 1]$  and acts on a qubit (i.e., a unit vector in  $\mathbb{C}^2$ ) by the infinite  $2 \times 2$  matrix product  $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where  $\theta = \arccos(x)$ ,

and  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  are the Pauli matrices.

**Question:** Given a function  $f : [0, 1] \rightarrow [-1, 1]$ , does there exist an infinite phase factor sequence  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  for which  $f(x)$  is the imaginary part of the upper left entry of  $U_{\infty}(x, \Psi)$  ?



# How to represent a function in a quantum computer?

**More precisely:** the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors*  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  in  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and an input variable  $x \in [0, 1]$  and acts on a qubit (i.e., a unit vector in  $\mathbb{C}^2$ ) by the infinite  $2 \times 2$  matrix product  $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where  $\theta = \arccos(x)$ ,

and  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  are the Pauli matrices.

**Question:** Given a function  $f : [0, 1] \rightarrow [-1, 1]$ , does there exist an infinite phase factor sequence  $\Psi = \{\psi_k\}_{k=0}^{\infty}$  for which  $f(x)$  is the imaginary part of the upper left entry of  $U_{\infty}(x, \Psi)$  ?

**Why should we care?** QSP is apparently a very simple and physically intuitive quantum algorithm.

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

$$\text{Note } e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix}$$

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$

# Making sense of of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}.$$

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$ . Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$$

# Making sense of of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$ . Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$ . Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix. Indeed

$$\begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$



# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$ . Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix. Indeed

$$\begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \\ = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}.$$

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$ . Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix.

# Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where  $\theta = \arccos(x)$ ,  $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

Note  $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$  and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$ . Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix.

$$\text{So } U_d(x, \Psi) = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}.$$

# Main Theorem

# Main Theorem

Because the truncated matrices  $U_d(x, \Psi)$  have upper left entries which are  $poly(x)$ , the QSP problem is actually about approximating  $f$  by polynomials generated in this fashion.

# Main Theorem

Because the truncated matrices  $U_d(x, \Psi)$  have upper left entries which are *poly*( $x$ ), the QSP problem is actually about approximating  $f$  by polynomials generated in this fashion.

## Theorem (A.-Mnatsakanyan-Thiele, 2023)

*For each  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , there exists a unique phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ , and we have the nonlinear Plancherel identity*

$$\sum_k \log(1 + \tan^2 \psi_{|k|}) = -\frac{2}{\pi} \int_0^1 \log |1 - f(x)^2| \frac{dx}{\sqrt{1-x^2}}.$$

# Main Theorem

Because the truncated matrices  $U_d(x, \Psi)$  have upper left entries which are *poly*( $x$ ), the QSP problem is actually about approximating  $f$  by polynomials generated in this fashion.

## Theorem (A.-Mnatsakanyan-Thiele, 2023)

*For each  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , there exists a unique phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ , and we have the nonlinear Plancherel identity*

$$\sum_k \log(1 + \tan^2 \psi_{|k|}) = -\frac{2}{\pi} \int_0^1 \log |1 - f(x)^2| \frac{dx}{\sqrt{1-x^2}}.$$

The key idea here is to use nonlinear Fourier analysis!

# Some quick basics



## Some quick basics

Given  $f$  a function on the unit disc  $\mathbb{D} \subset \mathbb{C}$ , define its reflection

$$f^*(z) := \overline{f\left(\frac{1}{\bar{z}}\right)}.$$

## Some quick basics

Given  $f$  a function on the unit disc  $\mathbb{D} \subset \mathbb{C}$ , define its reflection

$$f^*(z) := \overline{f\left(\frac{1}{\bar{z}}\right)}.$$

If  $f$  is holomorphic on  $\mathbb{D}$ , then  $f^*$  is then holomorphic on the reflected disk  $\mathbb{D}^*$  at  $\infty$ .

## Some quick basics

Given  $f$  a function on the unit disc  $\mathbb{D} \subset \mathbb{C}$ , define its reflection

$$f^*(z) := \overline{f\left(\frac{1}{\bar{z}}\right)}.$$

If  $f$  is holomorphic on  $\mathbb{D}$ , then  $f^*$  is then holomorphic on the reflected disk  $\mathbb{D}^*$  at  $\infty$ .

Given a Laurent series  $f(z) = \sum_k a_k z^k$ , the  $*$  map reflects its Fourier support about the origin.

## Some quick basics

Given  $f$  a function on the unit disc  $\mathbb{D} \subset \mathbb{C}$ , define its reflection

$$f^*(z) := \overline{f\left(\frac{1}{\bar{z}}\right)}.$$

If  $f$  is holomorphic on  $\mathbb{D}$ , then  $f^*$  is then holomorphic on the reflected disk  $\mathbb{D}^*$  at  $\infty$ .

Given a Laurent series  $f(z) = \sum_k a_k z^k$ , the  $*$  map reflects its Fourier support about the origin.

Recall the 2-dimensional unitary group  $SU(2)$  consists of matrices  $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$  satisfying the determinant condition  $|\alpha|^2 + |\beta|^2 = 1$ .

# The $SU(2)$ -nonlinear Fourier transform (NLFT)

# The $SU(2)$ -nonlinear Fourier transform (NLFT)

Fourier transform :  $(F_k)_k \mapsto$  Laurent series  $\sum_k F_k z^k$ .

# The $SU(2)$ -nonlinear Fourier transform (NLFT)

Fourier transform :  $(F_k)_k \mapsto$  Laurent series  $\sum_k F_k z^k$ .

nonlinear Fourier transform:  $(F_k)_k \mapsto$  a matrix of Laurent series.

# The $SU(2)$ -nonlinear Fourier transform (NLFT)

Fourier transform :  $(F_k)_k \mapsto$  Laurent series  $\sum_k F_k z^k$ .

nonlinear Fourier transform:  $(F_k)_k \mapsto$  a matrix of Laurent series.

Define the nonlinear Fourier transform of a sequence  $(F_k)_k$  as the  $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as  $j$  increases.



# The $SU(2)$ -nonlinear Fourier transform (NLFT)

Fourier transform :  $(F_k)_k \mapsto$  Laurent series  $\sum_k F_k z^k$ .

nonlinear Fourier transform:  $(F_k)_k \mapsto$  a matrix of Laurent series.

Define the nonlinear Fourier transform of a sequence  $(F_k)_k$  as the  $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as  $j$  increases.

Note this is well-defined for compactly supported  $\{F_k\}$ , since only finitely many matrices above are not the identity.

# The $SU(2)$ -nonlinear Fourier transform (NLFT)

Fourier transform :  $(F_k)_k \mapsto$  Laurent series  $\sum_k F_k z^k$ .

nonlinear Fourier transform:  $(F_k)_k \mapsto$  a matrix of Laurent series.

Define the nonlinear Fourier transform of a sequence  $(F_k)_k$  as the  $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as  $j$  increases.

Note this is well-defined for compactly supported  $\{F_k\}$ , since only finitely many matrices above are not the identity.

We abbreviate this  $SU(2)$  matrix as the pair  $(a, b)$ .

# The $SU(2)$ -nonlinear Fourier transform (NLFT)

Fourier transform :  $(F_k)_k \mapsto$  Laurent series  $\sum_k F_k z^k$ .

nonlinear Fourier transform:  $(F_k)_k \mapsto$  a matrix of Laurent series.

Define the nonlinear Fourier transform of a sequence  $(F_k)_k$  as the  $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as  $j$  increases.

Note this is well-defined for compactly supported  $\{F_k\}$ , since only finitely many matrices above are not the identity.

We abbreviate this  $SU(2)$  matrix as the pair  $(a, b)$ .

Note  $(a, b)$  then satisfies the determinant condition  $|a|^2 + |b|^2 = 1$  for  $z \in \mathbb{T}$ .

# Why call this a Fourier transform?

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} = \left( \prod_j (1 + |F_j|^2)^{-\frac{1}{2}} \right) \prod_j \begin{pmatrix} 1 & F_j z^j \\ -\bar{F}_j z^{-j} & 1 \end{pmatrix}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right)$$



# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{aligned} \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} &\approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ &= \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}. \end{aligned}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$
$n = 2 :$		$n = 3 :$
$n \text{ even} :$		$n \text{ odd} :$
$a =$		$b =$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\begin{array}{lcl} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \left| \begin{array}{l} n = 1 : \quad \sum_j \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 2 : \\ n \text{ odd} : \\ b = \end{array} \right. \end{array}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\begin{array}{lcl} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \left| \right. \begin{array}{l} n = 1 : \quad \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : \\ n \text{ odd} : \\ b = \end{array} \\ n = 2 : \\ n \text{ even} : \\ a = \end{array}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \begin{pmatrix} 0 & F_{j_2} z^{j_2} \\ -\overline{F_{j_2}} z^{-j_2} & 0 \end{pmatrix}$	$n = 3 :$	
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} -F_{j_1} \overline{F_{j_2}} z^{j_1 - j_2} & 0 \\ 0 & -\overline{F_{j_1}} F_{j_2} z^{j_2 - j_1} \end{pmatrix}$	$n = 3 :$	
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix} \begin{pmatrix} 0 & F_{j_3} z^{j_3} \\ -\overline{F_{j_3}} z^{-j_3} & 0 \end{pmatrix}$
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	



# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix}$
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$$\left. \begin{array}{ll} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ n = 2 : & \sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix} \\ n \text{ even} : & \sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix} \\ a = & \end{array} \right| \begin{array}{ll} n = 1 : & \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : & \sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix} \\ n \text{ odd} : & \\ b = & \end{array}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\left. \begin{array}{ll} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ n = 2 : & \sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix} \\ n \text{ even} : & \sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix} \\ a = & \end{array} \right| \begin{array}{ll} n = 1 : & \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : & \sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix} \\ n \text{ odd} : & \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & O(F)^n \\ O(F)^n & 0 \end{pmatrix} \\ b = & \end{array}$$

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix}$
$n \text{ even} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix}$	$n \text{ odd} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & O(F)^n \\ O(F)^n & 0 \end{pmatrix}$
$a =$	$1 + \text{higher order terms}$	$b =$	

# Why call this a Fourier transform?

We do an informal computation, assuming  $F$  is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1, j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix}$
$n \text{ even} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix}$	$n \text{ odd} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & O(F)^n \\ O(F)^n & 0 \end{pmatrix}$
$a =$	$1 + \text{higher order terms}$	$b =$	$\sum_j F_j z^j + \text{higher order terms}$

# The $L^2$ theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where  $\{z_k\}$  are the zeros of  $a^*$  in  $\mathbb{D}$ .

# The $L^2$ theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where  $\{z_k\}$  are the zeros of  $a^*$  in  $\mathbb{D}$ . This is known as nonlinear Plancherel.

# The $L^2$ theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where  $\{z_k\}$  are the zeros of  $a^*$  in  $\mathbb{D}$ . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from  $\ell^2([0, \infty))$  onto the set of pairs  $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$  for which

- $a^*(0) > 0$ .
- $aa^* + bb^* = 1$  a.e. on  $\mathbb{T}$ .
- $a^*$  and  $b$  share no common inner factor.



# The $L^2$ theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where  $\{z_k\}$  are the zeros of  $a^*$  in  $\mathbb{D}$ . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from  $\ell^2([0, \infty))$  onto the set of pairs  $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$  for which

- $a^*(0) > 0$ .
- $aa^* + bb^* = 1$  a.e. on  $\mathbb{T}$ .
- $a^*$  and  $b$  share no common inner factor.

Similarly, the NLFT extends to a bijection from  $\ell^2((-\infty, -1])$  to pairs  $(a, b)$  for which  $(a, b^*)$  is like above and  $b^*$  has a zero of order 1 at  $z = 0$ .

# The $L^2$ theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where  $\{z_k\}$  are the zeros of  $a^*$  in  $\mathbb{D}$ . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from  $\ell^2([0, \infty))$  onto the set of pairs  $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$  for which

- $a^*(0) > 0$ .
- $aa^* + bb^* = 1$  a.e. on  $\mathbb{T}$ .
- $a^*$  and  $b$  share no common inner factor.

Similarly, the NLFT extends to a bijection from  $\ell^2((-\infty, -1])$  to pairs  $(a, b)$  for which  $(a, b^*)$  is like above and  $b^*$  has a zero of order 1 at  $z = 0$ . Given a sequence  $F \in \ell^2(\mathbb{Z})$ , define its NLFT by

$$(a, b) := (a_-, b_-)(a_+, b_+)$$

where  $(a_-, b_-)$  is the NLFT of  $(F_k \mathbf{1}_{k < 0})_{k \in \mathbb{Z}}$ , and  $(a_+, b_+)$  is the NLFT of  $(F_k \mathbf{1}_{k \geq 0})_{k \in \mathbb{Z}}$ .

# The $L^2$ theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where  $\{z_k\}$  are the zeros of  $a^*$  in  $\mathbb{D}$ . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from  $\ell^2([0, \infty))$  onto the set of pairs  $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$  for which

- $a^*(0) > 0$ .
- $aa^* + bb^* = 1$  a.e. on  $\mathbb{T}$ .
- $a^*$  and  $b$  share no common inner factor.

Similarly, the NLFT extends to a bijection from  $\ell^2((-\infty, -1])$  to pairs  $(a, b)$  for which  $(a, b^*)$  is like above and  $b^*$  has a zero of order 1 at  $z = 0$ . Given a sequence  $F \in \ell^2(\mathbb{Z})$ , define its NLFT by

$$(a, b) := (a_-, b_-)(a_+, b_+)$$

where  $(a_-, b_-)$  is the NLFT of  $(F_k \mathbf{1}_{k < 0})_{k \in \mathbb{Z}}$ , and  $(a_+, b_+)$  is the NLFT of  $(F_k \mathbf{1}_{k \geq 0})_{k \in \mathbb{Z}}$ . Unknown whether the NLFT is injective on  $\ell^2(\mathbb{Z})$  however.

# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

**First step:** do a change of variable.

# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

**First step:** do a change of variable. Namely, set  $z = e^{2i\theta}$  where  $\theta = \arccos(x)$ , and define an even sequence  $F$  by  $F_k := i \tan \psi_{|k|}$ .

# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

**First step:** do a change of variable. Namely, set  $z = e^{2i\theta}$  where  $\theta = \arccos(x)$ , and define an even sequence  $F$  by  $F_k := i \tan \psi_{|k|}$ . Then for the matrix  $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$  we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

**First step:** do a change of variable. Namely, set  $z = e^{2i\theta}$  where  $\theta = \arccos(x)$ , and define an even sequence  $F$  by  $F_k := i \tan \psi_{|k|}$ . Then for the matrix  $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$  we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

where  $(a_d, b_d)$  is the NLFT of the truncated sequence  $(F_k \mathbf{1}_{-d \leq k \leq d})_k$ .



# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

**First step:** do a change of variable. Namely, set  $z = e^{2i\theta}$  where  $\theta = \arccos(x)$ , and define an even sequence  $F$  by  $F_k := i \tan \psi_{|k|}$ . Then for the matrix  $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$  we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

where  $(a_d, b_d)$  is the NLFT of the truncated sequence  $(F_k \mathbf{1}_{-d \leq k \leq d})_k$ . In particular, one gets  $-ib_d(z)$  equals the imaginary part of the upper left entry of  $U_d(x, \Psi)$ .

# Solving the QSP problem, proof of Theorem

**Initial problem:** given  $f$  with  $\|f\|_\infty < \frac{1}{\sqrt{2}}$ , find a phase factor sequence  $\Psi$  such that the imaginary parts of the upper left entries of  $U_d(x, \Psi)$  converge to  $f$  in  $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$ .

**First step:** do a change of variable. Namely, set  $z = e^{2i\theta}$  where  $\theta = \arccos(x)$ , and define an even sequence  $F$  by  $F_k := i \tan \psi_{|k|}$ . Then for the matrix  $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$  we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

where  $(a_d, b_d)$  is the NLFT of the truncated sequence  $(F_k \mathbf{1}_{-d \leq k \leq d})_k$ . In particular, one gets  $-ib_d(z)$  equals the imaginary part of the upper left entry of  $U_d(x, \Psi)$ .

**New problem:** given  $b$  “with symmetry” on  $\mathbb{T}$  with  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  satisfy  $b_d \rightarrow b$ .

# Solving the new problem

**New problem:** given  $b$  “with symmetry” on  $\mathbb{T}$  with  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  satisfy  $b_d \rightarrow b$ .

# Solving the new problem

**New problem:** given  $b$  “with symmetry” on  $\mathbb{T}$  with  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  satisfy  $b_d \rightarrow b$ .

**Key idea:** we’re only given  $b$ , meaning we can choose any  $a$  we want. Let’s choose  $a^*$  outer on  $\mathbb{D}$  so that  $|a|^2 + |b|^2 = 1$  on  $\mathbb{T}$ .

# Solving the new problem

**New problem:** given  $b$  “with symmetry” on  $\mathbb{T}$  with  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  satisfy  $b_d \rightarrow b$ .

**Key idea:** we’re only given  $b$ , meaning we can choose any  $a$  we want. Let’s choose  $a^*$  outer on  $\mathbb{D}$  so that  $|a|^2 + |b|^2 = 1$  on  $\mathbb{T}$ .

**Reduced problem:** Given  $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$  such that  $a^*$  outer,  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , and  $b$  “has symmetry”, find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  converge to  $(a, b)$ .

# Solving the new problem

**New problem:** given  $b$  “with symmetry” on  $\mathbb{T}$  with  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  satisfy  $b_d \rightarrow b$ .

**Key idea:** we’re only given  $b$ , meaning we can choose any  $a$  we want. Let’s choose  $a^*$  outer on  $\mathbb{D}$  so that  $|a|^2 + |b|^2 = 1$  on  $\mathbb{T}$ .

**Reduced problem:** Given  $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$  such that  $a^*$  outer,  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , and  $b$  “has symmetry”, find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  converge to  $(a, b)$ .

**Soln:** Use first assumption to set up a fixed point mapping to solve for the Fourier coefficients. The second assumption implies the fixed point mapping is a contraction mapping, so there exist unique Fourier coefficients  $F \in \ell^2(\mathbb{Z})$  corresponding to  $(a, b)$ .

# Solving the new problem

**New problem:** given  $b$  “with symmetry” on  $\mathbb{T}$  with  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  satisfy  $b_d \rightarrow b$ .

**Key idea:** we’re only given  $b$ , meaning we can choose any  $a$  we want. Let’s choose  $a^*$  outer on  $\mathbb{D}$  so that  $|a|^2 + |b|^2 = 1$  on  $\mathbb{T}$ .

**Reduced problem:** Given  $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$  such that  $a^*$  outer,  $\|b\|_\infty < \frac{1}{\sqrt{2}}$ , and  $b$  “has symmetry”, find an even sequence of pure imaginary nonlinear Fourier coefficients  $F$  whose truncated NLFTs  $(a_d, b_d)$  converge to  $(a, b)$ .

**Soln:** Use first assumption to set up a fixed point mapping to solve for the Fourier coefficients. The second assumption implies the fixed point mapping is a contraction mapping, so there exist unique Fourier coefficients  $F \in \ell^2(\mathbb{Z})$  corresponding to  $(a, b)$ .

Then argue  $F$  is even and pure imaginary because  $b$  is “symmetric.”

*Thank you for Listening!*