

How to represent a function in a quantum computer?

Michel Alexis

Joint work with Gevorg Mnatsakanyan and Christoph Thiele

University of Bonn

July 5

How to represent a function in a quantum computer?

How to represent a function in a quantum computer?

More precisely:

How to represent a function in a quantum computer?

More precisely: the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors* $\Psi = \{\psi_k\}_{k=0}^{\infty}$ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ and an input variable $x \in [0, 1]$ and acts on a qubit (i.e., a unit vector in \mathbb{C}^2) by the infinite 2×2 matrix product $U_{\infty}(x, \Psi) :=$

How to represent a function in a quantum computer?

More precisely: the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors* $\Psi = \{\psi_k\}_{k=0}^{\infty}$ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ and an input variable $x \in [0, 1]$ and acts on a qubit (i.e., a unit vector in \mathbb{C}^2) by the infinite 2×2 matrix product $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

How to represent a function in a quantum computer?

More precisely: the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors* $\Psi = \{\psi_k\}_{k=0}^{\infty}$ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ and an input variable $x \in [0, 1]$ and acts on a qubit (i.e., a unit vector in \mathbb{C}^2) by the infinite 2×2 matrix product $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where $\theta = \arccos(x)$,

How to represent a function in a quantum computer?

More precisely: the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors* $\Psi = \{\psi_k\}_{k=0}^{\infty}$ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ and an input variable $x \in [0, 1]$ and acts on a qubit (i.e., a unit vector in \mathbb{C}^2) by the infinite 2×2 matrix product $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where $\theta = \arccos(x)$,

and $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ are the Pauli matrices.

How to represent a function in a quantum computer?

More precisely: the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors* $\Psi = \{\psi_k\}_{k=0}^{\infty}$ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ and an input variable $x \in [0, 1]$ and acts on a qubit (i.e., a unit vector in \mathbb{C}^2) by the infinite 2×2 matrix product $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where $\theta = \arccos(x)$,

and $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ are the Pauli matrices.

Question: Given a function $f : [0, 1] \rightarrow [-1, 1]$, does there exist an infinite phase factor sequence $\Psi = \{\psi_k\}_{k=0}^{\infty}$ for which $f(x)$ is the imaginary part of the upper left entry of $U_{\infty}(x, \Psi)$?

How to represent a function in a quantum computer?

More precisely: the Quantum signal processing algorithm (QSP) takes a sequence of *phase factors* $\Psi = \{\psi_k\}_{k=0}^{\infty}$ in $(-\frac{\pi}{2}, \frac{\pi}{2})$ and an input variable $x \in [0, 1]$ and acts on a qubit (i.e., a unit vector in \mathbb{C}^2) by the infinite 2×2 matrix product $U_{\infty}(x, \Psi) :=$

$$\dots e^{i\psi_d Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} e^{i\psi_0 Z} e^{i\theta X} e^{i\psi_1 Z} e^{i\theta X} \dots e^{i\theta X} e^{i\psi_d Z} \dots,$$

where $\theta = \arccos(x)$,

and $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ are the Pauli matrices.

Question: Given a function $f : [0, 1] \rightarrow [-1, 1]$, does there exist an infinite phase factor sequence $\Psi = \{\psi_k\}_{k=0}^{\infty}$ for which $f(x)$ is the imaginary part of the upper left entry of $U_{\infty}(x, \Psi)$?

Why should we care? QSP is apparently a very simple and physically intuitive quantum algorithm.

Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix}$

Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

$$\text{Note } e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$$

Making sense of of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}.$$

Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$. Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$$

Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$. Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Making sense of of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$. Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix. Indeed

$$\begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Making sense of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$. Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix. Indeed

$$\begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix} \\ = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}.$$

Making sense of of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$. Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Multiplying two such matrices together yields another such matrix.

Making sense of of the matrix product

Consider the truncated product

$$U_d(x, \Psi) := e^{i\psi_d Z} e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} \dots e^{i\psi_0 Z} \dots e^{i\theta X} e^{i\psi_{d-1} Z} e^{i\theta X} e^{i\psi_d Z},$$

where $\theta = \arccos(x)$, $X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Note $e^{i\theta X} = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} x & i\sqrt{1-x^2} \\ i\sqrt{1-x^2} & x \end{pmatrix}$ and

$e^{i\psi_j Z} = \begin{pmatrix} e^{i\psi_j} & 0 \\ 0 & e^{-i\psi_j} \end{pmatrix}$. Then

$$e^{i\theta X} e^{i\psi_j Z} = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}$$

Mutlplying two such matrices together yields another such matrix.

$$\text{So } U_d(x, \Psi) = \begin{pmatrix} \text{poly}(x) & \text{poly}(x)\sqrt{1-x^2} \\ \text{poly}(x)\sqrt{1-x^2} & \text{poly}(x) \end{pmatrix}.$$

Main Theorem

Main Theorem

Because the truncated matrices $U_d(x, \Psi)$ have upper left entries which are $\text{poly}(x)$, the QSP problem is actually about approximating $f : [0, 1] \rightarrow [-1, 1]$ by polynomials generated in this fashion.

Main Theorem

Because the truncated matrices $U_d(x, \Psi)$ have upper left entries which are *poly*(x), the QSP problem is actually about approximating $f : [0, 1] \rightarrow [-1, 1]$ by polynomials generated in this fashion.

Theorem (A.-Mnatsakanyan-Thiele, 2023)

For each f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, there exists a unique phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$, and we have the nonlinear Plancherel identity

$$\sum_k \log(1 + \tan^2 \psi_{|k|}) = -\frac{2}{\pi} \int_0^1 \log(1 - f(x)^2) \frac{dx}{\sqrt{1-x^2}}.$$

Main Theorem

Because the truncated matrices $U_d(x, \Psi)$ have upper left entries which are $\text{poly}(x)$, the QSP problem is actually about approximating $f : [0, 1] \rightarrow [-1, 1]$ by polynomials generated in this fashion.

Theorem (A.-Mnatsakanyan-Thiele, 2023)

For each f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, there exists a unique phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$, and we have the nonlinear Plancherel identity

$$\sum_k \log(1 + \tan^2 \psi_{|k|}) = -\frac{2}{\pi} \int_0^1 \log(1 - f(x)^2) \frac{dx}{\sqrt{1-x^2}}.$$

The key idea here is to use nonlinear Fourier analysis!

Some quick basics, Hardy spaces

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D}) \quad |$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

$$f(z) = \sum_{k \geq 0} c_k z^k \text{ analytic on } \mathbb{D}$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$f(z) = \sum_{k \geq 0} c_k z^k \text{ analytic on } \mathbb{D}$$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

$$f(z) = \sum_{k \leq 0} c_k z^k \text{ analytic on } \mathbb{D}^*$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$f(z) = \sum_{k \geq 0} c_k z^k \text{ analytic on } \mathbb{D}$$

$$f(0) = \int_{-\pi}^{\pi} f \frac{d\theta}{2\pi} = c_0$$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

$$f(z) = \sum_{k \leq 0} c_k z^k \text{ analytic on } \mathbb{D}^*$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$f(z) = \sum_{k \geq 0} c_k z^k \text{ analytic on } \mathbb{D}$$

$$f(0) = \int_{-\pi}^{\pi} f \frac{d\theta}{2\pi} = c_0$$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

$$f(z) = \sum_{k \leq 0} c_k z^k \text{ analytic on } \mathbb{D}^*$$

$$f(\infty) = c_0$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$f(z) = \sum_{k \geq 0} c_k z^k \text{ analytic on } \mathbb{D}$$

$$f(0) = \int_{-\pi}^{\pi} f \frac{d\theta}{2\pi} = c_0$$

Have projection $P_{\mathbb{D}} : L^2(\mathbb{T}) \rightarrow H^2(\mathbb{D})$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

$$f(z) = \sum_{k \leq 0} c_k z^k \text{ analytic on } \mathbb{D}^*$$

$$f(\infty) = c_0$$

Some quick basics, Hardy spaces

Given $f(z) = \sum_k c_k z^k$ define its reflection $f^*(z) := \sum_k \overline{c_{-k}} z^k$.

f is holomorphic on \mathbb{D} iff f^* holomorphic on the reflected disk \mathbb{D}^* at ∞ .

Need two subspaces of $L^2(\mathbb{T})$ of functions $f(z)$ with Fourier series $\sum_k c_k z^k$.

$$H^2(\mathbb{D})$$

$$f(z) = \sum_{k \geq 0} c_k z^k \text{ analytic on } \mathbb{D}$$

$$f(0) = \int_{-\pi}^{\pi} f \frac{d\theta}{2\pi} = c_0$$

$$\text{Have projection } P_{\mathbb{D}} : L^2(\mathbb{T}) \rightarrow H^2(\mathbb{D})$$

$$H^2(\mathbb{D}^*) := H^2(\mathbb{D})^*$$

$$f(z) = \sum_{k \leq 0} c_k z^k \text{ analytic on } \mathbb{D}^*$$

$$f(\infty) = c_0$$

$$P_{\mathbb{D}^*} : L^2(\mathbb{T}) \rightarrow H^2(\mathbb{D}^*)$$

The $SU(2)$ -nonlinear Fourier transform (NLFT)

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform :

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform	:	$(F_k)_k$	\mapsto	Laurent series $\sum_k F_k z^k$
nonlinear Fourier transform	:			

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform	: $(F_k)_k$	\mapsto	Laurent series $\sum_k F_k z^k$
nonlinear Fourier transform	: $(F_k)_k$	\mapsto	

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform	: $(F_k)_k$	\mapsto	Laurent series $\sum_k F_k z^k$
nonlinear Fourier transform	: $(F_k)_k$	\mapsto	a matrix of Laurent series.

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform $: (F_k)_k \mapsto$ Laurent series $\sum_k F_k z^k$

nonlinear Fourier transform $: (F_k)_k \mapsto$ a matrix of Laurent series.

The nonlinear Fourier transform of a sequence $(F_k)_k$ is the $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\bar{F}_j z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as j increases.

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform : $(F_k)_k \mapsto$ Laurent series $\sum_k F_k z^k$

nonlinear Fourier transform : $(F_k)_k \mapsto$ a matrix of Laurent series.

The nonlinear Fourier transform of a sequence $(F_k)_k$ is the $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\bar{F}_j z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as j increases.

Note this is well-defined for compactly supported $\{F_k\}$, since only finitely many matrices above are not the identity.

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform : $(F_k)_k \mapsto$ Laurent series $\sum_k F_k z^k$

nonlinear Fourier transform : $(F_k)_k \mapsto$ a matrix of Laurent series.

The nonlinear Fourier transform of a sequence $(F_k)_k$ is the $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\bar{F}_j z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as j increases.

Note this is well-defined for compactly supported $\{F_k\}$, since only finitely many matrices above are not the identity.

We abbreviate this $SU(2)$ matrix as the pair (a, b) .

The $SU(2)$ -nonlinear Fourier transform (NLFT)

Recall the special unitary group $SU(2)$ consists of matrices $\begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}$ satisfying the determinant condition $|\alpha|^2 + |\beta|^2 = 1$.

linear Fourier transform : $(F_k)_k \mapsto$ Laurent series $\sum_k F_k z^k$

nonlinear Fourier transform : $(F_k)_k \mapsto$ a matrix of Laurent series.

The nonlinear Fourier transform of a sequence $(F_k)_k$ is the $SU(2)$ -valued function

$$\begin{pmatrix} a(z) & b(z) \\ -b^*(z) & a^*(z) \end{pmatrix} = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\bar{F}_j z^{-j} & 1 \end{pmatrix}$$

where the matrix product should be read left to right as j increases.

Note this is well-defined for compactly supported $\{F_k\}$, since only finitely many matrices above are not the identity.

We abbreviate this $SU(2)$ matrix as the pair (a, b) .

Note the determinant condition $|a|^2 + |b|^2 = 1$ then holds for $z \in \mathbb{T}$.

Why call this a Fourier transform?

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} = \left(\prod_j (1 + |F_j|^2)^{-\frac{1}{2}} \right) \prod_j \begin{pmatrix} 1 & F_j z^j \\ -\bar{F}_j z^{-j} & 1 \end{pmatrix}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right)$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{aligned} \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} &\approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ &= \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}. \end{aligned}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$
$n = 2 :$		$n = 3 :$
$n \text{ even} :$		$n \text{ odd} :$
$a =$		$b =$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\begin{array}{lcl} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \left| \right. \begin{array}{l} n = 1 : \quad \sum_j \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 2 : \\ n \text{ odd} : \\ b = \end{array} \end{array}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\begin{array}{lcl} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \left| \right. \begin{array}{l} n = 1 : \quad \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : \\ n \text{ odd} : \\ b = \end{array} \\ n = 2 : \\ n \text{ even} : \\ a = \end{array}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\left. \begin{array}{ll} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ n = 2 : & \sum_{j_1 < j_2} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \begin{pmatrix} 0 & F_{j_2} z^{j_2} \\ -\overline{F_{j_2}} z^{-j_2} & 0 \end{pmatrix} \\ n \text{ even} : & \\ a = & \end{array} \right| \begin{array}{ll} n = 1 : & \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : & \\ n \text{ odd} : & \\ b = & \end{array}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1 < j_2} \begin{pmatrix} -F_{j_1} \overline{F_{j_2}} z^{j_1 - j_2} & 0 \\ 0 & -\overline{F_{j_1}} F_{j_2} z^{j_2 - j_1} \end{pmatrix}$	$n = 3 :$	
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix} \begin{pmatrix} 0 & F_{j_3} z^{j_3} \\ -\overline{F_{j_3}} z^{-j_3} & 0 \end{pmatrix}$
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix}$
$n \text{ even} :$		$n \text{ odd} :$	
$a =$		$b =$	

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$$\left. \begin{array}{ll} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ n = 2 : & \sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix} \\ n \text{ even} : & \sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix} \\ a = & \end{array} \right| \begin{array}{ll} n = 1 : & \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : & \sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix} \\ n \text{ odd} : & \\ b = & \end{array}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix} .$$

$$\left. \begin{array}{ll} n = 0 : & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ n = 2 : & \sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix} \\ n \text{ even} : & \sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix} \\ a = & \end{array} \right| \begin{array}{ll} n = 1 : & \begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix} \\ n = 3 : & \sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix} \\ n \text{ odd} : & \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & O(F)^n \\ O(F)^n & 0 \end{pmatrix} \\ b = & \end{array}$$

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix}$
$n \text{ even} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix}$	$n \text{ odd} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & O(F)^n \\ O(F)^n & 0 \end{pmatrix}$
$a =$	$1 + \text{higher order terms}$	$b =$	

Why call this a Fourier transform?

We do an informal computation, assuming F is “small.”

$$\begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \approx \prod_j \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & F_j z^j \\ -\overline{F_j} z^{-j} & 0 \end{pmatrix} \right) \\ = \sum_{n \geq 0} \sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & F_{j_1} z^{j_1} \\ -\overline{F_{j_1}} z^{-j_1} & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & F_{j_n} z^{j_n} \\ -\overline{F_{j_n}} z^{-j_n} & 0 \end{pmatrix}.$$

$n = 0 :$	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$n = 1 :$	$\begin{pmatrix} 0 & \sum_j F_j z^j \\ -\sum_j \overline{F_j} z^{-j} & 0 \end{pmatrix}$
$n = 2 :$	$\sum_{j_1 < j_2} \begin{pmatrix} O(F)^2 & 0 \\ 0 & O(F)^2 \end{pmatrix}$	$n = 3 :$	$\sum_{j_1 < j_2 < j_3} \begin{pmatrix} 0 & O(F)^3 \\ O(F)^3 & 0 \end{pmatrix}$
$n \text{ even} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} O(F)^n & 0 \\ 0 & O(F)^n \end{pmatrix}$	$n \text{ odd} :$	$\sum_{j_1 < \dots < j_n} \begin{pmatrix} 0 & O(F)^n \\ O(F)^n & 0 \end{pmatrix}$
$a =$	$1 + \text{higher order terms}$	$b =$	$\sum_j F_j z^j + \text{higher order terms}$


The Laurent series a and b

The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b


The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a , b



The Laurent series a and b


A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a , b



Note that the NLFT plays well with truncations of sequences,

The Laurent series a and b


A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b



Note that the NLFT plays well with truncations of sequences, e.g., if (a_-, b_-) and (a_+, b_+) are the NLFTs of $(F_j \mathbf{1}_{j < 0})_j$ and $(F_j \mathbf{1}_{j \geq 0})_j$, then

The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b




Note that the NLFT plays well with truncations of sequences, e.g., if (a_-, b_-) and (a_+, b_+) are the NLFTs of $(F_j \mathbf{1}_{j < 0})_j$ and $(F_j \mathbf{1}_{j \geq 0})_j$, then

$$(a, b) = \prod_{j=-\infty}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b




Note that the NLFT plays well with truncations of sequences, e.g., if (a_-, b_-) and (a_+, b_+) are the NLFTs of $(F_j \mathbf{1}_{j < 0})_j$ and $(F_j \mathbf{1}_{j \geq 0})_j$, then

$$(a, b) = \prod_{j=-\infty}^{-1} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix} \prod_{j=0}^{\infty} \frac{1}{\sqrt{1 + |F_j|^2}} \begin{pmatrix} 1 & F_j z^j \\ -\overline{F_j} z^{-j} & 1 \end{pmatrix}$$

The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b




Note that the NLFT plays well with truncations of sequences, e.g., if (a_-, b_-) and (a_+, b_+) are the NLFTs of $(F_j \mathbf{1}_{j < 0})_j$ and $(F_j \mathbf{1}_{j \geq 0})_j$, then

$$(a, b) = (a_-, b_-)(a_+, b_+).$$

The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b




Note that the NLFT plays well with truncations of sequences, e.g., if (a_-, b_-) and (a_+, b_+) are the NLFTs of $(F_j \mathbf{1}_{j < 0})_j$ and $(F_j \mathbf{1}_{j \geq 0})_j$, then

$$(a, b) = (a_-, b_-)(a_+, b_+).$$

By multilinear expansion, we also have the formula $a^*(0) = \frac{1}{\prod_j (1 + |F_j|^2)^{\frac{1}{2}}}$.

The Laurent series a and b

A more careful multilinear expansion shows that the support of F dictates the Fourier support of Laurent polynomials a, b



Note that the NLFT plays well with truncations of sequences, e.g., if (a_-, b_-) and (a_+, b_+) are the NLFTs of $(F_j \mathbf{1}_{j < 0})_j$ and $(F_j \mathbf{1}_{j \geq 0})_j$, then

$$(a, b) = (a_-, b_-)(a_+, b_+).$$

By multilinear expansion, we also have the formula $a^*(0) = \frac{1}{\prod_j (1 + |F_j|^2)^{\frac{1}{2}}}$.

Thus $a^*(0) = a_-^*(0)a_+^*(0)$.

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} .

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel.

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H}

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

- $a^*(0) > 0$.

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

- $a^*(0) > 0$.
- $aa^* + bb^* = 1$ a.e. on \mathbb{T} .

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

- $a^*(0) > 0$.
- $aa^* + bb^* = 1$ a.e. on \mathbb{T} .
- a^* and b share no common inner factor.

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

- $a^*(0) > 0$.
- $aa^* + bb^* = 1$ a.e. on \mathbb{T} .
- a^* and b share no common inner factor.

Similarly, NLFT extends to a bijection from $\ell^2((-\infty, -1])$ to pairs (a, b) similar to above and b has vanishing constant Fourier coeff.

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

- $a^*(0) > 0$.
- $aa^* + bb^* = 1$ a.e. on \mathbb{T} .
- a^* and b share no common inner factor.

Similarly, NLFT extends to a bijection from $\ell^2((-\infty, -1])$ to pairs (a, b) similar to above and b has vanishing constant Fourier coeff.

Given a sequence $F \in \ell^2(\mathbb{Z})$, define its NLFT by

$$(a, b) := (a_-, b_-)(a_+, b_+)$$

where (a_-, b_-) is the NLFT of $(F_k \mathbf{1}_{k < 0})_{k \in \mathbb{Z}}$, and (a_+, b_+) is the NLFT of $(F_k \mathbf{1}_{k \geq 0})_{k \in \mathbb{Z}}$.

The L^2 theory for nonlinear Fourier transforms

A contour integral yields

$$-\sum_{n \in \mathbb{Z}} \log(1 + |F_n|^2) = \int_{\mathbb{T}} \log(1 - |b|^2) + 2 \sum_k \log |z_k|.$$

where $\{z_k\}$ are the zeros of a^* in \mathbb{D} . This is known as nonlinear Plancherel. Using nonlinear Plancherel, one can show the NLFT extends to a bijection from $\ell^2([0, \infty))$ onto the set \mathbf{H} of pairs $(a, b) \in H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$ for which

- $a^*(0) > 0$.
- $aa^* + bb^* = 1$ a.e. on \mathbb{T} .
- a^* and b share no common inner factor.

Similarly, NLFT extends to a bijection from $\ell^2((-\infty, -1])$ to pairs (a, b) similar to above and b has vanishing constant Fourier coeff.

Given a sequence $F \in \ell^2(\mathbb{Z})$, define its NLFT by

$$(a, b) := (a_-, b_-)(a_+, b_+)$$

where (a_-, b_-) is the NLFT of $(F_k \mathbf{1}_{k < 0})_{k \in \mathbb{Z}}$, and (a_+, b_+) is the NLFT of $(F_k \mathbf{1}_{k \geq 0})_{k \in \mathbb{Z}}$. Unknown whether the NLFT is injective on $\ell^2(\mathbb{Z})$ however.

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

First step: do a change of variable.

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

First step: do a change of variable. Namely, set $z = e^{2i\theta}$ where $\theta = \arccos(x)$, and define an even sequence F by $F_k := i \tan \psi_{|k|}$.

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

First step: do a change of variable. Namely, set $z = e^{2i\theta}$ where $\theta = \arccos(x)$, and define an even sequence F by $F_k := i \tan \psi_{|k|}$. Then for the matrix $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

First step: do a change of variable. Namely, set $z = e^{2i\theta}$ where $\theta = \arccos(x)$, and define an even sequence F by $F_k := i \tan \psi_{|k|}$. Then for the matrix $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

where (a_d, b_d) is the NLFT of the truncated sequence $(F_k \mathbf{1}_{-d \leq k \leq d})_k$.

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

First step: do a change of variable. Namely, set $z = e^{2i\theta}$ where $\theta = \arccos(x)$, and define an even sequence F by $F_k := i \tan \psi_{|k|}$. Then for the matrix $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

where (a_d, b_d) is the NLFT of the truncated sequence $(F_k \mathbf{1}_{-d \leq k \leq d})_k$. In particular, one gets $-ib_d(z)$ equals the imaginary part of the upper left entry of $U_d(x, \Psi)$.

Solving the QSP problem, proof of Theorem

Initial problem: given f with $\|f\|_\infty < \frac{1}{\sqrt{2}}$, find a phase factor sequence Ψ such that the imaginary parts of the upper left entries of $U_d(x, \Psi)$ converge to f in $L^2\left(\frac{dx}{\sqrt{1-x^2}}\right)$.

First step: do a change of variable. Namely, set $z = e^{2i\theta}$ where $\theta = \arccos(x)$, and define an even sequence F by $F_k := i \tan \psi_{|k|}$. Then for the matrix $M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ we have

$$MU_d(x, \Psi)M^{-1} := \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix} \begin{pmatrix} a_d & b_d \\ -b_d^* & a_d^* \end{pmatrix} \begin{pmatrix} z^{d/2} & 0 \\ 0 & z^{-d/2} \end{pmatrix}$$

where (a_d, b_d) is the NLFT of the truncated sequence $(F_k \mathbf{1}_{-d \leq k \leq d})_k$. In particular, one gets $-ib_d(z)$ equals the imaginary part of the upper left entry of $U_d(x, \Psi)$.

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Picking a

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Picking a

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Key idea: we’re only given b , meaning we can choose any a satisfying $|a|^2 + |b|^2 = 1$ that we want.

Picking a

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Key idea: we’re only given b , meaning we can choose any a satisfying $|a|^2 + |b|^2 = 1$ that we want. Let’s choose a as nice as possible:

Picking a

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Key idea: we’re only given b , meaning we can choose any a satisfying $|a|^2 + |b|^2 = 1$ that we want. Let’s choose a as nice as possible: let’s choose a so that a^* and $\frac{1}{a^*}$ are holomorphic on \mathbb{D} ,

Picking a

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Key idea: we’re only given b , meaning we can choose any a satisfying $|a|^2 + |b|^2 = 1$ that we want. Let’s choose a as nice as possible: let’s choose a so that a^* and $\frac{1}{a^*}$ are holomorphic on \mathbb{D} , for instance we can choose a^* to be outer,

$$a^*(z) := \exp \left(\int_{-\pi}^{\pi} \frac{e^{i\eta} + z}{e^{i\eta} - z} \log \sqrt{1 - |b(e^{i\eta})|^2} \frac{d\eta}{2\pi} \right).$$

Picking a

New problem: given b “with symmetry” on \mathbb{T} with $\|b\|_\infty < \frac{1}{\sqrt{2}}$, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) satisfy $b_d \rightarrow b$.

Key idea: we’re only given b , meaning we can choose any a satisfying $|a|^2 + |b|^2 = 1$ that we want. Let’s choose a as nice as possible: let’s choose a so that a^* and $\frac{1}{a^*}$ are holomorphic on \mathbb{D} , for instance we can choose a^* to be outer,

$$a^*(z) := \exp \left(\int_{-\pi}^{\pi} \frac{e^{i\eta} + z}{e^{i\eta} - z} \log \sqrt{1 - |b(e^{i\eta})|^2} \frac{d\eta}{2\pi} \right).$$

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) .

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1}$

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $a_- = a_+^* a + b b_+^*$ and $b_- = -b_+ a + a_+ b$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-}{a} = a_+^* + \frac{b}{a}b_+^*$ and $b_- = -b_+a + a_+b$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-}{a} = a_+^* + \frac{b}{a}b_+^*$ and $\frac{b_-}{a} = -b_+ + a_+\frac{b}{a}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-^*}{a^*} = a_+ + \frac{b^*}{a^*}b_+$ and $\frac{b_-}{a} = -b_+ + a_+\frac{b}{a}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-^*}{a^*} = a_+ + \frac{b^*}{a^*}b_+$ and $\frac{b_-}{a} = -b_+ + a_+\frac{b}{a}$.
Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-^*}{a^*} = a_+ + \frac{b_+^*}{a^*}b_+$ and $\frac{b_-}{a} = -b_+ + a_+\frac{b}{a}$.
Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$. So acting on the first equation by $P_{\mathbb{D}^*}$ should give the left side is $\frac{a_-^*(0)}{a^*(0)} =$

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-^*}{a^*} = a_+ + \frac{b_-^*}{a^*}b_+$ and $\frac{b_-}{a} = -b_+ + a_+\frac{b}{a}$.

Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$. So acting on the first equation by

$P_{\mathbb{D}^*}$ should give the left side is $\frac{a_-^*(0)}{a^*(0)} = \frac{a_-^*(0)}{a_-^*(0)a_+^*(0)}$

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{a_-^*}{a^*} = a_+ + \frac{b_-^*}{a^*}b_+$ and $\frac{b_-}{a} = -b_+ + a_+\frac{b}{a}$.
Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$. So acting on the first equation by $P_{\mathbb{D}^*}$ should give the left side is $\frac{a_-^*(0)}{a^*(0)} = \frac{a_-^*(0)}{a_-^*(0)a_+^*(0)} = \frac{1}{a_+^*(0)}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $P_{\mathbb{D}^*} \frac{a_-^*}{a^*} = P_{\mathbb{D}^*} a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$. Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$. So acting on the first equation by $P_{\mathbb{D}^*}$ should give the left side is $\frac{a_-^*(0)}{a^*(0)} = \frac{a_-^*(0)}{a_-^*(0)a_+^*(0)} = \frac{1}{a_+^*(0)}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = P_{\mathbb{D}^*} a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$.

Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$. So acting on the first equation by

$$P_{\mathbb{D}^*} \text{ should give the left side is } \frac{a_-^*(0)}{a^*(0)} = \frac{a_-^*(0)}{a_-^*(0)a_+^*(0)} = \frac{1}{a_+^*(0)}.$$

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$.

Because $1/a^*$ is holomorphic, so is the left side of the first equation, i.e., has Fourier support on $[0, \infty)$. So acting on the first equation by

$P_{\mathbb{D}^*}$ should give the left side is $\frac{a_-^*(0)}{a^*(0)} = \frac{a_-^*(0)}{a_-^*(0)a_+^*(0)} = \frac{1}{a_+^*(0)}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b_+}{a}$.

Again because $1/a$ is antiholomorphic, it has Fourier support on $(-\infty, 0]$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$.

Again because $1/a$ is antiholomorphic, it has Fourier support on $(-\infty, 0]$. When multiplied by b_- , which has Fourier support on $(-\infty, -1]$, the product has Fourier support on $(-\infty, -1]$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $\frac{b_-}{a} = -b_+ + a_+ \frac{b}{a}$.

Again because $1/a$ is antiholomorphic, it has Fourier support on $(-\infty, 0]$. When multiplied by b_- , which has Fourier support on $(-\infty, -1]$, the product has Fourier support on $(-\infty, -1]$. Thus acting on the second equation by $P_{\mathbb{D}}$ gives a left side of 0.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $P_{\mathbb{D}} \frac{b_-}{a} = -P_{\mathbb{D}} b_+ + P_{\mathbb{D}} a_+ \frac{b}{a}$.

Again because $1/a$ is antiholomorphic, it has Fourier support on $(-\infty, 0]$. When multiplied by b_- , which has Fourier support on $(-\infty, -1]$, the product has Fourier support on $(-\infty, -1]$. Thus acting on the second equation by $P_{\mathbb{D}}$ gives a left side of 0.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $0 = -P_{\mathbb{D}} b_+ + P_{\mathbb{D}} a_+ \frac{b}{a}$.

Again because $1/a$ is antiholomorphic, it has Fourier support on $(-\infty, 0]$. When multiplied by b_- , which has Fourier support on $(-\infty, -1]$, the product has Fourier support on $(-\infty, -1]$. Thus acting on the second equation by $P_{\mathbb{D}}$ gives a left side of 0.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $0 = -b_+ + P_{\mathbb{D}} a_+ \frac{b_+}{a_+}$.

Again because $1/a$ is antiholomorphic, it has Fourier support on $(-\infty, 0]$. When multiplied by b_- , which has Fourier support on $(-\infty, -1]$, the product has Fourier support on $(-\infty, -1]$. Thus acting on the second equation by $P_{\mathbb{D}}$ gives a left side of 0.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $0 = -b_+ + P_{\mathbb{D}} a_+ \frac{b}{a}$.

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $0 = -b_+ + P_{\mathbb{D}} a_+ \frac{b}{a}$.
- So a necessary condition for (a_+, b_+) is that the vector $(A, B) := a_+^*(0)(a_+, b_+)$ satisfies

Solving the nonlinear Fourier analysis problem, part 1

Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, and b “has symmetry”, find “nice” nonlinear Fourier coefficients F whose truncated NLFTs (a_d, b_d) converge to (a, b) .

“Solution:”

- (a, b) is a NLFT of $F \in \ell^2(\mathbb{Z})$ iff $(a, b) = (a_-, b_-)(a_+, b_+)$, where (a_-, b_-) is the NLFT of the left half of F , while (a_+, b_+) is the NLFT of the right half.
- To recover F , suffices to solve for (a_+, b_+) . Indeed, once we have (a_+, b_+) , we compute $(a_-, b_-) = (a, b)(a_+, b_+)^{-1} = (a, b)(a_+^*, -b_+)$.
- So $\frac{1}{a_+^*(0)} = a_+ + P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} b_+$ and $0 = -b_+ + P_{\mathbb{D}} a_+ \frac{b}{a}$.
- So a necessary condition for (a_+, b_+) is that the vector $(A, B) := a_+^*(0)(a_+, b_+)$ satisfies

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b_+^*}{a_+^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Very Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, factor $(a, b) = (a_-, b_-)(a_+, b_+)$.
 If we can solve for $(A, B) = a_+^*(0)(a_+, b_+)$ in

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

then this will basically give us (a_+, b_+) and show that (a, b) is an NLFT.

Very Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, factor $(a, b) = (a_-, b_-)(a_+, b_+)$.
 If we can solve for $(A, B) = a_+^*(0)(a_+, b_+)$ in

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

then this will basically give us (a_+, b_+) and show that (a, b) is an NLFT.

But $\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} = I + \begin{pmatrix} 0 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 0 \end{pmatrix} =: I + M.$

Very Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, factor $(a, b) = (a_-, b_-)(a_+, b_+)$.
 If we can solve for $(A, B) = a_+^*(0)(a_+, b_+)$ in

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

then this will basically give us (a_+, b_+) and show that (a, b) is an NLFT.

But $\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} = I + \begin{pmatrix} 0 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 0 \end{pmatrix} =: I + M.$

By the determinant identity $|a|^2 + |b|^2 = 1$ on \mathbb{T} and the assumption $\|b\|_\infty < \frac{1}{\sqrt{2}}$, we get $\|\frac{b}{a}\|_\infty < 1$.

Very Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, factor $(a, b) = (a_-, b_-)(a_+, b_+)$.
If we can solve for $(A, B) = a_+^*(0)(a_+, b_+)$ in

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

then this will basically give us (a_+, b_+) and show that (a, b) is an NLFT.

But $\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} = I + \begin{pmatrix} 0 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 0 \end{pmatrix} =: I + M.$

By the determinant identity $|a|^2 + |b|^2 = 1$ on \mathbb{T} and the assumption $\|b\|_\infty < \frac{1}{\sqrt{2}}$, we get $\|\frac{b}{a}\|_\infty < 1$.

So have the operator norm estimate $\|M\| \leq \|\frac{b}{a}\|_\infty < 1$ (viewing M as an operator on $H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$).

Very Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, factor $(a, b) = (a_-, b_-)(a_+, b_+)$.
If we can solve for $(A, B) = a_+^*(0)(a_+, b_+)$ in

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

then this will basically give us (a_+, b_+) and show that (a, b) is an NLFT.

But $\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} = I + \begin{pmatrix} 0 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 0 \end{pmatrix} =: I + M.$

By the determinant identity $|a|^2 + |b|^2 = 1$ on \mathbb{T} and the assumption $\|b\|_\infty < \frac{1}{\sqrt{2}}$, we get $\|\frac{b}{a}\|_\infty < 1$.

So have the operator norm estimate $\|M\| \leq \|\frac{b}{a}\|_\infty < 1$ (viewing M as an operator on $H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$).

Thus by geometric sum, we have $(I + M)^{-1} = \sum_{k=0}^{\infty} (-M)^k$ on

$H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$. So $\begin{pmatrix} A \\ B \end{pmatrix} = \sum_{k=0}^{\infty} (-M)^k \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$

Very Reduced problem: Given $(a, b) \in H^2(\mathbb{D}^*) \times L^\infty(\mathbb{T})$ such that a^* outer, $\|b\|_\infty < \frac{1}{\sqrt{2}}$, factor $(a, b) = (a_-, b_-)(a_+, b_+)$.
If we can solve for $(A, B) = a_+^*(0)(a_+, b_+)$ in

$$\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

then this will basically give us (a_+, b_+) and show that (a, b) is an NLFT.

But $\begin{pmatrix} 1 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 1 \end{pmatrix} = I + \begin{pmatrix} 0 & P_{\mathbb{D}^*} \frac{b^*}{a^*} \\ -P_{\mathbb{D}} \frac{b}{a} & 0 \end{pmatrix} =: I + M.$

By the determinant identity $|a|^2 + |b|^2 = 1$ on \mathbb{T} and the assumption $\|b\|_\infty < \frac{1}{\sqrt{2}}$, we get $\|\frac{b}{a}\|_\infty < 1$.

So have the operator norm estimate $\|M\| \leq \|\frac{b}{a}\|_\infty < 1$ (viewing M as an operator on $H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$).

Thus by geometric sum, we have $(I + M)^{-1} = \sum_{k=0}^{\infty} (-M)^k$ on

$H^2(\mathbb{D}^*) \times H^2(\mathbb{D})$. So $\begin{pmatrix} A \\ B \end{pmatrix} = \sum_{k=0}^{\infty} (-M)^k \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$



Thank you for Listening!