

# A Two-Weight T1 Theorem for Singular Integral Operators with respect to Doubling Measures

Michel Alexis

McMaster University

CMS Summer Meeting 2022

# Basic Defintions

# Basic Definitions

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

# Basic Definitions

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

## Definition (CZO)

$T$  is a (smooth) Calderón-Zygmund operator on functions if

# Basic Definitions

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

## Definition (CZO's)

$T$  is a (smooth) Calderón-Zygmund operator on functions if

$$Tf(x) \equiv \int K(x, y)f(y) dy,$$

# Basic Definitions

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

## Definition (CZO)

$T$  is a (smooth) Calderón-Zygmund operator on functions if

$$Tf(x) \equiv \int K(x, y)f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

# Basic Definitions

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

## Definition (CZO)

$T$  is a (smooth) Calderón-Zygmund operator on functions if

$$Tf(x) \equiv \int K(x, y)f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition.

# Basic Definitions

## Definition (Doubling)

A measure  $\mu$  is doubling if there exists a constant  $C$  such that  $\mu(B(x, 2r)) \leq C\mu(B(x, r))$  for all  $x, r$ .

## Definition (CZO)

$T$  is a (smooth) Calderón-Zygmund operator on functions if

$$Tf(x) \equiv \int K(x, y)f(y) dy,$$

where  $K$  is smooth, satisfies the decay condition

$$|\nabla_x^j K(x, y)|, |\nabla_y^j K(x, y)| \lesssim \frac{1}{|x - y|^{n+j}},$$

and some continuity condition. Example:  $K(x, y) = \frac{1}{x-y}$  in dimension 1.



# A T1 theorem for doubling measures

# A T1 theorem for doubling measures

Given a measure  $\mu$ , write  $T_\mu(f) \equiv T(f\mu)$ .

# A T1 theorem for doubling measures

Given a measure  $\mu$ , write  $T_\mu(f) \equiv T(f\mu)$ .

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

*Assume  $\omega, \sigma$  are doubling measures and  $T$  is a Calderón-Zygmund operator. Then for all functions  $f \in L^2(\sigma)$ , we have*

$$\|T_\sigma f(x)\|_{L^2(\omega)} \lesssim_{\sigma, \omega} (\mathfrak{T} + \mathfrak{T}^* + \sqrt{A_2(\omega, \sigma)}) \|f\|_{L^2(\sigma)},$$

# A T1 theorem for doubling measures

Given a measure  $\mu$ , write  $T_\mu(f) \equiv T(f\mu)$ .

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

*Assume  $\omega, \sigma$  are doubling measures and  $T$  is a Calderón-Zygmund operator. Then for all functions  $f \in L^2(\sigma)$ , we have*

$$\|T_\sigma f(x)\|_{L^2(\omega)} \lesssim_{\sigma, \omega} (\mathfrak{T} + \mathfrak{T}^* + \sqrt{A_2(\omega, \sigma)}) \|f\|_{L^2(\sigma)},$$

where

$$A_2(\omega, \sigma) \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\omega(Q)}{|Q|} \frac{\sigma(Q)}{|Q|},$$

# A T1 theorem for doubling measures

Given a measure  $\mu$ , write  $T_\mu(f) \equiv T(f\mu)$ .

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

Assume  $\omega, \sigma$  are doubling measures and  $T$  is a Calderón-Zygmund operator. Then for all functions  $f \in L^2(\sigma)$ , we have

$$\|T_\sigma f(x)\|_{L^2(\omega)} \lesssim_{\sigma, \omega} (\mathfrak{T} + \mathfrak{T}^* + \sqrt{A_2(\omega, \sigma)}) \|f\|_{L^2(\sigma)},$$

where

$$A_2(\omega, \sigma) \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\omega(Q)}{|Q|} \frac{\sigma(Q)}{|Q|},$$
$$\mathfrak{T}^2 \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\int_Q |T_\sigma 1_Q(x)|^2 d\omega(x)}{\sigma(Q)},$$

# A T1 theorem for doubling measures

Given a measure  $\mu$ , write  $T_\mu(f) \equiv T(f\mu)$ .

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

Assume  $\omega, \sigma$  are doubling measures and  $T$  is a Calderón-Zygmund operator. Then for all functions  $f \in L^2(\sigma)$ , we have

$$\|T_\sigma f(x)\|_{L^2(\omega)} \lesssim_{\sigma, \omega} (\mathfrak{T} + \mathfrak{T}^* + \sqrt{A_2(\omega, \sigma)}) \|f\|_{L^2(\sigma)},$$

where

$$A_2(\omega, \sigma) \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\omega(Q)}{|Q|} \frac{\sigma(Q)}{|Q|},$$

$$\mathfrak{T}^2 \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\int_Q |T_\sigma 1_Q(x)|^2 d\omega(x)}{\sigma(Q)},$$

and  $\mathfrak{T}^*$  is defined in a similar fashion (swap  $\sigma$  and  $\omega$  and use operator  $T^*$ ).

# A T1 theorem for doubling measures

Given a measure  $\mu$ , write  $T_\mu(f) \equiv T(f\mu)$ .

Theorem (A. -Sawyer-Uriarte-Tuero, 2021)

Assume  $\omega, \sigma$  are doubling measures and  $T$  is a Calderón-Zygmund operator. Then for all functions  $f \in L^2(\sigma)$ , we have

$$\|T_\sigma f(x)\|_{L^2(\omega)} \lesssim_{\sigma, \omega} (\mathfrak{T} + \mathfrak{T}^* + \sqrt{A_2(\omega, \sigma)}) \|f\|_{L^2(\sigma)},$$

where

$$A_2(\omega, \sigma) \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\omega(Q)}{|Q|} \frac{\sigma(Q)}{|Q|},$$

$$\mathfrak{T}^2 \equiv \sup_{Q \subset \mathbb{R}^n} \frac{\int_Q |T_\sigma 1_Q(x)|^2 d\omega(x)}{\sigma(Q)},$$

and  $\mathfrak{T}^*$  is defined in a similar fashion (swap  $\sigma$  and  $\omega$  and use operator  $T^*$ ).

Note that the statement is only meaningful if  $A_2(\omega, \sigma), \mathfrak{T}, \mathfrak{T}^*$  are all finite.

# List of Tools



# List of Tools

Tools given by the problem:

- $A_2$  (good for separated cubes).
- Testing conditions (good for controlling the singularity).
- Doubling (allows us to make  $\lesssim$  approximations without constants blowing up).

# List of Tools

Tools given by the problem:

- $A_2$  (good for separated cubes).
- Testing conditions (good for controlling the singularity).
- Doubling (allows us to make  $\lesssim$  approximations without constants blowing up).

Tools we bring in to help:

- Haar wavelets.
- Random dyadic grids.
- Stopping time arguments.

# List of Tools

Tools given by the problem:

- $A_2$  (good for separated cubes).
- Testing conditions (good for controlling the singularity).
- Doubling (allows us to make  $\lesssim$  approximations without constants blowing up).

Tools we bring in to help:

- Haar wavelets.
- Random dyadic grids.
- Stopping time arguments.

# Start of proof: a Haar decomposition

## Start of proof: a Haar decomposition

By duality, for all  $f$  and  $g$  with compact support we want to show

$$|\langle T_\sigma f, g \rangle_\omega| \lesssim \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}$$

## Start of proof: a Haar decomposition

By duality, for all  $f$  and  $g$  with compact support we want to show

$$|\langle T_\sigma f, g \rangle_\omega| \lesssim \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}$$

Fix a dyadic grid  $\mathcal{D}$ .

# Start of proof: a Haar decomposition

By duality, for all  $f$  and  $g$  with compact support we want to show

$$|\langle T_\sigma f, g \rangle_\omega| \lesssim \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}$$

Fix a dyadic grid  $\mathcal{D}$ . Expand  $f$  into  $L^2(\sigma)$ -Haar wavelets and  $g$  into  $L^2(\omega)$ -Haar wavelets, i.e. write each as the orthogonal sum

$$f = \sum_{l \in \mathcal{D}} \Delta_l^\sigma f, \quad g = \sum_{l \in \mathcal{D}} \Delta_l^\omega g$$


# Start of proof: a Haar decomposition

By duality, for all  $f$  and  $g$  with compact support we want to show

$$|\langle T_\sigma f, g \rangle_\omega| \lesssim \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}$$

Fix a dyadic grid  $\mathcal{D}$ . Expand  $f$  into  $L^2(\sigma)$ -Haar wavelets and  $g$  into  $L^2(\omega)$ -Haar wavelets, i.e. write each as the orthogonal sum

$$f = \sum_{I \in \mathcal{D}} \Delta_I^\sigma f, \quad g = \sum_{I \in \mathcal{D}} \Delta_I^\omega g$$






# Start of proof: a Haar decomposition

By duality, for all  $f$  and  $g$  with compact support we want to show

$$|\langle T_\sigma f, g \rangle_\omega| \lesssim \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}$$

Fix a dyadic grid  $\mathcal{D}$ . Expand  $f$  into  $L^2(\sigma)$ -Haar wavelets and  $g$  into  $L^2(\omega)$ -Haar wavelets, i.e. write each as the orthogonal sum

$$f = \sum_{I \in \mathcal{D}} \Delta_I^\sigma f, \quad g = \sum_{I \in \mathcal{D}} \Delta_I^\omega g$$



Then we want to show

$$\left| \sum_{I, J \in \mathcal{D}} \langle T_\sigma \Delta_I^\sigma f, \Delta_J^\omega g \rangle_\omega \right| \lesssim \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}.$$

# Why Haar wavelets?

# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

$$\begin{aligned} \left| \int_{(2Q)^c} \int_Q K(x, y) b_Q(y) dy dx \right| &= \left| \int_{(2Q)^c} \int_Q (K(x, y) - K(x, c_Q)) b_Q(y) dy dx \right| \\ &\leq \int_Q \left| \int_{(2Q)^c} K(x, y) - K(x, c_Q) dx \right| |b_Q(y)| dy \approx \int_{(2Q)^c} |\nabla K(x, c_Q)| dx \int_Q |b_Q(y)| dy \end{aligned}$$

# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

$$\begin{aligned} \left| \int_{(2Q)^c} \int_Q K(x, y) b_Q(y) dy dx \right| &= \left| \int_{(2Q)^c} \int_Q (K(x, y) - K(x, c_Q)) b_Q(y) dy dx \right| \\ &\leq \int_Q \left| \int_{(2Q)^c} K(x, y) - K(x, c_Q) dx \right| |b_Q(y)| dy \approx \int_{(2Q)^c} |\nabla K(x, c_Q)| dx \int_Q |b_Q(y)| dy \end{aligned}$$

Same principle with Haar wavelets.

# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

$$\begin{aligned} \left| \int_{(2Q)^c} \int_Q K(x, y) b_Q(y) dy dx \right| &= \left| \int_{(2Q)^c} \int_Q (K(x, y) - K(x, c_Q)) b_Q(y) dy dx \right| \\ &\leq \int_Q \left| \int_{(2Q)^c} K(x, y) - K(x, c_Q) dx \right| |b_Q(y)| dy \approx \int_{(2Q)^c} |\nabla K(x, c_Q)| dx \int_Q |b_Q(y)| dy \end{aligned}$$

Same principle with Haar wavelets. For instance, for any measure  $\nu$

$$|\langle T(1_{(2J)^c} \nu), \Delta_J^\omega g \rangle_\omega| \lesssim P_1(J, 1_{(2J)^c} \nu) \sqrt{\omega(J)} \|\Delta_J^\omega g\|_{L^2(\omega)},$$

# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

$$\begin{aligned} \left| \int_{(2Q)^c} \int_Q K(x, y) b_Q(y) dy dx \right| &= \left| \int_{(2Q)^c} \int_Q (K(x, y) - K(x, c_Q)) b_Q(y) dy dx \right| \\ &\leq \int_Q \left| \int_{(2Q)^c} K(x, y) - K(x, c_Q) dx \right| |b_Q(y)| dy \approx \int_{(2Q)^c} |\nabla K(x, c_Q)| dx \int_Q |b_Q(y)| dy \end{aligned}$$

Same principle with Haar wavelets. For instance, for any measure  $\nu$

$$|\langle T(1_{(2J)^c} \nu), \Delta_J^\omega g \rangle_\omega| \lesssim P_1(J, 1_{(2J)^c} \nu) \sqrt{\omega(J)} \|\Delta_J^\omega g\|_{L^2(\omega)},$$

where  $P_1(J, \nu)$  denotes a Poisson integral with kernel centered and scaled to fit  $J$ , with decay like  $x^{-n-1}$ .

# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

$$\begin{aligned} \left| \int_{(2Q)^c} \int_Q K(x, y) b_Q(y) dy dx \right| &= \left| \int_{(2Q)^c} \int_Q (K(x, y) - K(x, c_Q)) b_Q(y) dy dx \right| \\ &\leq \int_Q \left| \int_{(2Q)^c} K(x, y) - K(x, c_Q) dx \right| |b_Q(y)| dy \approx \int_{(2Q)^c} |\nabla K(x, c_Q)| dx \int_Q |b_Q(y)| dy \end{aligned}$$

Same principle with Haar wavelets. For instance, for any measure  $\nu$

$$|\langle T(1_{(2J)^c} \nu), \Delta_J^\omega g \rangle_\omega| \lesssim P_1(J, 1_{(2J)^c} \nu) \sqrt{\omega(J)} \|\Delta_J^\omega g\|_{L^2(\omega)},$$

where  $P_1(J, \nu)$  denotes a Poisson integral with kernel centered and scaled to fit  $J$ , with decay like  $x^{-n-1}$ .

This type of estimate pervades most of the proof.



# Why Haar wavelets?

Recall the classical CZ decomposition: we have  $T$  act on a “bad” function  $b_Q$  supported on a cube  $Q$  with mean 0, and then exploit cancellation of the kernel via a Hörmander-type condition away from the singularity:

$$\begin{aligned} \left| \int_{(2Q)^c} \int_Q K(x, y) b_Q(y) dy dx \right| &= \left| \int_{(2Q)^c} \int_Q (K(x, y) - K(x, c_Q)) b_Q(y) dy dx \right| \\ &\leq \int_Q \left| \int_{(2Q)^c} K(x, y) - K(x, c_Q) dx \right| |b_Q(y)| dy \approx \int_{(2Q)^c} |\nabla K(x, c_Q)| dx \int_Q |b_Q(y)| dy \end{aligned}$$

Same principle with Haar wavelets. For instance, for any measure  $\nu$

$$|\langle T(1_{(2J)^c} \nu), \Delta_J^\omega g \rangle_\omega| \lesssim P_1(J, 1_{(2J)^c} \nu) \sqrt{\omega(J)} \|\Delta_J^\omega g\|_{L^2(\omega)},$$

where  $P_1(J, \nu)$  denotes a Poisson integral with kernel centered and scaled to fit  $J$ , with decay like  $x^{-n-1}$ .

This type of estimate pervades most of the proof. Note that for each  $P_1(J, 1_{(2J)^c} \nu)$  we have only  $\sqrt{\omega(J)}$ .

## Base case: “flat” measures and the pivotal condition


Using the Haar cancellation condition leads to lots of terms like

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_Q \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$

# Base case: “flat” measures and the pivotal condition

Using the Haar cancellation condition leads to lots of terms like


$$\sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$



## Base case: “flat” measures and the pivotal condition

Using the Haar cancellation condition leads to lots of terms like

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$



Need to make sense of these/simplify.

## Base case: “flat” measures and the pivotal condition

Using the Haar cancellation condition leads to lots of terms like

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_Q \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$



Need to make sense of these/simplify.

Assume (for now)  $\omega, \sigma$  are “flat.”

## Base case: “flat” measures and the pivotal condition

Using the Haar cancellation condition leads to lots of terms like

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$



Need to make sense of these/simplify.


Assume (for now)  $\omega, \sigma$  are “flat.” Then  $\omega, \sigma$  satisfy the Pivotal condition

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r) \lesssim \sum_{r=1}^{\infty} (\sigma(Q_r)/|Q_r|)^2 \omega(Q_r) \lesssim A_2(\omega, \sigma) \sum_{r=1}^{\infty} \sigma(Q_r)$$

## Base case: “flat” measures and the pivotal condition

Using the Haar cancellation condition leads to lots of terms like

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$



Need to make sense of these/simplify.

Assume (for now)  $\omega, \sigma$  are “flat.” Then  $\omega, \sigma$  satisfy the Pivotal condition

$$\begin{aligned} \sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r) &\lesssim \sum_{r=1}^{\infty} (\sigma(Q_r)/|Q_r|)^2 \omega(Q_r) \lesssim A_2(\omega, \sigma) \sum_{r=1}^{\infty} \sigma(Q_r) \\ &\leq A_2 \sigma(Q). \end{aligned}$$

## Base case: “flat” measures and the pivotal condition

Using the Haar cancellation condition leads to lots of terms like

$$\sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r), \text{ where } Q = \sqcup Q_r.$$



Need to make sense of these/simplify.

Assume (for now)  $\omega, \sigma$  are “flat.” Then  $\omega, \sigma$  satisfy the Pivotal condition

$$\begin{aligned} \sum_{r=1}^{\infty} P_1(Q_r, 1_{Q_r} \sigma)^2 \omega(Q_r) &\lesssim \sum_{r=1}^{\infty} (\sigma(Q_r)/|Q_r|)^2 \omega(Q_r) \lesssim A_2(\omega, \sigma) \sum_{r=1}^{\infty} \sigma(Q_r) \\ &\leq A_2 \sigma(Q). \end{aligned}$$

When  $\omega, \sigma$  are “flat,” we can control the above terms by the  $A_2$  condition.



# How (else) do we use $A_2$ ?

## How (else) do we use $A_2$ ?

$A_2$  is good for controlling terms where the cubes have some **separation**:

## How (else) do we use $A_2$ ?

$A_2$  is good for controlling terms where the cubes have some **separation**: indeed, if  $I, J$  have sidelength  $L$  and are distance at least  $L$  apart, then

## How (else) do we use $A_2$ ?

$A_2$  is good for controlling terms where the cubes have some **separation**: indeed, if  $I, J$  have sidelength  $L$  and are distance at least  $L$  apart, then

$$\begin{aligned} |\langle T_\sigma(\Delta_I^\sigma f), \Delta_J^\omega g \rangle_\omega| &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \int_J \int_I \frac{d\sigma(y) d\omega(x)}{|x-y|^n} \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} (\omega(J)\sigma(I))/L^n \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \sqrt{A_2(\omega, \sigma)} \sqrt{\omega(J)} \sqrt{\sigma(I)} \\ &\lesssim \sqrt{A_2} \|\Delta_I^\sigma f\|_{L^2(\sigma)} \|\Delta_J^\omega g\|_{L^2(\omega)}. \end{aligned}$$

## How (else) do we use $A_2$ ?

$A_2$  is good for controlling terms where the cubes have some **separation**: indeed, if  $I, J$  have sidelength  $L$  and are distance at least  $L$  apart, then

$$\begin{aligned} |\langle T_\sigma(\Delta_I^\sigma f), \Delta_J^\omega g \rangle_\omega| &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \int_J \int_I \frac{d\sigma(y) d\omega(x)}{|x - y|^n} \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} (\omega(J)\sigma(I))/L^n \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \sqrt{A_2(\omega, \sigma)} \sqrt{\omega(J)} \sqrt{\sigma(I)} \\ &\lesssim \sqrt{A_2} \|\Delta_I^\sigma f\|_{L^2(\sigma)} \|\Delta_J^\omega g\|_{L^2(\omega)}. \end{aligned}$$

(The last inequality is a property of Haar wavelets).

## How (else) do we use $A_2$ ?

$A_2$  is good for controlling terms where the cubes have some **separation**: indeed, if  $I, J$  have sidelength  $L$  and are distance at least  $L$  apart, then

$$\begin{aligned} |\langle T_\sigma(\Delta_I^\sigma f), \Delta_J^\omega g \rangle_\omega| &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \int_J \int_I \frac{d\sigma(y) d\omega(x)}{|x-y|^n} \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} (\omega(J)\sigma(I))/L^n \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \sqrt{A_2(\omega, \sigma)} \sqrt{\omega(J)} \sqrt{\sigma(I)} \\ &\lesssim \sqrt{A_2} \|\Delta_I^\sigma f\|_{L^2(\sigma)} \|\Delta_J^\omega g\|_{L^2(\omega)}. \end{aligned}$$

(The last inequality is a property of Haar wavelets).

In fact, if we work much harder, we can estimate

$$\left| \sum_{\substack{I, J \in \mathcal{D} \\ I \nsubseteq 3J \text{ and } J \nsubseteq 3I}} \langle T_\sigma(\Delta_I^\sigma f), \Delta_J^\omega g \rangle_\omega \right| \lesssim \sqrt{A_2} \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}.$$

## How (else) do we use $A_2$ ?

$A_2$  is good for controlling terms where the cubes have some **separation**: indeed, if  $I, J$  have sidelength  $L$  and are distance at least  $L$  apart, then

$$\begin{aligned} |\langle T_\sigma(\Delta_I^\sigma f), \Delta_J^\omega g \rangle_\omega| &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \int_J \int_I \frac{d\sigma(y) d\omega(x)}{|x-y|^n} \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} (\omega(J)\sigma(I))/L^n \\ &\leq \|\Delta_I^\sigma f\|_{L^\infty} \|\Delta_J^\omega g\|_{L^\infty} \sqrt{A_2(\omega, \sigma)} \sqrt{\omega(J)} \sqrt{\sigma(I)} \\ &\lesssim \sqrt{A_2} \|\Delta_I^\sigma f\|_{L^2(\sigma)} \|\Delta_J^\omega g\|_{L^2(\omega)}. \end{aligned}$$

(The last inequality is a property of Haar wavelets).

In fact, if we work much harder, we can estimate

$$\left| \sum_{\substack{I, J \in \mathcal{D} \\ I \nsubseteq 3J \text{ and } J \nsubseteq 3I}} \langle T_\sigma(\Delta_I^\sigma f), \Delta_J^\omega g \rangle_\omega \right| \lesssim \sqrt{A_2} \|f\|_{L^2(\sigma)} \|g\|_{L^2(\omega)}.$$

So left with estimating terms where  $I, J$  are overlapping or close.

# What to do when cubes are nearby?



# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?

# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?

A: We don't!

# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?


A: We don't! Given  $f, g$ , choose a random dyadic grid  $\mathcal{D}$  so that there exists  $\rho \gg 1$  such that for all  $K, L$  in the Haar support of  $f, g$ , if  $K \subset L$  and  $\ell(K) \leq 2^{-\rho} \ell(L)$ , then  $K \subseteq L$  ("deeply embedded" as on right below):

# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?

A: We don't! Given  $f, g$ , choose a random dyadic grid  $\mathcal{D}$  so that there exists  $\rho \gg 1$  such that for all  $K, L$  in the Haar support of  $f, g$ , if  $K \subset L$  and  $\ell(K) \leq 2^{-\rho} \ell(L)$ , then  $K \Subset L$  ("deeply embedded" as on right below):




# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?

A: We don't! Given  $f, g$ , choose a random dyadic grid  $\mathcal{D}$  so that there exists  $\rho \gg 1$  such that for all  $K, L$  in the Haar support of  $f, g$ , if  $K \subset L$  and  $\ell(K) \leq 2^{-\rho} \ell(L)$ , then  $K \Subset L$  ("deeply embedded" as on right below):




If cube  $J$  is close to  $I$  and tiny with respect to  $I$ , then have lots of separation  $\Rightarrow$  use  $P_1$  and  $A_2$  to control these terms.

# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?

A: We don't! Given  $f, g$ , choose a random dyadic grid  $\mathcal{D}$  so that there exists  $\rho \gg 1$  such that for all  $K, L$  in the Haar support of  $f, g$ , if  $K \subset L$  and  $\ell(K) \leq 2^{-\rho} \ell(L)$ , then  $K \Subset L$  ("deeply embedded" as on right below):



If cube  $J$  is close to  $I$  and tiny with respect to  $I$ , then have lots of separation  $\Rightarrow$  use  $P_1$  and  $A_2$  to control these terms.


So only have to deal with  $\sum_{I, J : J \Subset I} + \sum_{I, J : \ell(I) \sim \ell(J) \text{ and } J \subset 3I}$

# What to do when cubes are nearby?

Issue: many  $J$ 's are near  $I$ , especially as  $\ell(J) \rightarrow 0$ . Too many to sum, and with no separation can't get a  $P_1$  term.

Q: How do we handle cubes which are near each other?

A: We don't! Given  $f, g$ , choose a random dyadic grid  $\mathcal{D}$  so that there exists  $\rho \gg 1$  such that for all  $K, L$  in the Haar support of  $f, g$ , if  $K \subset L$  and  $\ell(K) \leq 2^{-\rho} \ell(L)$ , then  $K \Subset L$  ("deeply embedded" as on right below):



If cube  $J$  is close to  $I$  and tiny with respect to  $I$ , then have lots of separation  $\Rightarrow$  use  $P_1$  and  $A_2$  to control these terms.

So only have to deal with  $\sum_{I, J : J \Subset I} + \sum_{I, J : \ell(I) \sim \ell(J) \text{ and } J \subset 3I}$  (and the same terms with  $I, J$  swapped, which can be handled similarly by duality).



# Getting rid of cubes of comparable size

# Getting rid of cubes of comparable size

When cubes  $I, J$  are of similar size and are near each other, we estimate this with testing!

## Getting rid of cubes of comparable size

When cubes  $I, J$  are of similar size and are near each other, we estimate this with testing!

Say for instance  $J, I$  are adjacent and of same size:

$$|\langle T_\sigma 1_I, \Delta_J^\omega g \rangle_\omega| \leq \left( \int_J |T_\sigma 1_I(x)|^2 d\omega(x) \right)^{1/2} \|\Delta_J^\omega g\|_{L^2(\omega)}.$$

# Getting rid of cubes of comparable size

When cubes  $I, J$  are of similar size and are near each other, we estimate this with testing!

Say for instance  $J, I$  are adjacent and of same size:

$$|\langle T_\sigma 1_I, \Delta_J^\omega g \rangle_\omega| \leq \left( \int_J |T_\sigma 1_I(x)|^2 d\omega(x) \right)^{1/2} \|\Delta_J^\omega g\|_{L^2(\omega)}.$$

But then  $\int_J |T_\sigma 1_I(x)|^2 d\omega(x) \leq \int_{3I} |T_\sigma 1_I(x)|^2 d\omega(x) \lesssim \int_I |T_\sigma 1_I(x)|^2 d\omega(x) + A_2(\omega, \sigma) + O(1)$  because the measures are doubling.  
We've boiled this down to testing!

## Getting rid of cubes of comparable size

When cubes  $I, J$  are of similar size and are near each other, we estimate this with testing!

Say for instance  $J, I$  are adjacent and of same size:

$$|\langle T_\sigma 1_I, \Delta_J^\omega g \rangle_\omega| \leq \left( \int_J |T_\sigma 1_I(x)|^2 d\omega(x) \right)^{1/2} \|\Delta_J^\omega g\|_{L^2(\omega)}.$$

But then  $\int_J |T_\sigma 1_I(x)|^2 d\omega(x) \leq \int_{3I} |T_\sigma 1_I(x)|^2 d\omega(x) \lesssim \int_I |T_\sigma 1_I(x)|^2 d\omega(x) + A_2(\omega, \sigma) + O(1)$  because the measures are doubling.

We've boiled this down to testing!

If the cubes  $I, J$  outright overlap, then even easier!

# Getting rid of cubes of comparable size

When cubes  $I, J$  are of similar size and are near each other, we estimate this with testing!

Say for instance  $J, I$  are adjacent and of same size:

$$|\langle T_\sigma 1_I, \Delta_J^\omega g \rangle_\omega| \leq \left( \int_J |T_\sigma 1_I(x)|^2 d\omega(x) \right)^{1/2} \|\Delta_J^\omega g\|_{L^2(\omega)}.$$

But then  $\int_J |T_\sigma 1_I(x)|^2 d\omega(x) \leq \int_{3I} |T_\sigma 1_I(x)|^2 d\omega(x) \lesssim \int_I |T_\sigma 1_I(x)|^2 d\omega(x) + A_2(\omega, \sigma) + O(1)$  because the measures are doubling.

We've boiled this down to testing!

If the cubes  $I, J$  outright overlap, then even easier!

So now we only have to bound

$$\left| \sum_{I \in \mathcal{D}} \sum_{\substack{J \in \mathcal{D} \\ J \subseteq I}} \langle T_\sigma \Delta_J^\sigma f, \Delta_J^\omega g \rangle_\omega \right|$$

# A stopping time argument glossed over

# A stopping time argument glossed over

The rest of the proof here uses lots more deep mathematics, using both tools we both have and haven't discussed yet:



# A stopping time argument glossed over

The rest of the proof here uses lots more deep mathematics, using both tools we both have and haven't discussed yet:

- A stopping time argument

# A stopping time argument glossed over

The rest of the proof here uses lots more deep mathematics, using both tools we both have and haven't discussed yet:

- A stopping time argument ( $J \subseteq I$  means that we can partition our cubes into “chunks” that don't interact too much with each other).

# A stopping time argument glossed over

The rest of the proof here uses lots more deep mathematics, using both tools we both have and haven't discussed yet:

- A stopping time argument ( $J \subseteq I$  means that we can partition our cubes into “chunks” that don't interact too much with each other).
- Haar wavelets have a nice telescoping property.

# A stopping time argument glossed over

The rest of the proof here uses lots more deep mathematics, using both tools we both have and haven't discussed yet:

- A stopping time argument ( $J \subseteq I$  means that we can partition our cubes into “chunks” that don't interact too much with each other).
- Haar wavelets have a nice telescoping property.
- Use of the Pivotal Condition.

# A stopping time argument glossed over

The rest of the proof here uses lots more deep mathematics, using both tools we both have and haven't discussed yet:

- A stopping time argument ( $J \subseteq I$  means that we can partition our cubes into “chunks” that don't interact too much with each other).
- Haar wavelets have a nice telescoping property.
- Use of the Pivotal Condition.

But I don't have time to discuss :(

# What about arbitrary doubling measures?

# What about arbitrary doubling measures?

Recall we assumed the measures  $\omega, \sigma$  were “flat.”

# What about arbitrary doubling measures?

Recall we assumed the measures  $\omega, \sigma$  were “flat.”

Arbitrary doubling measures don't satisfy the pivotal condition with  $P_1 \dots$



# What about arbitrary doubling measures?

Recall we assumed the measures  $\omega, \sigma$  were “flat.”

Arbitrary doubling measures don't satisfy the pivotal condition with  $P_1 \dots$

But they do satisfy a Pivotal condition with  $P_\kappa$ , a Poisson integral with kernel decay like  $x^{-n-\kappa}$  for some  $\kappa$  sufficiently large!

# What about arbitrary doubling measures?

Recall we assumed the measures  $\omega, \sigma$  were “flat.”

Arbitrary doubling measures don't satisfy the pivotal condition with  $P_1 \dots$

But they do satisfy a Pivotal condition with  $P_\kappa$ , a Poisson integral with kernel decay like  $x^{-n-\kappa}$  for some  $\kappa$  sufficiently large!

We get

$$\sum_{r=1}^{\infty} P_\kappa(Q_r, 1_Q \sigma)^2 \omega(Q_r) \lesssim A_2(\omega, \sigma) \sigma(Q).$$

# How does the proof skeleton change?

# How does the proof skeleton change?

Recall we originally got the  $P_1$  terms by saying

$$\int_{(2Q)^c} \int K(x, y) b_Q(y) dy dx \approx \left| \int_{(2Q)^c} \nabla K(x, c_Q) dx \right| \int |b_Q(y)| dy$$

# How does the proof skeleton change?

Recall we originally got the  $P_1$  terms by saying

$$\int_{(2Q)^c} \int K(x, y) b_Q(y) dy dx \approx \left| \int_{(2Q)^c} \nabla K(x, c_Q) dx \right| \int |b_Q(y)| dy$$

To prove the analogous estimate with  $P_\kappa$  for  $\kappa > 1$ , you need to get

$\int \nabla^\kappa K$ . To get this, we must subtract the  $(\kappa - 1)$ st Taylor approximation of  $K$  from itself.

# How does the proof skeleton change?

Recall we originally got the  $P_1$  terms by saying

$$\int_{(2Q)^c} \int K(x, y) b_Q(y) dy dx \approx \left| \int_{(2Q)^c} \nabla K(x, c_Q) dx \right| \int |b_Q(y)| dy$$

To prove the analogous estimate with  $P_\kappa$  for  $\kappa > 1$ , you need to get  $\int \nabla^\kappa K$ . To get this, we must subtract the  $(\kappa - 1)$ st Taylor approximation of  $K$  from itself.

To do this, you need to consider functions  $b_Q$  to not just have mean 0, but rather  $\int y^d b_Q(y) dy = 0$  for  $d = 0, \dots, \kappa - 1$ . Need the same properties of  $\Delta_I^\omega g$ .

# Cue Alpert Wavelets!

# Cue Alpert Wavelets!

Let  $\mu$  be a measure. A  $\kappa$ -Alpert wavelet with support  $I$ , say  $h_{I;\kappa}$ , is piecewise a polynomial of degree  $\leq \kappa - 1$  on each dyadic child of  $I$  ( $I^-$ ,  $I^+$  in 1-dimension), such that

$$\int x^m h_{I;\kappa}(x) d\mu(x) = 0, \text{ for } m = 0, \dots, \kappa - 1.$$



# Cue Alpert Wavelets!

Let  $\mu$  be a measure. A  $\kappa$ -Alpert wavelet with support  $I$ , say  $h_{I;\kappa}$ , is piecewise a polynomial of degree  $\leq \kappa - 1$  on each dyadic child of  $I$  ( $I^-$ ,  $I^+$  in 1-dimension), such that

$$\int x^m h_{I;\kappa}(x) d\mu(x) = 0, \text{ for } m = 0, \dots, \kappa - 1.$$

i.e. Alpert wavelets are piecewise polynomials of degree strictly less than  $\kappa$  whose first  $\kappa$  moments vanish.

# Cue Alpert Wavelets!

Let  $\mu$  be a measure. A  $\kappa$ -Alpert wavelet with support  $I$ , say  $h_{I;\kappa}$ , is piecewise a polynomial of degree  $\leq \kappa - 1$  on each dyadic child of  $I$  ( $I^-$ ,  $I^+$  in 1-dimension), such that

$$\int x^m h_{I;\kappa}(x) d\mu(x) = 0, \text{ for } m = 0, \dots, \kappa - 1.$$

i.e. Alpert wavelets are piecewise polynomials of degree strictly less than  $\kappa$  whose first  $\kappa$  moments vanish. *Note:  $\kappa = 1$  is exactly Haar!*

# Cue Alpert Wavelets!

Let  $\mu$  be a measure. A  $\kappa$ -Alpert wavelet with support  $I$ , say  $h_{I;\kappa}$ , is piecewise a polynomial of degree  $\leq \kappa - 1$  on each dyadic child of  $I$  ( $I^-$ ,  $I^+$  in 1-dimension), such that

$$\int x^m h_{I;\kappa}(x) d\mu(x) = 0, \text{ for } m = 0, \dots, \kappa - 1.$$


i.e. Alpert wavelets are piecewise polynomials of degree strictly less than  $\kappa$  whose first  $\kappa$  moments vanish. *Note:  $\kappa = 1$  is exactly Haar!*

# Cue Alpert Wavelets!

Let  $\mu$  be a measure. A  $\kappa$ -Alpert wavelet with support  $I$ , say  $h_{I;\kappa}$ , is piecewise a polynomial of degree  $\leq \kappa - 1$  on each dyadic child of  $I$  ( $I^-$ ,  $I^+$  in 1-dimension), such that

$$\int x^m h_{I;\kappa}(x) d\mu(x) = 0, \text{ for } m = 0, \dots, \kappa - 1.$$

i.e. Alpert wavelets are piecewise polynomials of degree strictly less than  $\kappa$  whose first  $\kappa$  moments vanish. *Note:  $\kappa = 1$  is exactly Haar!*




# Cue Alpert Wavelets!

Let  $\mu$  be a measure. A  $\kappa$ -Alpert wavelet with support  $I$ , say  $h_{I;\kappa}$ , is piecewise a polynomial of degree  $\leq \kappa - 1$  on each dyadic child of  $I$  ( $I^-$ ,  $I^+$  in 1-dimension), such that

$$\int x^m h_{I;\kappa}(x) d\mu(x) = 0, \text{ for } m = 0, \dots, \kappa - 1.$$

i.e. Alpert wavelets are piecewise polynomials of degree strictly less than  $\kappa$  whose first  $\kappa$  moments vanish. *Note:  $\kappa = 1$  is exactly Haar!*



So to accomodate arbitrary doubling constants and the  $P_\kappa$ , we decompose  $f$  into  $L^2(\sigma)$ -Alpert wavelets:  $f = \sum_{I \in \mathcal{D}} \Delta_{I;\kappa}^\sigma f$  (and similarly for  $g$ ).

# Final changes in the proof to accomodate Alpert wavelets

# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues)...

# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues). . . Except for a step of the proof following the stopping time argument.



# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues). . . Except for a step of the proof following the stopping time argument.

In Haar case, we could write  $1_{I-}\Delta_I^\sigma f = c1_{I-}$  and so

$$|\langle T1_{I-}\Delta_I^\sigma f, \Delta_J^\omega g \rangle_\omega| \leq |c| |\langle T1_{I-}, \Delta_J^\omega g \rangle_\omega|.$$

Can't do this in the Alpert wavelet case, since can't pull polynomials out of operators.

# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues)... Except for a step of the proof following the stopping time argument.

In Haar case, we could write  $1_I \Delta_I^\sigma f = c 1_I$  and so

$$|\langle T 1_I \Delta_I^\sigma f, \Delta_J^\omega g \rangle_\omega| \leq |c| |\langle T 1_I, \Delta_J^\omega g \rangle_\omega|.$$

Can't do this in the Alpert wavelet case, since can't pull polynomials out of operators.

But commuting polynomials with CZO's is actually nice! For example, the Hilbert transform  $H$  with kernel  $K(x, y) = \frac{1}{x-y}$ , we have  $[H, x]f$  equals

# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues). . . Except for a step of the proof following the stopping time argument.

In Haar case, we could write  $1_{I-}\Delta_I^\sigma f = c1_{I-}$  and so

$$|\langle T1_{I-}\Delta_I^\sigma f, \Delta_J^\omega g \rangle_\omega| \leq |c| |\langle T1_{I-}, \Delta_J^\omega g \rangle_\omega|.$$

Can't do this in the Alpert wavelet case, since can't pull polynomials out of operators.

But commuting polynomials with CZO's is actually nice! For example, the Hilbert transform  $H$  with kernel  $K(x, y) = \frac{1}{x-y}$ , we have  $[H, x]f$  equals

$$H(xf) - xHf = \int \frac{yf(y)}{x-y} dy - x \int \frac{f(y)}{x-y} dy = \int f(y) \frac{y-x}{x-y} dy = - \int f(y)$$

# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues). . . Except for a step of the proof following the stopping time argument.

In Haar case, we could write  $1_{I-}\Delta_I^\sigma f = c1_{I-}$  and so

$$|\langle T1_{I-}\Delta_I^\sigma f, \Delta_J^\omega g \rangle_\omega| \leq |c| |\langle T1_{I-}, \Delta_J^\omega g \rangle_\omega|.$$

Can't do this in the Alpert wavelet case, since can't pull polynomials out of operators.

But commuting polynomials with CZO's is actually nice! For example, the Hilbert transform  $H$  with kernel  $K(x, y) = \frac{1}{x-y}$ , we have  $[H, x]f$  equals

$$H(xf) - xHf = \int \frac{yf(y)}{x-y} dy - x \int \frac{f(y)}{x-y} dy = \int f(y) \frac{y-x}{x-y} dy = - \int f(y)$$

Since you make the singularity of the CZO more regular, you can in fact just control the commutator term by an  $\sqrt{A_2} \|f\|_{L^2(\sigma)} \|g\|_{L^2(\sigma)}$ .

# Final changes in the proof to accomodate Alpert wavelets

Most of what I mentioned previously still works for Alpert wavelets (with straightforward analogues). . . Except for a step of the proof following the stopping time argument.

In Haar case, we could write  $1_{I-}\Delta_I^\sigma f = c1_{I-}$  and so

$$|\langle T1_{I-}\Delta_I^\sigma f, \Delta_J^\omega g \rangle_\omega| \leq |c| |\langle T1_{I-}, \Delta_J^\omega g \rangle_\omega|.$$

Can't do this in the Alpert wavelet case, since can't pull polynomials out of operators.

But commuting polynomials with CZO's is actually nice! For example, the Hilbert transform  $H$  with kernel  $K(x, y) = \frac{1}{x-y}$ , we have  $[H, x]f$  equals

$$H(xf) - xHf = \int \frac{yf(y)}{x-y} dy - x \int \frac{f(y)}{x-y} dy = \int f(y) \frac{y-x}{x-y} dy = - \int f(y)$$

Since you make the singularity of the CZO more regular, you can in fact just control the commutator term by an  $\sqrt{A_2} \|f\|_{L^2(\sigma)} \|g\|_{L^2(\sigma)}$ . (In particular, no testing needed!)

*Thank you for Listening!*