

Esempio di verifica

$$\forall \varepsilon \in \mathbb{R} |f(x+\varepsilon) - f(x)| < \mathscr{S}$$


$$A\widehat{B}C = 120^{\circ} - \int_{x=0}^{\infty} \frac{1}{x^2} + \binom{5}{2} - \sum_{i=1}^{N} \frac{i(i+1)}{2} - \overline{x}$$

 \Box °Ce

€



Figura 1: Logo della scuola 1



Logo della scuola 2

$$\begin{array}{ccc}
A1 & A2 \\
B12 & AB3
\end{array}$$

Prova in verde

$$2x^2 - 6 = 0$$
$$2(x - \sqrt{3})(x + \sqrt{3}) = 0$$

prova prova prova prova

abcdefghijklmnopqstuvwxyz

abcdefghijklmnopqrstuvwxyz

ABCDEFGHIJKLMNOPQRSTUVWXYZ

ABCDEFGHIJKLMNOPQRSTUVWXYZ

ABCDEFGHIJKLMNOPQRSTUVWXYZ1

$$\operatorname{Nm} \cdot \mathbf{s}^{-2} \Omega \xrightarrow{1c\varsigma} \xrightarrow{1c\varsigma} \qquad \frac{\mathbf{m}}{\mathbf{s}^2} \quad \Longrightarrow \quad \overrightarrow{v} \neq \overrightarrow{v_1} \binom{5+2}{2} = 10 \qquad \begin{pmatrix} 1\\3\\6 \end{pmatrix} \quad \Longrightarrow \quad A \widehat{B} C$$

Consideriamo i triangoli ABC e A'B'C':

$$\left. \begin{array}{l} AB \cong A'B' \\ BC \cong B'C' \\ CA \cong C'A' \end{array} \right\} \stackrel{3^{\circ} \text{cc}}{\Longrightarrow} ABC \cong A'B'C'$$