

ATLAS activities at Sydney

Carl Suster, Shyam Balaji, Kevin Varvell, Bruce Yabsley

The University of Sydney

May 3, 2019



THE UNIVERSITY OF
SYDNEY

LHC Page1 Fill: 7495 E: 0 Z GeV t(SB): 00:00:00 02-05-19 09:40:12



SHUTDOWN: NO BEAM

BIS status and SMP flags		B1	B2
Comments (25-Apr-2019 04:57:28) LS2	Link Status of Beam Permits	Except	Except
	Global Beam Permit	Except	Except
	Setup Beam	false	false
	Beam Presence	false	false
	Moveable Devices Allowed In	false	false
	Stable Beams	false	false
AFS: 75_150ns_733Pb_733_702_468_42bpi_20inj		PM Status B1	PM Status B2
		ENABLED	ENABLED

LHC long-term schedule



The ATLAS pp datasets in run 2



- Supersymmetry searches
- Exotics searches
- Higgs and diboson searches
- Higgs physics
- Standard Model
- Top physics
- Heavy ion physics
- B physics and light states
- Physics modelling


+ hardware, upgrades, operations, organisation, ...

- Supersymmetry searches
- Exotics searches
- Higgs and diboson searches – search for a heavy neutral Higgs
- Higgs physics
- Standard Model
- Top physics – single top tW production, AIDA
- Heavy ion physics
- B physics and light states – quarkonium production spectroscopy
- Physics modelling


+ hardware, upgrades, operations, organisation, ...

Search for a heavy neutral Higgs [Shyam]


- 2HDM has 5 Higgs bosons after EWSB: h and H (CP-even), A (CP-odd), and H^\pm (charged).
- For 2HDM EW baryogenesis we need $125 \text{ GeV} < m_H < m_A \lesssim 800 \text{ GeV}$.
- Search for $A \rightarrow ZH$ with $Z \rightarrow \ell^+ \ell^-$ (clean) and $H \rightarrow b\bar{b}$ (high branching fraction).




gluon-gluon fusion




associated $b\bar{b}$




Phys. Lett. B 783 (2018)



- Select events with m_{bb} in a window around m_H .
- Then scan over m_A , searching for resonances in $m_{\ell\ell bb}$.
- Main backgrounds are Z +jets and $t\bar{t}/tW$.
- Scan over mass points ensuring $m_A - m_H \geq 100 \text{ GeV}$.






- [Shyam](#) is involved in the update of this search to the full dataset:
 - optimising the event selection using machine learning, and
 - signal interpolation.
- Recent [improvements in \$b\$ -jet tagging](#) will help the analysis.
- There will be a focus on making the results [re-interpretable](#).


-
- Shyam is also involved in improving photon reconstruction at ATLAS.
 - He is also pursuing several [theory](#) projects:
 - B physics: SU(4) unified models to explain the $R_{K^{(*)}}$ and $R_{D^{(*)}}$ anomalies,
 - Higgs portal dark matter models, and
 - simplified dark matter models.

Single top tW production [Carl]


- Top quark production at the LHC is predominantly via $t\bar{t}$.
- Rarer **weak force** processes produce top quarks singly, proportional to $|V_{tb}|$.
- The tW channel was first **observed in run 1** of the LHC at ATLAS and CMS.
- We use events with exactly two oppositely-charged electrons or muons (**dilepton**).



	t -channel	tW	s -channel
Branching fraction	73%	24%	3%
<i>Sensitive to...</i>			
4-fermion operators	✓	×	✓
tWb vertex corrections	✓	✓	✓





$$\sigma_{tW} = 71.7 \pm 1.8 (\text{scale}) \pm 3.4 (\text{PDF}) \text{ pb} \quad (\text{NNLO+NNLL})$$




EPJC 78 (2018) 186

- First measurement of the **shape** of tW production.
- Differential with respect to 6 quantities.




An inclusive dilepton analysis (LHC run 1)

- Our involvement in top physics began as a collaboration with Duke (Mark Kruse).
- Simultaneous measurement of $t\bar{t}$, W^+W^- and $Z/\gamma^* \rightarrow \tau\tau$: Phys. Rev. D 91 (2015) 052005
- We've been planning to resume this analysis, subject to funding for a new postdoc.



NLO



NNLO