

# Exam 2 - Newton - KEY

Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

Class Number: PHYS 1061 H002 Instructor: Michael Haas

Name: \_\_\_\_\_

Please read and follow all instructions carefully. Use the back of the sheet if necessary.

Definitions: (incompressible: not able to be compressed, incomprehensible: not able to be understood; not intelligible.)

Score: \_\_\_\_\_ / 42 + Bonus: \_\_\_\_\_ / 6 = Total: \_\_\_\_\_ / 42 || Final: \_\_\_\_\_ % -> [A, B, C, D, F]

## Problem #1 (3 points)

A truck with mass 12.3Mg collides with a bug with mass 0.432g. How does the magnitude of the bug's force on the truck compare to the magnitude of the truck's force on the bug?

- A. The bug's force on the truck is greater than the truck's force on the bug.
- B. The truck's force on the bug is greater than the bug's force on the truck.
- C. **The two forces are equal in magnitude.**
- D. It is impossible to determine without more information.

## Problem #2 (3 points)

A 10.0kg box is at rest on an inclined ramp. Which of the following correctly describes the magnitude of force due to static friction?

- A.  $|\vec{f}_s| = |\vec{W}|$
- B.  $|\vec{f}_s| = |\vec{N}|$
- C.  $|\vec{f}_s| = |\vec{W}_\perp|$
- D.  **$|\vec{f}_s| = |\vec{W}_\parallel|$**
- E.  $|\vec{f}_s| = 0$

## Problem #3 (3 points)

A baby is pushing on a walker with a force of 10.0N at an angle of 30.0° above the horizontal. What is the magnitude of the horizontal component of the force he is applying?

- A.  **$F_x = 10.0\text{N} \cos(30.0^\circ)$**
- B.  $F_x = 10.0\text{N} \sin(30.0^\circ)$
- C.  $F_x = 10.0\text{N}$
- D.  $F_x = 10.0\text{N} \tan(30.0^\circ)$

### Problem #4 (3 points)

Which of the following is not one of Newton's laws of motion?

- A.  $\vec{F} = m\vec{a}$
- B. When in equilibrium:  $\vec{F}_{\text{net}} = 0$
- C.  $f = \mu N$
- D.  $\vec{F}_{AB} = -\vec{F}_{BA}$


### Problem #5 (3 points)

Arthur Dent is in an elevator on the planet Squornshellous Zeta where the gravitational acceleration is  $g_{\text{Squornshellous}} = 5.70 \frac{\text{m}}{\text{s}^2}$ . The elevator is accelerating upwards at  $2.00 \frac{\text{m}}{\text{s}^2}$ . What is the magnitude of the normal force on Arthur if his mass is 70.0kg?

- A.  $N = mg = 399. \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$
- B.  $N = mg + ma = 539. \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$
- C.  $N = mg - ma = 259. \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$
- D.  $N = ma = 140. \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$

### Problem #6 (3 points)

Boudreaux is moving a box along a horizontal surface, but he can't decide if he should push the box by exerting force  $\vec{F}_1$ , or pull the box by exerting force  $\vec{F}_2$ . Assuming both forces are at the same angle from the horizontal, which one gives the least friction between the box and the surface?



- A.  $\vec{F}_1$
- B.  $\vec{F}_2$
- C. Both forces result in the same friction.
- D. It is impossible to determine without more information.

### Problem #7 (3 points)

An object is hanging by a string from the ceiling of an elevator. The elevator is slowing down while moving upward. What can be said about the magnitude of the tension in the string?

- A. **The magnitude of the tension is less than the magnitude of the resting weight of the object.**
- B. The tension in the string cannot be determined without knowing the speed of the elevator.
- C. The magnitude of the tension is greater than the magnitude of the resting weight of the object.
- D. The tension in the string is zero.
- E. The magnitude of the tension in the string is equal to the magnitude of the resting weight of the object.

### Bonus #8 (3 points) **Severus Snape, $3.20e + 05$ , 1, 1, 2, 3, 5 or 0, 1, 1, 2, 3**


Bonus est omnis divisa in partes tres: Who is the half blood prince? How many centimeters is  $3.20\text{km}$ ? Write out the first 5 numbers in the Fibonacci sequence.

### Problem #9 (9 points)

Baby Godric (who currently has a mass of 9.98kg) is being held up by his mom as in the picture. Treat each of his arms as ideal tension ropes.

#### Question #1 (3 points)

Draw a free body diagram for Godric showing all forces acting on him.



#### Question #2 (3 points)

What is the tension in his right arm?

$$T_L + T_R = mg = 9.98\text{kg} * 9.81 \frac{\text{m}}{\text{s}^2}$$

$$T_L = 97.9 \frac{\text{kg}\cdot\text{m}}{\text{s}^2} - T_R$$

$$\text{Notice: } T_L = T_R$$

$$T_L = T_R = \frac{mg}{2} = 49.0 \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$$

#### Question #3 (3 points)

What would the tension in the right arm be if they were in an elevator accelerating upward at  $2.00 \frac{\text{m}}{\text{s}^2}$ ?

$$T_L + T_R = m(g + a) = 9.98\text{kg} * (9.81 \frac{\text{m}}{\text{s}^2} + 2.00 \frac{\text{m}}{\text{s}^2})$$

$$T_L = 118. \frac{\text{kg}\cdot\text{m}}{\text{s}^2} - T_R$$

$$\text{Notice: } T_L = T_R$$

$$T_L = T_R = \frac{m(g+a)}{2} = 58.9 \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$$

**Bonus #10 (3 points)** Gravity on Mars is  $3.70 \frac{\text{m}}{\text{s}^2}$ . (from test instructions),

$$T_1 = T_2 = \frac{mg_{\text{mars}}}{2} = 18.5 \frac{\text{kg}\cdot\text{m}}{\text{s}^2}$$



What would the tension in the left arm be if they were on the surface of Mars?

### Problem #11 (12 points)

Two boxes are connected by a string (assume massless, frictionless string that never breaks) over a pulley. Box 1 has mass  $10.0\text{kg}$  and is on a table where the coefficient of kinetic friction is  $\mu_k = 0.200$ . Box 2 has mass  $5.00\text{kg}$  and is hanging from the string. The system is in motion (meaning  $v_{x\text{Box1}} \neq 0$ ).

#### Question #1 (3 points)

Draw a free body diagram for each box showing all forces acting on it.



#### Question #2 (3 points)

What is the acceleration of the system? (only give algebraic answer, no numbers yet)

$$\sum_{\text{ext}} F_x = m_1 a_x = T_1 - f_k \text{ and } \sum_{\text{ext}} F_y = m_2 a_y = m_2 g - T_2$$

$$a_x = a_y = a \text{ and } T_1 = T_2 \implies T = f_k + m_1 a = T = m_2 g - m_2 a$$

$$m_1 a + m_2 a = m_2 g - f_k \implies a = \frac{m_2 g - f_k}{m_1 + m_2}$$

Notice this implies the acceleration is positive when the block on the table is moving to the left.

#### Question #3 (3 points)

What is the acceleration of the system? (give a number with units, 3 significant figures)

$$a = \frac{m_2 g - f_k}{m_1 + m_2}, f_k = \mu_k N, N = m_1 g, f_k = \mu_k m_1 g$$

$$a = \frac{(5.00\text{kg})(9.81 \frac{\text{m}}{\text{s}^2}) - (0.200)(10.0\text{kg})(9.81 \frac{\text{m}}{\text{s}^2})}{10.0\text{kg} + 5.00\text{kg}}$$

$$a = 1.96 \frac{\text{m}}{\text{s}^2}$$

#### Question #4 (3 points)

What is the tension in the string?

$$T = f_k + m_1 a = \mu_k m_1 g + m_1 a = (0.200)(10.0\text{kg})(9.81 \frac{\text{m}}{\text{s}^2}) + (10.0\text{kg})(1.96 \frac{\text{m}}{\text{s}^2})$$

$$T = 39.2 \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$$