

Exam 1 - Units, Vectors, and Kinematics

Date: ____ / ____ / ____

Class Number: PHYS 1061 H002 Instructor: Michael Haas

Name: _____

Please read and follow all instructions carefully. Use the back of the sheet if necessary. Mars has $g = 3.72 \frac{\text{m}}{\text{s}^2}$.

Score: _____ / 42 + Bonus: _____ / 6 = Total: _____ / 42 || Final: _____ % -> [A, B, C, D, F]

Problem #1 (3 points)

A truck with mass 12.3Mg collides with a bug with mass 0.432g. How does the magnitude of the bug's force on the truck compare to the magnitude of the truck's force on the bug?

- A. The bug's force on the truck is greater than the truck's force on the bug.
- B. The truck's force on the bug is greater than the bug's force on the truck.
- C. The two forces are equal in magnitude.
- D. It is impossible to determine without more information.

Problem #2 (3 points)

A 10.0kg box is at rest on an inclined ramp. Which of the following correctly describes the magnitude of force due to static friction?

- A. $|\vec{f}_s| = |\vec{W}|$
- B. $|\vec{f}_s| = |\vec{N}|$
- C. $|\vec{f}_s| = |\vec{W}_\perp|$
- D. $|\vec{f}_s| = |\vec{W}_\parallel|$
- E. $|\vec{f}_s| = 0$

Problem #3 (3 points)

A baby is pushing on a walker with a force of 10.0N at an angle of 30.0° above the horizontal. What is the magnitude of the horizontal component of the force he is applying?

- A. $F_x = 10.0\text{N} \cos(30.0^\circ)$
- B. $F_x = 10.0\text{N} \sin(30.0^\circ)$
- C. $F_x = 10.0\text{N}$
- D. $F_x = 10.0\text{N} \tan(30.0^\circ)$

Problem #4 (3 points)

Which of the following is not one of Newton's laws of motion?

- A. $\vec{F} = m\vec{a}$
- B. When in equilibrium: $\vec{F}_{\text{net}} = 0$
- C. $f = \mu N$
- D. $\vec{F}_{\text{AB}} = -\vec{F}_{\text{BA}}$


Problem #5 (3 points)

Arthur Dent is in an elevator on the planet Squornshellous Zeta where the gravitational acceleration is $g_{\text{Squornshellous}} = 5.70 \frac{\text{m}}{\text{s}^2}$. The elevator is accelerating upwards at $2.00 \frac{\text{m}}{\text{s}^2}$. What is the magnitude of the normal force on Arthur if his mass is 70.0kg ?

- A. $N = mg = 399. \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$
- B. $N = mg + ma = 539. \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$
- C. $N = mg - ma = 259. \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$
- D. $N = ma = 140. \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$

Problem #6 (3 points)

Boudreaux is moving a box along a horizontal surface, but he can't decide if he should push the box by exerting force \vec{F}_1 , or pull the box by exerting force \vec{F}_2 . Assuming both forces are at the same angle from the horizontal, which one gives the least friction between the box and the surface?



- A. \vec{F}_1
- B. \vec{F}_2
- C. Both forces result in the same friction.
- D. It is impossible to determine without more information.

Problem #7 (3 points)

An object is hanging by a string from the ceiling of an elevator. The elevator is slowing down while moving upward. What can be said about the magnitude of the tension in the string?

- A. The magnitude of the tension is less than the magnitude of the resting weight of the object.
- B. The tension in the string cannot be determined without knowing the speed of the elevator.
- C. The magnitude of the tension is greater than the magnitude of the resting weight of the object.
- D. The tension in the string is zero.
- E. The magnitude of the tension in the string is equal to the magnitude of the resting weight of the object.

Bonus #8 (3 points)

Bonus est omnis divisa in partes tres: Who is the half blood prince? How many centimeters is 3.20km ? Write out the first 5 numbers in the Fibonacci sequence.

Problem #9 (9 points)

Baby Godric (who currently has a mass of 9.98kg) is being held up by his mom as in the picture. Treat each of his arms as ideal tension ropes.


Question #1 (3 points)

Draw a free body diagram for Godric showing all forces acting on him.



Question #2 (3 points)

What is the tension in his right arm?



Question #3 (3 points)

What would the tension in the right arm be if they were in an elevator accelerating upward at $2.00 \frac{\text{m}}{\text{s}^2}$?



Bonus #10 (3 points)


What would the tension in the left arm be if they were on the surface of Mars?

Problem #11 (12 points)

Two boxes are connected by a string (assume massless, frictionless string that never breaks) over a pulley. Box 1 has mass 10.0kg and is on a table where the coefficient of kinetic friction is $\mu_k = 0.200$. Box 2 has mass 5.00kg and is hanging from the string. The system is in motion (meaning $v_{x_{\text{Box1}}} \neq 0$).

Question #1 (3 points)

Draw a free body diagram for each box showing all forces acting on it.



Question #2 (3 points)

What is the acceleration of the system? (only give algebraic answer, no numbers yet)



Question #3 (3 points)

What is the acceleration of the system? (give a number with units, 3 significant figures)



Question #4 (3 points)

What is the tension in the string?

