

# Mathematics Journal

Michael C.R. Byrd Jr.

July 12, 2018

# Contents

<b>I</b>	<b>Calculus</b>	<b>2</b>
<b>1</b>	<b>Derivatives</b>	<b>3</b>
1.1	Rates of Change . . . . .	3
<b>II</b>	<b>Abstract Algebra</b>	<b>4</b>
<b>1</b>	<b>Groups</b>	<b>5</b>
1.1	Introduction - Groups . . . . .	5
<b>III</b>	<b>Combinatorics</b>	<b>6</b>
<b>1</b>	<b>Graphs</b>	<b>7</b>
1.1	Introduction - Graphs . . . . .	7
<b>IV</b>	<b>Quantum Computation</b>	<b>8</b>
<b>1</b>	<b>Quantum Computation - Intro</b>	<b>9</b>
1.1	Quantum Bits . . . . .	9
1.1.1	Multiple Qubits . . . . .	11

Part I

Calculus

# Chapter 1

## Derivatives

This is a chapter on Derivatives.

### 1.1 Rates of Change

This is the section on Rates of Change.

**Part II**

**Abstract Algebra**

# Chapter 1

# Groups

This is the chapter on Groups.

## 1.1 Introduction - Groups

**Part III**

**Combinatorics**

# Chapter 1

# Graphs

This is the chapter on Graphs.

## 1.1 Introduction - Graphs



## Part IV

# Quantum Computation

# Chapter 1

## Quantum Computation - Intro

This chapter will consist of quantum bits and gates.

### 1.1 Quantum Bits

Classical computation is built upon *bits*. In the world of Quantum computation and quantum information, we have an analogous concept known as a *quantum bit*, or *qubit*.

**Definition 1** (Classical Bit). A *classical bit* is a unit of information that takes on one of two states, commonly denoted 0 or 1. The term bit is a portmanteau of the term binary digit.

**Definition 2** (Quantum Bit). A quantum bit, also known as a qubit has a state. Two possible states for a qubit are  $|0\rangle$  and  $|1\rangle$ . The notation “ $|\rangle$ ” is known as *Dirac notation* named after Paul Dirac.

The main difference between bits and qubits is that a qubit can take on a state other than  $|0\rangle$  or  $|1\rangle$ . A qubit can take the form that is a linear combination of these two states, this is referred to as a superposition.

An example of a superposition is the following

$$|\psi\rangle = \alpha |0\rangle + \beta |1\rangle, \quad (1.1)$$

where  $\alpha, \beta \in \mathbb{C}$ .

We can consider the state of a qubit as a vector in  $\mathbb{C}^2$ . The states  $|0\rangle$  and  $|1\rangle$  are known as computation basis states and form an orthogonal basis for this vector space.

While a bit's state can be determined by examination, the same is not true for a qubit. That is, one cannot determine the values  $\alpha$  and  $\beta$ . When we measure a qubit we either get the result 0, with probability  $|\alpha|^2$ , or the result 1, with probability  $|\beta|^2$ . Since the total probability must sum to 1, we have that  $|\alpha|^2 + |\beta|^2 = 1$ . Returning to the vector space interpretation, we have that the qubit's state is a unit vector in  $\mathbb{C}^2$ .

**Example 1.** A qubit can have the following state,

$$\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle,$$

when measured the state gives the result 0 fifty percent of the time, as well as the result 1 fifty percent of the time. This state is commonly denoted  $|+\rangle$ .

We can think about qubits using a different geometric interpretation by rewriting Equation (1.1) as

$$|\psi\rangle = e^{i\gamma} \left[ \cos\left(\frac{\theta}{2}\right) |0\rangle + e^{i\phi} \sin\left(\frac{\theta}{2}\right) |1\rangle \right], \quad (1.2)$$

where  $\theta, \phi, \gamma \in \mathbb{R}$ . We will later see that we can ignore the factor of  $e^{i\gamma}$ , because it has no observable effects. We can now write

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right) |0\rangle + e^{i\phi} \sin\left(\frac{\theta}{2}\right) |1\rangle, \quad (1.3)$$

where the numbers  $\theta$  and  $\phi$  define a point on the three-dimensional unit sphere. We will refer to this as the *Bloch sphere*.




Figure 1.1: Bloch Sphere REFERENCE

### 1.1.1 Multiple Qubits

With two classical bits, there are four possible states: 00, 01, 10, and 11. Similarly, qubits have the four corresponding computation basis states:  $|00\rangle$ ,  $|01\rangle$ ,  $|10\rangle$ , and  $|11\rangle$ .

A pair of qubits can also exist in superposition. This can be represented as the following

$$|\phi\rangle = \alpha_{00} |00\rangle + \alpha_{01} |01\rangle + \alpha_{10} |10\rangle + \alpha_{11} |11\rangle.$$

Where each  $\alpha_{ij}$  is a complex coefficient, known as the *amplitude*. The measurement result for each of the four states occurs with probability  $|\alpha_{ij}|^2$