

Notation 1 (*Preliminaries*). Let \mathcal{L} be a countable language and let T be a stable theory in \mathcal{L} which permits infinite models. We will also let \mathcal{A} and \mathcal{B} denote \mathcal{L} -structures with universes A and B respectively. When relevant, let $\mathcal{A} \models T$. We will also be using the monster model $\mathbb{M} \succeq \mathcal{A}$ for many of the following results.

- Let S be the set of all \mathcal{L} -formulas. Throughout this write-up, we will let Δ be a set of \mathcal{L} -formulas, so $\Delta \subseteq S$.
- Let S_n be the set of all \mathcal{L} -formulas with at most one free variable x .
- For any set X , let $\mathcal{L}_X = \mathcal{L} \cup \{c_x : x \in X\}$, where we augment \mathcal{L} with constant symbols for every element of X .
- Let $S(X)$ be the set of all \mathcal{L}_X formulas.
- Let $S_1(X)$ be defined similarly.
- If $X \subseteq A$, we define $\text{Th}(\mathcal{A}, X)$ to be the theory of \mathcal{A} interpreted as an \mathcal{L}_X -structure. In particular, $\text{Th}(\mathcal{A}, A)$ is the elementary diagram of \mathcal{A} .
- An \mathcal{A} -instance of a formula φ is any formula obtained from φ by substituting $c_a \in \mathcal{L}_A$ for each variable.

Definition 1 (*Boolean closure*). Let Δ be a set of \mathcal{L} -formulas and let X be a set. We define $\Delta^b(X)$ to be the set of formulas in $S_1(X)$ obtained from formulas in Δ using conjunction, disjunction, negation, and substituting instances of c_x for a variable. Trivially, $S^b(A) = S(A)$.

Definition 2 (*Partitioning*). Let $\varphi \in S_1(B)$ and $\Gamma = \{\psi_1, \dots, \psi_n\} \subset S_1(B)$. We say that Γ *partitions* φ if we have the following.

$$\begin{aligned} \mathcal{B} &\models \forall x(\psi_1(x) \vee \dots \vee \psi_n(x) \vee \neg\varphi(x)) \\ \mathcal{B} &\models \forall x((\psi_i(x) \wedge \psi_j(x)) \rightarrow \neg\varphi(x)) \quad \text{for all } i \neq j \\ \mathcal{B} &\models \exists x(\varphi(x) \wedge \psi_i(x)) \quad \text{for all } i \end{aligned}$$

In other words, whenever $\varphi(x)$ is true in \mathcal{B} , exactly one of the $\psi_i(x)$'s is true.

Definition 3 (*Generalized rank and degree*). Let $\Delta \subseteq A$. In order to define a generalized version of Morley rank and degree, we will need to define two sets $S^\alpha(\mathcal{A}, \Delta)$ and $\text{Tr}^\alpha(\mathcal{A}, \Delta)$ for any ordinal α inductively.

- Let $S^0(\mathcal{A}, \Delta) = \{\varphi \in S_1(A) : \mathcal{A} \models \exists x\varphi(x)\}$, i.e. the set of all \mathcal{L} -formulas that have witnesses in \mathcal{A} .
- If $S^\alpha(\mathcal{A}, \Delta)$ is already defined, we may define $\text{Tr}^\alpha(\mathcal{A}, \Delta)$ as follows. Given a formula

φ , we say that $\varphi \in \text{Tr}^\alpha(\mathcal{A}, \Delta)$ if and only if there exists a finite k such that for all $\mathcal{B} \succeq \mathcal{A}$ and any finite $\Gamma \subset \Delta^b(\mathcal{B})$ partitioning \mathcal{A} , there are no more than k formulas ψ in Γ such that $(\varphi \wedge \psi) \in S^\alpha(\mathcal{A}, \Delta)$.

- Let $S^{\alpha+1}(\mathcal{A}, \Delta) = S^\alpha(\mathcal{A}, \Delta) \setminus \text{Tr}^\alpha(\mathcal{A}, \Delta)$.
- For α limit, let $S^\alpha(\mathcal{A}, \Delta) = \bigcap_{\beta < \alpha} S^\beta(\mathcal{A}, \Delta)$.
- The Δ -rank of φ in \mathcal{A} is the least (and unique) ordinal α such that $\varphi \in \text{Tr}^\alpha(\mathcal{A}, \Delta)$. When Δ is unambiguous, we denote this as $\text{rank}_{\mathcal{A}}(\varphi)$.
- The Δ -degree of φ in \mathcal{A} is the least number k witnessing that $\varphi \in \text{Tr}^\alpha(\mathcal{A}, \Delta)$. When Δ is unambiguous, we denote this as $\text{deg}_{\mathcal{A}}(\varphi)$.

Proposition 1 (Δ -rank generalizes Morley rank). Let p be a 1-type in $\text{Th}(\mathcal{A}, A)$. The Morley rank of p is the least α such that $p \cap \text{Tr}^\alpha(\mathcal{A}, S) \neq \emptyset$. The Morley degree of p is the minimum of the S -degrees of the formulas in $p \cap \text{Tr}^\alpha(\mathcal{A}, S)$.

Proof. Let $\mathcal{B} \succeq \mathcal{A}$ be an \aleph_0 -saturated model, so p is realized in \mathcal{B} . To prove the first part, we show by induction on α that for all $\varphi \in p$,

$$\text{RM}(\varphi) = \text{RM}^{\mathcal{B}}(\varphi) \geq \alpha \Leftrightarrow \varphi \in S^\alpha(\mathcal{A}, S).$$

If $\alpha = 0$, we clearly have by definition that $\mathcal{A} \models \exists x \varphi(x)$ if and only if $\varphi(\mathcal{A})$ is nonempty. The limit case follows easily by induction.

As for the successor case, we need to show that $\text{RM}(\varphi) \geq \alpha + 1$ if and only if φ is in $S^\alpha(\mathcal{A}, S) \setminus \text{Tr}^\alpha(\mathcal{A}, S)$. First suppose that $\text{RM}(\varphi) \geq \alpha + 1$, so there exist infinitely many $\mathcal{L}_{\mathcal{B}}$ -formulas ψ_1, ψ_2, \dots defining pairwise disjoint subsets of $\varphi(\mathcal{B})$ such that $\text{RM}(\psi_i) \geq \alpha$ for all $i < \omega$. Let k be any finite number. Then, define the following finite set of formulas $\Gamma \subset S^b(\mathcal{B})$.

$$\Gamma = \left\{ \psi_1, \dots, \psi_{k+1}, \left(\varphi \wedge \bigwedge_{i=1}^{k+1} \neg \psi_i \right) \right\}$$

This set clearly partitions φ . We can see that for at least $k + 1$ formulas $\psi_i \in \Gamma$, $\psi_i \wedge \varphi = \psi_i$ has rank α , which by induction means $\psi_i \in S^\alpha(\mathcal{B}, S)$. Thus, we conclude that φ cannot be in $\text{Tr}^\alpha(\mathcal{A}, S)$.

Now suppose that $\text{RM}(\varphi) = \alpha$. Then, we can see that the Morley degree of φ as an $\mathcal{L}_{\mathcal{B}}$ -formula is a k witnessing that $\varphi \in \text{Tr}^\alpha(\mathcal{A}, S)$. In fact, by the definition of the Morley degree of a type, the second part of the proposition is trivial. \square

Definition 4 (*Minimality*). A formula φ is *minimal* in $\text{Tr}^\alpha(\mathcal{A}, \Delta)$ if $\varphi \in \text{Tr}^\alpha(\mathcal{A}, \Delta)$ and there is no $\psi \in \Delta$ with some \mathcal{A} -instance ψ' such that $(\varphi \wedge \psi')$ and $(\varphi \wedge \neg \psi')$ are both in $\text{Tr}^\alpha(\mathcal{A}, \Delta)$.

Lemma 1 (*Properties of rank*). Let $\varphi, \varphi_0, \varphi_1$, and ψ be formulas in $S_1(A)$.

- (i) Suppose $\varphi(\mathcal{A}) \subseteq \psi(\mathcal{A})$. Then $\text{rank}_{\mathcal{A}}(\varphi) \leq \text{rank}_{\mathcal{A}}(\psi)$ and, if $\text{rank}_{\mathcal{A}}(\varphi) = \text{rank}_{\mathcal{A}}(\psi)$, then $\deg_{\mathcal{A}}(\varphi) \leq \deg_{\mathcal{A}}(\psi)$.
- (ii) If $\mathcal{B} \succeq \mathcal{A}$, then $\text{rank}_{\mathcal{A}}(\varphi) = \text{rank}_{\mathcal{B}}(\varphi)$.
- (iii) If $(\varphi_0 \vee \varphi_1)(\mathcal{A}) \supseteq \varphi(\mathcal{A})$ and $\text{rank}_{\mathcal{A}}(\varphi) \geq \alpha$, then at least one of φ_0 and φ_1 has a Δ -rank of at least α . In particular, if $\text{rank}_{\mathcal{A}}(\varphi) = \alpha$ for any ordinal α , then $\deg_{\mathcal{A}}(\varphi) \geq 1$.

Proof. (i) Clearly $\varphi \in S^0(\mathcal{A}, \Delta)$ implies that $\psi(\mathcal{A}) \supseteq \varphi(\mathcal{A}) \neq \emptyset$, so $\psi \in S^0(\mathcal{A}, \Delta)$. For the successor case, given a set which partitions φ with k formulas in $S^\alpha(\mathcal{A}, \Delta)$, we easily also have a set which partitions ψ with k formulas in $S^\alpha(\mathcal{A}, \Delta)$. Thus, if $\varphi \notin \text{Tr}^\alpha(\mathcal{A}, \Delta)$, then $\psi \notin \text{Tr}^\alpha(\mathcal{A}, \Delta)$. The limit case follows easily from induction. Also, if k formulas of rank α partition φ , then these same formulas partition ψ , giving the second part of the statement easily.

(ii) Since $\mathcal{A} \models \exists x \varphi(x)$ if and only if $\mathcal{B} \models \exists x \varphi(x)$, we immediately have the case of $\alpha = 0$. In fact, since the property of a set partitioning a formula depends only on the satisfaction of finitely many sentences, a set partitioning φ in \mathcal{A} with k formulas in $S^\alpha(\mathcal{A}, \Delta)$ will also be a set partitioning φ in \mathcal{B} with k formulas in $S^\alpha(\mathcal{B}, \Delta)$ and vice versa by induction. The limit case also follows easily.

(iii) We have the following deduction to show that this holds for the case of $\alpha = 0$.

$$\mathcal{A} \models \exists x \varphi(x) \Rightarrow \mathcal{A} \models \exists x (\varphi_0(x) \vee \varphi_1(x)) \Rightarrow \mathcal{A} \models \exists x \varphi_0(x) \vee \exists x \varphi_1(x)$$

Suppose the lemma holds for α , suppose $\varphi \in S^{\alpha+1}(\mathcal{A}, \Delta)$, and let $k \in \omega$. There exists a set Γ which partitions φ with formulas $\psi_1, \dots, \psi_{2k} \in \Gamma$ such that each ψ_i is in $S^\alpha(\mathcal{A}, \Delta)$. For each ψ_i , at least one of $(\psi_i \wedge \varphi_0)$ and $(\psi_i \wedge \varphi_1)$ is in $S^\alpha(\mathcal{A}, \Delta)$ by induction. Since we have $2k$ formulas, for at least one of the φ_j 's there are at least k formulas of the form $(\psi_i \wedge \varphi_j)$ in $S^\alpha(\mathcal{A}, \Delta)$. Then we can easily construct a partitioning set showing that φ_j cannot have degree k . We are able to show that for any $k \in \omega$ one of φ_0 and φ_1 is unable to have degree k . Thus, for one of the φ_j 's, φ_j fails to have degree k for infinitely many k , meaning $\varphi_j \in S^{\alpha+1}(\mathcal{A}, \Delta)$.

Finally, for the limit case, assuming the lemma holds for all $\beta < \alpha$, it is true for at least one of the φ_j 's that there is a cofinal sequence of ordinals in α where for each entry β , $\varphi_j \in S^\beta(\mathcal{A}, \Delta)$. Thus, $\varphi_j \in S^\alpha(\mathcal{A}, \Delta)$. \square

Definition 5 (*Weak satisfiability*). First, for any set of formulas Θ , we define

$$\Theta^- = \{\neg\phi : \phi \in \Theta\}$$

to be the set of negations of all formulas in Θ . The set $\Gamma \subset S(A)$ is *weakly satisfiable* in \mathcal{A} if no finite disjunctions of formulas in Γ^- is valid in \mathcal{A} .

Remark 1. A set $\Gamma \subset S(A)$ is weakly satisfiable in \mathcal{A} if and only if Γ is satisfiable in some $\mathcal{B} \succeq \mathcal{A}$.

Proof. For the rightward direction, $\Gamma \cup \text{Diag}_{\text{el}}(\mathcal{A})$ is finitely satisfiable and therefore satisfiable. For the leftward direction, if $\Gamma \cup \text{Diag}_{\text{el}}(\mathcal{A})$ were not satisfiable, then there would exist formulas $\psi_1, \dots, \psi_n \in \Gamma^-$ such that $\text{Diag}_{\text{el}}(\mathcal{A}) \models \forall \bar{x}(\psi_1 \vee \dots \vee \psi_n)$, so Γ would not be weakly satisfiable in \mathcal{A} . \square

Lemma 2 (*A characterization of rank*). Let \mathcal{A} and \mathcal{B} be \mathcal{L} -structures. Let $\varphi \in S^\alpha(\mathcal{A}, \Delta)$. Let Γ be the set of all formulas obtained by switching around variables in the formulas of $\{\varphi\} \cup \Delta \cup \Delta^-$. There exists $\Gamma^* \subseteq \Gamma$ weakly satisfiable in \mathcal{A} such that if $\psi \in S_1(\mathcal{B})$ and Γ^+ is weakly satisfiable in \mathcal{B} , where Γ^+ is obtained from Γ^* by replacing each instance of φ by the corresponding instance of ψ , then $\psi \in S^\alpha(\mathcal{B}, \Delta)$.

Proof. We proceed by induction on α . For $\alpha = 0$, the set $\Gamma' = \{\varphi\}$ works, as well as its obvious counterpart $\Gamma^+ = \{\psi\}$.

If α is a limit ordinal, suppose the lemma holds for all $\beta < \alpha$. Then, for each β , there exists some $\Gamma_\beta^* \subseteq \Gamma$ weakly satisfiable in \mathcal{A} such that the corresponding set Γ_β^+ has the desired property of the lemma. We may change the variables in each Γ_β such that if $\beta, \gamma < \alpha$ and $\beta \neq \gamma$, then Γ_β and Γ_γ have no variables in common. Then, let $\Gamma^* = \bigcup_{\beta < \alpha} \Gamma_\beta^*$, so $\Gamma^+ = \bigcup_{\beta < \alpha} \Gamma_\beta^+$. Clearly, Γ^* is weakly satisfiable in \mathcal{A} . If Γ^+ is weakly satisfiable in \mathcal{B} , then each Γ_β^+ is weakly satisfiable. By induction, that means $\psi \in S^\beta(\mathcal{B}, \Delta)$ for all $\beta < \alpha$ and so $\psi \in S^\alpha(\mathcal{B}, \Delta)$.

Now suppose that α is a successor with $\alpha = \beta + 1$. We will want to construct a nice sequence of formulas in order to construct Γ^* .

Claim. There exists $\mathcal{A}' \succeq \mathcal{A}$ and a sequence $\langle \varphi_n : n \in \omega \rangle$ of formulas in $\Delta^b(\mathcal{A}')$ such that, for all $m, n \in \omega$,

- $(\varphi \wedge \varphi_n) \in S^\beta(\mathcal{A}', \Delta)$,
- $\neg(\varphi_m \wedge \varphi_n)$ is a tautology, i.e. valid in every structure, if $m \neq n$, and
- φ_n is a conjunction of \mathcal{A}' -instances of formulas in $\Delta \cup \Delta^-$.

First, since $\varphi \in S^\alpha(\mathcal{A}, \Delta)$, there exists $\mathcal{A}_0 \succeq \mathcal{A}$ and mutually exclusive $\theta_0, \theta_1 \in \Delta^b(\mathcal{A}_0)$ such that $(\varphi \wedge \theta_0), (\varphi \wedge \theta_1) \in S^\beta(\mathcal{A}_0, \Delta)$. By **Lemma 1**(iii), we can let θ be one of θ_0, θ_1 , and $(\neg\theta_0 \wedge \neg\theta_1)$ such that $(\varphi \wedge \theta) \in S^\alpha(\mathcal{A}_0, \Delta)$. If $\theta = \theta_0$, let $\varphi_0^* = \theta_1$. Otherwise, let $\varphi_0^* = \theta_0$.

By repeated use of **Lemma 1**(iii) and laws of Boolean algebra, we can find φ_0 , a conjunction of \mathcal{A}_0 -instances of formulas in $\Delta \cup \Delta^-$ such that $\varphi_0(\mathcal{A}_0) \subseteq \varphi_0^*(\mathcal{A}_0)$ and $\varphi \in S^\beta(\mathcal{A}_0, \Delta)$.

Since $\theta \in S^\alpha(\mathcal{A}, \Delta)$, we can repeat this process on θ instead of φ and, indeed, continue it indefinitely. This gives us a sequence of formulas $\langle \varphi_n : n \in \omega \rangle$ and an increasing chain of models

$$\mathcal{A} \preceq \mathcal{A}_0 \preceq \mathcal{A}_1 \preceq \dots$$

where, for all n , φ_n is a conjunction of \mathcal{A}_n -instances of formulas in $\Delta \cup \Delta^-$. Finally, let $\mathcal{A}' = \bigcup_{n \in \omega} \mathcal{A}_n$. Then, for all n , $(\varphi \wedge \varphi_n) \in S^\beta(\mathcal{A}', \Delta)$ by **Lemma 1**(ii). The sequence also satisfies the other two desired properties, meaning we have proven the claim.

There is an issue with this proof: I have shown that these formulas are mutually exclusive in \mathcal{A}' , but this is a weaker notion than the second part of the claim. To salvage this: make it so φ_{n+1} has the negation of a formula $\delta_m \in \Delta \cup \Delta^-$ for all $m \leq n$, where δ_m appears in φ_m but not φ_{m-1} .

For each n , let Γ_n be the set of all formulas obtained by switching around variables in the formulas of $\{(\varphi \wedge \varphi_n)\} \cup \Delta \cup \Delta^-$. Since we have assumed the lemma for β , there exists $\Gamma'_n \subseteq \Gamma_n$ weakly satisfiable in \mathcal{A}' with our desired property. We can also assume that Γ_m and Γ_n share no variables if $m \neq n$ and that there are infinitely many variables not occurring in any Γ_n .

For each $c_a \in \mathcal{L}_A$ which occurs in one of our formulas φ_n we can set aside a variable x_a ; this variable is independent of n . Let Γ_n^* be obtained from Γ_n by replacing each occurrence of c_a with a corresponding variable x_a . Let $\Gamma^* = \bigcup_{n \in \omega} \Gamma_n^*$. We can see that Γ^* is weakly satisfiable in \mathcal{A} . Now, let Γ^\dagger be the set obtained from Γ^* by replacing each instance of φ with ψ .

Suppose Γ^\dagger is weakly satisfiable in \mathcal{B} . Then, by **Remark 1**, there exists $\mathcal{B}' \succeq \mathcal{B}$ where Γ^\dagger is satisfied. By replacing each variable v_a with a constant symbol in $\mathcal{L}_{\mathcal{B}'}$ appropriately, we have a sequence of formulas $\langle \psi_n : n \in \omega \rangle$ corresponding to each φ_n . This gives us infinitely many disjoint formulas $(\psi \wedge \psi_n) \in S^\beta(\mathcal{B}', \Delta)$, witnessing that $\psi \in S^\alpha(\mathcal{B}', \Delta)$. By **Lemma 1**(ii), $\psi \in S^\alpha(\mathcal{B}, \Delta)$. \square

Lemma 3. Let Δ be finite, $n \in \omega$, and let ψ have Δ -rank n and Δ -degree 1. For each $\varphi \in (\Delta \cup \Delta^-) \cap S_{k+1}$ and for any $\mathcal{B} \succeq \mathcal{A}$, the set

$$D = \{\bar{b} \in \mathcal{B}^k : \text{rank}_{\mathcal{B}}(\psi(x) \wedge \varphi(x, \bar{b})) = n\}$$

is definable over \mathcal{A} .

Proof. We may first suppose that \mathcal{B} is the monster model \mathbb{M} . We assume that each $\varphi \in \Delta$ is stable since the theory T is stable. Letting φ be any formula in Δ or Δ^- with $k+1$ free variables, we define the following global φ -type.

$$p = \{\varphi(x, \bar{b}) : \bar{b} \in D\} \cup \{\neg\varphi(x, \bar{c}) : \bar{c} \in \mathbb{M}^k \setminus D\}$$

First, we show that this type is consistent, i.e. that $\text{Diag}_{\text{el}}(\mathcal{A}) \cup p$ is satisfiable. Let $p_0 \subset p$ be finite; we'll write this partial φ -type as

$$\{\varphi(x, \bar{b}_1), \dots, \varphi(x, \bar{b}_r), \neg\varphi(x, \bar{c}_1), \dots, \neg\varphi(x, \bar{c}_s)\}$$

where $\bar{b}_1, \dots, \bar{b}_r \in D$ and $\bar{c}_1, \dots, \bar{c}_s \in \mathbb{M}^k \setminus D$. We will show that

$$\theta_i(x) = \psi(x) \wedge \varphi(x, \bar{b}_1) \wedge \dots \wedge \varphi(x, \bar{b}_i)$$

has Δ -rank n for every $i \in \{1, \dots, r\}$. It will follow, since $n \geq 0$, that this formula has a witness. First, we have that $\text{rank}(\psi(x) \wedge \varphi(x, \bar{b}_1)) = n$ by definition. Now, let $i \geq 2$ and suppose $\text{rank}(\theta_{i-1}(x)) = n$. We know that $\deg(\theta_{i-1}(x)) = 1$ by [Lemma 1](#)(i) and (iii). Thus, if $\text{rank}(\theta_{i-1}(x) \wedge \varphi(x, \bar{b}_i)) < n$, then it must be the case that $\text{rank}(\theta_{i-1}(x) \wedge \neg\varphi(x, \bar{b}_i)) = n$. However, by [Lemma 1](#)(i), this means that $\text{rank}(\psi(x) \wedge \neg\varphi(x, \bar{b}_i)) = n$, contradicting that ψ has Δ -degree 1. So, it can only be the case that $\text{rank}(\theta_i(x)) = n$. This along with analogous logic for each $\neg\varphi(x, \bar{c}_i)$ shows that

$$\text{rank}_{\mathbb{M}} \left(\psi(x) \wedge \bigwedge \varphi(x, \bar{b}_i) \wedge \bigwedge \neg\varphi(x, \bar{c}_i) \right) = n$$

In particular, since $n \geq 0$, the formula has a witness in \mathbb{M} , meaning that p_0 is satisfiable and so p is realized in \mathbb{M} .

Since φ is stable, the φ -type p is definable by some formula $\theta(\bar{y})$ with parameters in the monster. This formula also defines D . To finish our proof, we only need to show that D is A -invariant. Let σ be any automorphism on \mathbb{M} which fixes A pointwise. We must show that for any $\bar{b} \in D$, $\sigma(\bar{b}) \in D$. Since the only names in ψ are names of elements of A and the only names in $\varphi(x, \bar{b})$ are \bar{b} , we obtain the following.

$$\begin{aligned} \text{rank}(\sigma(\psi(x) \wedge \varphi(x, \bar{b}))) &= \text{rank}(\sigma(\psi(x)) \wedge \sigma(\varphi(x, \bar{b}))) \\ &= \text{rank}(\psi(x) \wedge \varphi(x, \sigma(\bar{b}))) = n \end{aligned}$$

Thus, $\sigma(\bar{b}) \in D$, meaning D is A -invariant. Thus, D is A -definable. Since we are working in the monster model, whatever formula defines D in \mathbb{M} will also define any \mathcal{B} extending \mathcal{A} , completing our proof. \square

Lemma 4. If Δ is finite then $\text{Tr}^\alpha(\mathcal{A}, \Delta) = \emptyset$ if $\alpha \geq \omega$.

Proof. No proof but explains some analogous results. □

Definition 6 (Δ -stable). To be added. Generalization of stability (duh).

Lemma 5. Let $\mathcal{A} \preceq \mathcal{B}$, let φ be a formula in $S_1(A)$ which is Δ -stable, and let $\psi \in \Delta^b(B)$. Then there exists $\theta \in S_1(A)$ such that $\varphi(\mathcal{A}) \cap \psi(\mathcal{B}) = \theta(\mathcal{A})$.

Proof. No proof but some references listed, including Shelah. □

Lemma 6. If φ is minimal in $\text{Tr}^\alpha(\mathcal{A}, S)$, then the S -degree of φ is 1.

Proof. Long, seemingly full proof given. □

That last lemma says that all transcendental points in the Stone space of a model have degree 1.

Lemma 7. If Δ is finite and φ is minimal in $\text{Tr}^n(\mathcal{A}, \Delta)$, then the Δ -degree of φ in \mathcal{A} is 1.

Proof. Lachlan's experimental notation really shines here. □

Theorem 1 (*Model extension*). Let \mathcal{A} and \mathcal{B} be models of a countable stable theory and suppose that $\mathcal{A} \prec \mathcal{B}$ and $P(\mathcal{A}) = P(\mathcal{B})$ where P is a unary predicate symbol. There exists $\mathcal{C} \succ \mathcal{B}$ such that $P(\mathcal{C}) = P(\mathcal{B})$.

Proof. Long! Uses a theorem of Ehrenfeucht. Check Reference 8. □