
Primary and core sensitivity analyses
Primary
Exclude Cevik
Include Iachetta and Kleibl
Include Boisdron−Celle overlap
Exclude Onesti
Exclude genotype−informed carriers
Exclude PK−informed Morel
Exclude all treatment−moderated
Wigle early window
Cremolini first four cycles
Compound lower−function allocation
Compound reciprocal allocation
1.0
1.5
2.0
2.5
3.0
3.5
Odds ratio


WT comparator−classification sensitivities
Loose: HapB3 vs WT_Clean
Loose: HapB3 vs WT_Biased
Strict: HapB3 vs WT_Clean
Strict: HapB3 vs WT_Biased
0.5
1.0
2.0
5.0
Odds ratio


Tag/causal−SNP and sparse−data sensitivities
Exclude carrier arms with N <= 2
Bozina 2021: best case
Bozina 2021: worst case
Wigle 2021 prospective: best case
Wigle 2021 prospective: worst case
Rosmarin 2015 arm B: best case
Rosmarin 2015 arm B: worst case
1.0
1.5
2.0
2.5
3.0
Odds ratio


Grouped leave−one−source−out sensitivities
Exclude Boige 2016 FFCD 2000 05
Exclude Boige 2016 PETACC 8
Exclude Botticelli 2017
Exclude Bozina 2021
Exclude Braun 2009
Exclude Cevik 2022
Exclude Cremolini 2018
Exclude Etienne Grimaldi 2017
Exclude Falvella 2015
Exclude Froehlich 2015
Exclude Ikonnikova 2024
Exclude Jennings 2013
Exclude Largillier 2006
Exclude Lee N0147
Exclude Loganayagam 2013
Exclude Morel 2006
Exclude Onesti 2017
Exclude Ragia 2023
Exclude Rosmarin 2015
Exclude Salgueiro 2004
Exclude Schwab 2008
Exclude Toffoli 2015
Exclude Wigle 2021 prospective
1.0
1.5
2.0
2.5
3.0
3.5
Odds ratio