

# CU Denver Math Camp - Limits & Derivatives

## Day 2

Michael R. Karas

August 6, 2024

University of Colorado, Boulder

# Day 2 Topics

- Practice Problems
- Increasing/Decreasing Functions
- Concave/Convex Functions
- Implicit Differentiation
- Partial Derivatives
- Taylor Series

# Day 1 Practice Problems

$$f(x) = \frac{2x}{x^2 + 2}$$

# Day 1 Practice Problems

$$f(x) = e^{x^2 + \ln(x)}$$

## Day 1 Practice Problems

$$f(x) = x(x^2 + 1)$$

Find  $f'(x)$  and  $f''(x)$

# Day 1 Practice Problems

$$f(x) = \ln(x^2 + 1)$$

Find  $f'(x)$  and  $f''(x)$

# Day 1 Practice Problems

Prove that  $\frac{d}{dx}e^x = e^x$

# Increasing/Decreasing Functions

Let  $f$  be a differentiable function defined on an interval  $[a, b]$

- $f$  is increasing on  $[a, b]$  if, for every  $a \leq x \leq b$ ,  $f'(x) > 0$
- $f$  is decreasing on  $[a, b]$  if, for every  $a \leq x \leq b$ ,  $f'(x) \leq 0$

Could replace  $\leq$  with  $<$  and say “strictly increasing” (no flat parts of  $f$  on  $[a, b]$ ), likewise with decreasing. When  $f'(x) = 0$  the function is not changing, an “optimal” point.



# Increasing/Decreasing Functions Practice Problems

$$f(x) = x^2$$

Find which intervals which  $f(x)$  is increasing and which intervals which  $f(x)$  is decreasing.

# Increasing/Decreasing Functions Practice Problems

$$f(x) = \ln(x)$$

Find which intervals which  $f(x)$  is increasing and which intervals which  $f(x)$  is decreasing.

# Increasing/Decreasing Functions Practice Problems

$$f(x) = e^x$$

Find which intervals which  $f(x)$  is increasing and which intervals which  $f(x)$  is decreasing.

# Concave/Convex Functions



Often it is not enough to know if a function is increasing/decreasing. We need to know its shape. Let  $f$  and  $I$  be as before. Then,

- $f$  is convex on  $[a, b]$  if, for every  $a \leq x \leq b$ ,  $f''(x) > 0$
- $f$  is concave on  $[a, b]$  if, for every  $a \leq x \leq b$ ,  $f''(x) < 0$

Convex: function is increasing at an increasing rate to the bottom of a “valley.” Concave: function is increasing at a decreasing rate to the peak of a “mountain.”

- If  $f''(x) = 0$ ,  $f$  is at an inflection point – the acceleration switches (e.g. skiing)

# Concave/Convex Functions



# Concave/Convex Functions Practice Problems

$$f(x) = x - 2\ln(x + 1)$$

# Implicit Differentiation

Sometimes we can't solve for  $y = f(x)$  (or it would be very difficult) but still need to differentiate. Implicit differentiation allows us to find  $f'(x)$  without a “closed-form” solution for  $y = f(x)$ .

- Differentiate both sides of the equation using all the typical rules of derivatives, treating  $x$  and  $y$  as variables that are both changing
- If you differentiate something involving  $y$ , multiply the answer by  $y'$  (the derivative of  $y = f(x)$ )
- If you differentiate something involving  $x$ , do not multiply by anything new
- After everything is differentiated, solve for  $y'$  using algebra
- The answer for  $y'$  should be a function of  $x$  (and potentially  $y$ )

# Implicit Differentiation

Suppose  $x + y + xy = 5$ .

$$1 + 1 \cdot y' + \underbrace{1 \cdot y}_{f'(x)g(x)} + \underbrace{x \cdot 1 \cdot y'}_{f(x)g'(x)} = 0$$

Using product rule:

$$1 + y' + y + xy' = 0$$

$$y'(1 + x) = -1 - y$$

$$y' = \frac{-1 - y}{1 + x}$$

The derivative is a function of both  $x$  and  $y$ .



# Implicit Differentiation Practice Problems

$$x^2 + y^2 = 1$$

# Partial Derivatives

The function  $z = f(x, y)$  takes two inputs and outputs a third. We can evaluate what happens to that output when we change  $x$  or  $y$  (but not both).

- $\frac{\partial z}{\partial x} =$  “the derivative of  $z$  with respect to  $x$ ”
  - The slope of  $f$  in the cardinal direction of  $x$  (e.g. “north-south”)
  - The tangent of  $f$  in the  $x$  direction at a point  $(x_0, y_0)$
- $\frac{\partial z}{\partial y} =$  “the derivative of  $z$  with respect to  $y$ ”
  - The slope of  $f$  in the cardinal direction of  $y$  (e.g. “east-west”)
  - The tangent of  $f$  in the  $y$  direction at a point  $(x_0, y_0)$

# Partial Derivatives

All of these mean “take the partial derivative of  $f$  with respect to  $x$ ”:

- $f_x(x, y)$
- $\frac{\partial f(x, y)}{\partial x}$
- $\frac{\partial}{\partial x} f(x, y)$
- $\frac{\partial z}{\partial x}$  if  $z = f(x, y)$

# Partial Derivatives

Interpret  $\frac{\partial z}{\partial x}$  as “what happens to  $z$  if I change  $x$ , holding  $y$  constant.” Likewise for  $\frac{\partial z}{\partial y}$ .

Suppose  $z = x + y + xy$ , find  $\frac{\partial z}{\partial x}$ .

$$\frac{\partial z}{\partial x} = \frac{\partial}{\partial x}[x + y + xy] = \frac{\partial}{\partial x}x + \frac{\partial}{\partial x}y + \frac{\partial}{\partial x}xy = 1 + 0 + y \cdot 1 = 1 + y$$

# Partial Derivatives

The first partial derivatives  $f_x(x, y)$  and  $f_y(x, y)$  are themselves functions of  $x$  and  $y$ . We could differentiate each of them with respect to either  $x$  or  $y$ , so there are four second-order partial derivatives.

$$\begin{array}{cccc} f(x, y) & & & \\ \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} & & \\ \frac{\partial^2 f}{\partial x^2} & \frac{\partial^2 f}{\partial y \partial x} & \frac{\partial^2 f}{\partial x \partial y} & \frac{\partial^2 f}{\partial y^2} \end{array}$$

# Partial Derivatives

All of these mean “take the partial derivative of  $f$  with respect to  $x$  twice”:

- $f_{xx}(x, y)$
- $\frac{\partial^2 f(x, y)}{\partial x^2}$
- $\frac{\partial^2}{\partial x^2} f(x, y)$
- $\frac{\partial^2 z}{\partial x^2}$  if  $z = f(x, y)$

# Partial Derivatives

All of these mean “take the partial derivative of  $f$  with respect to  $x$ , then with respect to  $y$ ”:

- $f_{xy}(x, y)$
- $\frac{\partial^2 f(x, y)}{\partial y \partial x}$
- $\frac{\partial^2}{\partial y \partial x} f(x, y)$
- $\frac{\partial^2 z}{\partial y \partial x}$  if  $z = f(x, y)$

# Young's Theorem

If  $f(x, y)$  is a twice differentiable function and continuous at the point  $(x_0, y_0)$ , then

$$f_{xy}(x_0, y_0) = f_{yx}(x_0, y_0)$$

The cross-partial derivatives are the same, regardless of the order in which you take them.



# Partial Derivatives Practice Problems

$$f(x, y) = x^2 + y^2 + x^2y^2$$

# Partial Derivatives Practice Problems

$$f(x, y) = x^2y + \ln(x)y^3$$

# Taylor Series

Sometimes it is too difficult (or not possible) to differentiate a function. We can use a Taylor Series expansion to approximate the value of a function around different points.

For a function  $f(x)$  with derivatives of all orders at  $x = a$ , the Taylor series is:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

When  $a = 0$ , it is called the Maclaurin series.

# Taylor Series

Taylor Series for  $f(x) = \frac{1}{x}$

$$f(x) = \frac{1}{x} \quad f'(x) = -\frac{1}{x^2} \quad f''(x) = \frac{2}{x^3} \quad f^{(k)}(x) = \frac{k!}{x^{k+1}}$$

$$P_k(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(k)}(a)}{k!}(x-a)^k$$


At  $a = 1$  of order  $k = 2$ :

$$P_2(x) = 1 - (x-1) + (x-1)^2$$

At  $a = 3$  of order  $k = 2$ :

$$P_3(x) = \frac{1}{3} - \frac{x-3}{3^2} + \frac{(x-3)^2}{3^3}$$

# Taylor Series



# Taylor Series Practice Problems

Write the Taylor Series Expansion of order  $k$  generated for the function  $f(x) = e^x$  around the point  $x = 0$