

# Introduction to parton showers, matching and merging

Marek Schönherr

Institute for Particle Physics Phenomenology, Durham University

CERN, 11 Jun 2024



THE  
ROYAL  
SOCIETY

## Before we begin

Parton showers are an **active field of research**, though we have the experience of over four decades of development.

Many issues are currently actively debated and developed. In many cases, there is no final answer yet.

**I am an author of the SHERPA** Monte-Carlo event generator. Although I endeavour to be agnostic, this will invariably influence my point of view and choice of examples to some extent.

Many thanks to S. Höche for letting me steal many plots/sketches/illustrations from his lectures in the MCnet School '21.

## What to expect

- A basic understanding of what a parton shower is, its features and its limitations.
- The underlying concepts of matching and merging, used in most theory predictions for collider experiments today.
- The background that allows you to follow the discussions in the past, present, and (hopefully) future parton shower literature.

## What not to expect

- All the latest and greatest plots, as well as a survey of all possible algorithms. This could fill the entire time of the school.

## Literature

- ① R. K. Ellis, W. J. Stirling, B. R. Webber  
**QCD and Collider Physics**  
Cambridge University Press, 2003
- ② R. D. Field  
**Applications of Perturbative QCD**  
Addison-Wesley, 1995
- ③ M. E. Peskin, D. V. Schroeder  
**An Introduction to Quantum Field Theory**  
Westview Press, 1995
- ④ T. Sjöstrand, S. Mrenna, P. Z. Skands  
**PYTHIA 6.4 Physics and Manual**  
JHEP 05 (2006) 026
- ⑤ S. Höche,  
**Introduction to parton-shower event generators**  
TASI lectures, 2014

# Overview of lectures

- 1) Introduction to parton showers
  - approximate higher-order corrections
  - building a parton shower
- 2) Improving parton showers
  - assessing the properties of a parton shower
  - NLL accuracy and beyond
- 3) Matching and merging
  - matching
  - merging

# Introduction to parton showers

- ① Approximate higher-order corrections
- ② The parton branching process
- ③ Monte-Carlo methods
- ④ Effects
- ⑤ Summary

## Approximate higher-order corrections

## Leading order cross section

- hadron collider cross section for production of system  $Y$   
(think  $Y = \ell^+\ell^-$ ,  $t\bar{t}$ ,  $W^+W^-$ , dijets, ...)

$$d\sigma_{pp \rightarrow Y+X} = \sum_{a,b \in \{q,g\}} dx_a dx_b f_a(x_a, \mu_F^2) f_b(x_b, \mu_F^2) d\Phi_n \frac{d\hat{\sigma}_{ab \rightarrow Y+X}(\Phi, \mu_F^2)}{d\Phi_n}$$

- PDFs  $f_i(x_i, \mu_F^2)$ ,  $n$ -particle phase space element  $d\Phi_n$
- partonic cross section at LO

$$d\hat{\sigma}_{ab \rightarrow Y+X} \propto |\mathcal{M}_{ab \rightarrow Y}^{\text{tree}}|^2$$

**Note:** every cross-section is inclusive in some additional particles.  
The leading order cross section does not contain them explicitly.  
Higher-order corrections must allow additional radiation.



## Leading order cross section

- hadron collider cross section for production of system  $Y$  (think  $Y = \ell^+\ell^-$ ,  $t\bar{t}$ ,  $W^+W^-$ , dijets, ...)

$$d\sigma_{pp \rightarrow Y+X} = \sum_{a,b \in \{q,g\}} dx_a dx_b f_a(x_a, \mu_F^2) f_b(x_b, \mu_F^2) d\Phi_n \frac{d\hat{\sigma}_{ab \rightarrow Y+X}(\Phi, \mu_F^2)}{d\Phi_n}$$

- PDFs  $f_i(x_i, \mu_F^2)$ ,  $n$ -particle phase space element  $d\Phi_n$
- partonic cross section at LO

$$d\hat{\sigma}_{ab \rightarrow Y+X} \propto |\mathcal{M}_{ab \rightarrow Y}^{\text{tree}}|^2$$

**Note:** every cross-section is inclusive in some additional particles. The leading order cross section does not contain them explicitly.

Higher-order corrections must allow additional radiation.

## Leading order cross section

- hadron collider cross section for production of system  $Y$  (think  $Y = \ell^+\ell^-$ ,  $t\bar{t}$ ,  $W^+W^-$ , dijets, ...)

$$d\sigma_{pp \rightarrow Y+X} = \sum_{a,b \in \{q,g\}} dx_a dx_b f_a(x_a, \mu_F^2) f_b(x_b, \mu_F^2) d\Phi_n \frac{d\hat{\sigma}_{ab \rightarrow Y+X}(\Phi, \mu_F^2)}{d\Phi_n}$$

- PDFs  $f_i(x_i, \mu_F^2)$ ,  $n$ -particle phase space element  $d\Phi_n$
- partonic cross section at LO

$$d\hat{\sigma}_{ab \rightarrow Y+X} \propto |\mathcal{M}_{ab \rightarrow Y}^{\text{tree}}|^2$$

**Note:** every cross-section is inclusive in some additional particles. The leading order cross section does not contain them explicitly. Higher-order corrections must allow additional radiation.

# Approximate NLO corrections

- partonic cross section at NLO

$$d\hat{\sigma}_{ab \rightarrow n+X} \propto \underbrace{|\mathcal{M}_{ab \rightarrow n}^{\text{tree}}|^2}_{\text{Born}} + \underbrace{2\text{Re} \left\{ \mathcal{M}_{ab \rightarrow n}^{\text{loop}} \mathcal{M}_{ab \rightarrow n}^{\text{tree}*} \right\}}_{\text{virtual corr.}} + \underbrace{|\mathcal{M}_{ab \rightarrow n+1}^{\text{tree}}|^2}_{\text{real corr.}}$$

real and virtual correction separately diverging  
(infrared singularities caused by soft or collinear parton emission)  
sum is finite due to Kinoshita-Lee-Nauenberg (KLN) theorem

- infrared limit is universal, depends only on external states, construct

$$d\hat{\sigma}_{n+1}^{\text{approx}} = d\hat{\sigma}_n \otimes \sum_{i,k} dV_{ik}$$

some splitting function  $V_{ik}$ ,  $ik \rightarrow ijk$

# Approximate NLO corrections

- partonic cross section at NLO

$$d\hat{\sigma}_{ab \rightarrow n+X} \propto \underbrace{|\mathcal{M}_{ab \rightarrow n}^{\text{tree}}|^2}_{\text{Born}} + \underbrace{2\text{Re} \left\{ \mathcal{M}_{ab \rightarrow n}^{\text{loop}} \mathcal{M}_{ab \rightarrow n}^{\text{tree}*} \right\}}_{\text{virtual corr.}} + \underbrace{|\mathcal{M}_{ab \rightarrow n+1}^{\text{tree}}|^2}_{\text{real corr.}}$$

real and virtual correction separately diverging  
(infrared singularities caused by soft or collinear parton emission)  
sum is finite due to Kinoshita-Lee-Nauenberg (KLN) theorem

- infrared limit is universal, depends only on external states, construct

$$d\hat{\sigma}_{n+1}^{\text{approx}} = d\hat{\sigma}_n \otimes \sum_{i,k} dV_{ik}$$

some **splitting function**  $V_{ik}$ ,  $ik \rightarrow ijk$

# Approximate NLO corrections

## Collinear approximation

- collinear splitting function  $F_{ab}(z, \phi)$ ,  $a \rightarrow bj$

$$dV_{ak} \rightarrow \frac{dt}{t} dz \frac{d\phi}{2\pi} \frac{\alpha_s}{2\pi} F_{ab}(z, \phi) \xrightarrow{\phi \text{ av.}} \frac{dt}{t} dz \frac{\alpha_s}{2\pi} P_{ab}(z)$$

- azimuthal average:  $F_{ab}(z, \phi) \rightarrow P_{ab}(z)$

Altarelli-Parisi splitting functions

- azimuthally averaged collinear limit of  $n + 1$  matrix element
- dropped spin-correlations in splitting,  
 $\rightarrow dV_{ak}$  is purely multiplicative factor

# Approximate NLO corrections

## Soft approximation

- limit of soft gluon emission

$$dV_{ik} \rightarrow \omega d\omega \frac{d\Omega}{2\pi} \frac{\alpha_s}{2\pi} C_{ik} \frac{p_i \cdot p_k}{p_i \cdot q \, p_k \cdot q}$$

- kinematics described by **Eikonal**
- colour factor in general matrix valued, but

$$C_{ik} = -\mathbf{T}_i \mathbf{T}_k \xrightarrow{\text{large-}N_c} \left\{ \begin{array}{ll} \mathbf{T}_i^2 + \mathcal{O}(1/N_c^2) & \text{for } i = q \\ \frac{1}{2} \mathbf{T}_i^2 + \mathcal{O}(1/N_c^2) & \text{for } i = g \end{array} \right\} \equiv C_i$$

- large- $N_c$  colour factor not matrix-valued any longer, and only depends on parton  $i$

# Approximate NLO corrections

## Soft approximation

- limit of soft gluon emission

$$dV_{ik} \rightarrow \omega d\omega \frac{d\Omega}{2\pi} \frac{\alpha_s}{2\pi} C_{ik} \frac{p_i \cdot p_k}{p_i \cdot q \, p_k \cdot q}$$

- kinematics described by **Eikonal**
- colour factor in general matrix valued, but

$$C_{ik} = -\mathbf{T}_i \mathbf{T}_k \xrightarrow{\text{large-}N_c} \left\{ \begin{array}{ll} \mathbf{T}_i^2 + \mathcal{O}(1/N_c^2) & \text{for } i = q \\ \frac{1}{2} \mathbf{T}_i^2 + \mathcal{O}(1/N_c^2) & \text{for } i = g \end{array} \right\} \equiv C_i$$

- large- $N_c$  colour factor not matrix-valued any longer, and only depends on parton i

# Approximate NLO corrections

- partial-fractioning the Eikonal

$$\frac{p_i \cdot p_k}{p_i \cdot q \, p_k \cdot q} \rightarrow \frac{1}{p_i \cdot q} \frac{p_i \cdot p_k}{(p_i + p_k) \cdot q} + \frac{1}{p_k \cdot q} \frac{p_i \cdot p_k}{(p_i + p_k) \cdot q}$$

The first term contains the soft singularity associated with the region collinear to  $p_i$ , while the second that collinear to  $p_k$ .

- with this, we get

$$dV_{ik} \rightarrow dV_i = \omega d\omega \frac{d\Omega}{2\pi} \frac{\alpha_s}{2\pi} C_i W_{ijk}^i$$

a real-number-valued multiplicative factor of the soft gluon-emission correction in the large- $N_c$  limit

- combine with coll. limit to soft-collinear (dipole) splitting functions



# Approximate NLO corrections

- partial-fractioning the Eikonal

$$\frac{p_i \cdot p_k}{p_i \cdot q \, p_k \cdot q} \rightarrow \frac{1}{p_i \cdot q} \frac{p_i \cdot p_k}{(p_i + p_k) \cdot q} + \frac{1}{p_k \cdot q} \frac{p_i \cdot p_k}{(p_i + p_k) \cdot q}$$

The first term contains the soft singularity associated with the region collinear to  $p_i$ , while the second that collinear to  $p_k$ .

- with this, we get

$$dV_{ik} \rightarrow dV_i = \omega d\omega \frac{d\Omega}{2\pi} \frac{\alpha_s}{2\pi} C_i W_{ijk}^i$$

a real-number-valued multiplicative factor of the soft gluon-emission correction in the large- $N_c$  limit

- combine with coll. limit to soft-collinear (dipole) splitting functions

## Approximate NLO corrections

- partial-fractioning the Eikonal

$$\frac{p_i \cdot p_k}{p_i \cdot q \, p_k \cdot q} \rightarrow \frac{1}{p_i \cdot q} \frac{p_i \cdot p_k}{(p_i + p_k) \cdot q} + \frac{1}{p_k \cdot q} \frac{p_i \cdot p_k}{(p_i + p_k) \cdot q}$$

The first term contains the soft singularity associated with the region collinear to  $p_i$ , while the second that collinear to  $p_k$ .

- with this, we get

$$dV_{ik} \rightarrow dV_i = \omega d\omega \frac{d\Omega}{2\pi} \frac{\alpha_s}{2\pi} C_i W_{ijk}^i$$

a real-number-valued multiplicative factor of the soft gluon-emission correction in the large- $N_c$  limit


- combine with coll. limit to soft-collinear (dipole) splitting functions

# Higher-order corrections and parton branchings

## The heuristic view


## Radiative corrections as a branching process

- parton branchings are IR divergent, introduce a resolution parameter to regulate the branching process  $t_{\text{res}}$



- resolvable,  $t > t_c$ , finite

- include




- unresolvable,  $t < t_c$ , finite

- Assumption:** corrections from resolvable and unresolvable branchings add up to zero, true for divergent leading logarithms (KLN theorem), amounts to saying that integrated higher-order corrections vanish

⇒ parton branchings can be interpreted probabilistically, either a parton branches resolvably with a probability given by the resolvable branching process or it does not


## Radiative corrections as a branching process

- parton branchings are IR divergent, introduce a resolution parameter to regulate the branching process  $t_{\text{res}}$



- resolvable,  $t > t_c$ , finite

- include




- unresolvable,  $t < t_c$ , finite

- Assumption:** corrections from resolvable and unresolvable branchings add up to zero, true for divergent leading logarithms (KLN theorem), amounts to saying that integrated higher-order corrections vanish
- ⇒ parton branchings can be interpreted probabilistically, either a parton branches resolvably with a probability given by the resolvable branching process or it does not

→ same as nuclear decay


## Radiative corrections as a branching process

- parton branchings are IR divergent, introduce a resolution parameter to regulate the branching process  $t_{\text{res}}$



- resolvable,  $t > t_c$ , finite

- include



- unresolvable,  $t < t_c$ , finite

- Assumption:** corrections from resolvable and unresolvable branchings add up to zero, true for divergent leading logarithms (KLN theorem), amounts to saying that integrated higher-order corrections vanish
- ⇒ parton branchings can be interpreted probabilistically, either a parton branches resolvably with a probability given by the resolvable branching process or it does not

→ same as nuclear decay

## Radiative corrections as a branching process

- The consequence is Poisson statistics
  - let the branching probability be  $\lambda$
  - assume indistinguishable particles  $\rightarrow$  naïve probability for  $n$  emissions

$$P_{\text{naïve}}(n, \lambda) = \frac{\lambda^n}{n!}$$

- probability conservation (unitarity) implies a no-emission probability

$$P(n, \lambda) = \frac{\lambda^n}{n!} \exp\{-\lambda\} \quad \longrightarrow \quad \sum_{n=0}^{\infty} P(n, \lambda) = 1$$

- introduce Sudakov form factor  $\Delta = \exp\{-\lambda\}$

## Radiative corrections as a branching process

- branching probability for parton state at scale  $Q^2$  in collinear limit in terms of resolution variable  $t$

$$\lambda \rightarrow \int_t^{Q^2} d\bar{t} \frac{d}{d\bar{t}} \left[ \frac{\sigma_{n+1}(\bar{t})}{\sigma_n} \right] \approx \sum_{\text{jets}} \int_t^{Q^2} d\bar{t} \int dz \frac{\alpha_s}{2\pi\bar{t}} P(z)$$

- Altarelli-Parisi splitting functions  $P(z)$ , spin- and colour dependent

$$P_{qq}(z) = C_F \left[ \frac{2z}{1-z} + (1-z) \right] \quad P_{gq}(z) = T_R \left[ z^2 + (1-z)^2 \right]$$

$$P_{gg}(z) = C_A \left[ \frac{2z}{1-z} + z(1-z) \right] + (z \leftrightarrow 1-z)$$


- branching process conserves momentum, colour, and on-shellness



# Radiative corrections as a branching process

- branching probability for parton state at scale  $Q^2$  in collinear limit in terms of resolution variable  $t$

$$\lambda \rightarrow \int_t^{Q^2} d\bar{t} \frac{d}{d\bar{t}} \left[ \frac{\sigma_{n+1}(\bar{t})}{\sigma_n} \right] \approx \sum_{\text{jets}} \int_t^{Q^2} d\bar{t} \int dz \frac{\alpha_s}{2\pi\bar{t}} P(z)$$



- Altarelli-Parisi splitting functions  $P(z)$ , spin- and colour dependent

$$P_{qq}(z) = C_F \left[ \frac{2z}{1-z} + (1-z) \right] \quad P_{gq}(z) = T_R \left[ z^2 + (1-z)^2 \right]$$


$$P_{gg}(z) = C_A \left[ \frac{2z}{1-z} + z(1-z) \right] + (z \leftrightarrow 1-z)$$

- branching process conserves momentum, colour, and on-shellness

## Radiative corrections as a branching process

- branching probability for parton state at scale  $Q^2$  in collinear limit in terms of resolution variable  $t$

$$\lambda \rightarrow \int_t^{Q^2} d\bar{t} \frac{d}{d\bar{t}} \left[ \frac{\sigma_{n+1}(\bar{t})}{\sigma_n} \right] \approx \sum_{\text{jets}} \int_t^{Q^2} d\bar{t} \int dz \frac{\alpha_s}{2\pi\bar{t}} P(z)$$



- Altarelli-Parisi splitting functions  $P(z)$ , spin- and colour dependent

$$P_{qq}(z) = C_F \left[ \frac{2z}{1-z} + (1-z) \right] \quad P_{gq}(z) = T_R \left[ z^2 + (1-z)^2 \right]$$

$$P_{gg}(z) = C_A \left[ \frac{2z}{1-z} + z(1-z) \right] + (z \leftrightarrow 1-z)$$

- branching process conserves momentum, colour, and on-shellness

## The improved large- $N_c$ approximation

# Colour flow

- quark propagator in fundamental representation  $\delta_{ij}$  contains  $N_c = 3$  colour states
- gluon propagator in adjoint representation  $\delta^{ab}$  contains  $N_c^2 - 1 = 8$  colour states

using completeness relations

$$\underbrace{\delta^{ab}}_{\text{standard}} = 2 \text{Tr}(T^a T^b) = 2 T_{ij}^a T_{ji}^b = T_{ij}^a \underbrace{2 \delta_{ik} \delta_{jl}}_{\text{colour flow}} T_{lk}^b$$

## Colour flow


- quark propagator in fundamental representation  $\delta_{ij}$   
contains  $N_c = 3$  colour states
- gluon propagator in adjoint representation  $\delta^{ab}$   
contains  $N_c^2 - 1 = 8$  colour states

using completeness relations


$$\underbrace{\delta^{ab}}_{\text{standard}} = 2 \text{Tr}(T^a T^b) = 2 T_{ij}^a T_{ji}^b = T_{ij}^a \underbrace{2 \delta_{ik} \delta_{jl}}_{\text{colour flow}} T_{lk}^b$$

## Colour flow

- Quark-gluon vertex  $T_{ij}^a T_{kl}^a = \frac{1}{2} \left( \delta_{il} \delta_{jk} - \frac{1}{N_c} \delta_{ij} \delta_{kl} \right)$




- Gluon-gluon vertex  $f^{abc} T_{ij}^a T_{kl}^b T_{mn}^c = \delta_{il} \delta_{kn} \delta_{mj} - \delta_{in} \delta_{ml} \delta_{kj}$




# The improved large- $N_c$ approximation

- leading colour approximation

$$T_{ij}^a T_{kl}^a \rightarrow \frac{1}{2} \delta_{il} \delta_{jk} \quad \leftrightarrow$$



- this overestimates the colour charge of the quark:  
Consider process  $q \rightarrow qg$  attached to some larger diagram  $|\mathcal{M}|^2$



$$\propto T_{ij}^a T_{jk}^a = C_F \delta_{ik} \quad (\text{QCD, } N_c = 3)$$


$$\rightarrow \frac{1}{2} \delta_{il} \delta_{jm} \delta_{mj} \delta_{lk} = \frac{C_A}{2} \delta_{ik} \quad (N_c \rightarrow \infty)$$

- improved large- $N_c$  approx.: keep colour charge of quarks at  $C_F$


# The improved large- $N_c$ approximation

- leading colour approximation

$$T_{ij}^a T_{kl}^a \rightarrow \frac{1}{2} \delta_{il} \delta_{jk} \quad \leftrightarrow$$



- this overestimates the colour charge of the quark:  
Consider process  $q \rightarrow qg$  attached to some larger diagram  $|\mathcal{M}|^2$



$$\propto T_{ij}^a T_{jk}^a = C_F \delta_{ik} \quad (\text{QCD, } N_c = 3)$$

$$\rightarrow \frac{1}{2} \delta_{il} \delta_{jm} \delta_{mj} \delta_{lk} = \frac{C_A}{2} \delta_{ik} \quad (N_c \rightarrow \infty)$$

- improved large- $N_c$  approx.: keep colour charge of quarks at  $C_F$



# Monte-Carlo methods for parton showers

## — The veto algorithm

## Monte-Carlo methods: Poisson distributions

- assume branching process described by  $g(t)$
- branching can happen only if it has not happened already, must account for survival probability  $\leftrightarrow$  Poisson distribution

$$\mathcal{G}(t) = g(t)\Delta(t, t_0) \quad \text{where} \quad \Delta(t, t_0) = \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- if  $G(t)$  is known, then we also know the integral of  $g(t)$

$$\int_t^{t_0} dt' \mathcal{G}(t') = \int_t^{t_0} dt' \frac{d\Delta(t', t_0)}{dt'} = 1 - \Delta(t, t_0)$$

- can generate events by requiring  $1 - \Delta(t, t_0) = 1 - R$  ( $R \in [0, 1]$ )

$$t = G^{-1} \left[ G(t_0) + \log R \right]$$

# Veto algorithm – importance sampling for Poisson dists

Parton shower branching probability  $f(t) \propto \frac{\alpha_s(t)}{t} P(z)$

**Problem:** we do not know  $F(t)$

**Solution:** veto algorithm

- 1 find overestimate  $g(t) \geq f(t) \forall t \in [t_c, t_0]$ ,  
generate event according to

$$G(t) = g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- 2 accept with  $w(t) = f(t)/g(t)$
- 3 if rejected, continue starting from  $t$

## Veto algorithm – importance sampling for Poisson dists

Parton shower branching probability  $f(t) \propto \frac{\alpha_s(t)}{t} P(z)$

**Problem:** we do not know  $F(t)$

**Solution:** veto algorithm

- 1 find overestimate  $g(t) \geq f(t) \forall t \in [t_c, t_0]$ ,  
generate event according to

$$G(t) = g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- 2 accept with  $w(t) = f(t)/g(t)$
- 3 if rejected, continue starting from  $t$

# Veto algorithm – importance sampling for Poisson dists

## Does this give the correct distribution?

- probability for immediate acceptance of emission at scale  $t$

$$\frac{f(t)}{g(t)} g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- probability for acceptance after one rejection

$$\frac{f(t)}{g(t)} g(t) \int_t^{t_0} dt_1 \exp \left\{ - \int_t^{t_1} dt' g(t') \right\} \left( 1 - \frac{f(t_1)}{g(t_1)} \right) g(t_1) \exp \left\{ - \int_{t_1}^{t_0} dt' g(t') \right\}$$

- For  $n$  rejections we obtain  $n$  nested integrals  $\int_t^{t_0} \int_{t_1}^{t_0} \dots \int_{t_{n-1}}^{t_0}$
- disentangling yields  $1/n!$ , summing over all possible rejections gives

$$\begin{aligned} & f(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\} \sum_{n=0}^{\infty} \frac{1}{n!} \left[ \int_t^{t_0} dt' [g(t') - f(t')] \right]^n \\ &= f(t) \exp \left\{ - \int_t^{t_0} dt' f(t') \right\} \end{aligned}$$

# Veto algorithm – importance sampling for Poisson dists

## Does this give the correct distribution?

- probability for immediate acceptance of emission at scale  $t$

$$\frac{f(t)}{g(t)} g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- probability for acceptance after one rejection

$$\frac{f(t)}{g(t)} g(t) \int_t^{t_0} dt_1 \exp \left\{ - \int_t^{t_1} dt' g(t') \right\} \left( 1 - \frac{f(t_1)}{g(t_1)} \right) g(t_1) \exp \left\{ - \int_{t_1}^{t_0} dt' g(t') \right\}$$

- For  $n$  rejections we obtain  $n$  nested integrals  $\int_t^{t_0} \int_{t_1}^{t_0} \dots \int_{t_{n-1}}^{t_0}$
- disentangling yields  $1/n!$ , summing over all possible rejections gives

$$\begin{aligned} f(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\} \sum_{n=0}^{\infty} \frac{1}{n!} \left[ \int_t^{t_0} dt' [g(t') - f(t')] \right]^n \\ = f(t) \exp \left\{ - \int_t^{t_0} dt' f(t') \right\} \end{aligned}$$

# Veto algorithm – importance sampling for Poisson dists

## Does this give the correct distribution?

- probability for immediate acceptance of emission at scale  $t$

$$\frac{f(t)}{g(t)} g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- probability for acceptance after one rejection

$$\frac{f(t)}{g(t)} g(t) \int_t^{t_0} dt_1 \exp \left\{ - \int_t^{t_1} dt' g(t') \right\} \left( 1 - \frac{f(t_1)}{g(t_1)} \right) g(t_1) \exp \left\{ - \int_{t_1}^{t_0} dt' g(t') \right\}$$

- For  $n$  rejections we obtain  $n$  nested integrals  $\int_t^{t_0} \int_{t_1}^{t_0} \dots \int_{t_{n-1}}^{t_0}$

- disentangling yields  $1/n!$ , summing over all possible rejections gives

$$\begin{aligned} f(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\} \sum_{n=0}^{\infty} \frac{1}{n!} \left[ \int_t^{t_0} dt' [g(t') - f(t')] \right]^n \\ = f(t) \exp \left\{ - \int_t^{t_0} dt' f(t') \right\} \end{aligned}$$

# Veto algorithm – importance sampling for Poisson dists

## Does this give the correct distribution?

- probability for immediate acceptance of emission at scale  $t$

$$\frac{f(t)}{g(t)} g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- probability for acceptance after one rejection

$$\frac{f(t)}{g(t)} g(t) \int_t^{t_0} dt_1 \exp \left\{ - \int_t^{t_1} dt' g(t') \right\} \left( 1 - \frac{f(t_1)}{g(t_1)} \right) g(t_1) \exp \left\{ - \int_{t_1}^{t_0} dt' g(t') \right\}$$

- For  $n$  rejections we obtain  $n$  nested integrals  $\int_t^{t_0} \int_{t_1}^{t_0} \dots \int_{t_{n-1}}^{t_0}$
- disentangling yields  $1/n!$ , summing over all possible rejections gives

$$\begin{aligned} & f(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\} \sum_{n=0}^{\infty} \frac{1}{n!} \left[ \int_t^{t_0} dt' [g(t') - f(t')] \right]^n \\ &= f(t) \exp \left\{ - \int_t^{t_0} dt' f(t') \right\} \quad \checkmark \end{aligned}$$



# Veto algorithm – importance sampling for Poisson dists

## Does this give the correct distribution?

- probability for immediate acceptance of emission at scale  $t$

$$\frac{f(t)}{g(t)} g(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\}$$

- probability for acceptance after one rejection

$$\frac{f(t)}{g(t)} g(t) \int_t^{t_0} dt_1 \exp \left\{ - \int_t^{t_1} dt' g(t') \right\} \left( 1 - \frac{f(t_1)}{g(t_1)} \right) g(t_1) \exp \left\{ - \int_{t_1}^{t_0} dt' g(t') \right\}$$

- For  $n$  rejections we obtain  $n$  nested integrals  $\int_t^{t_0} \int_{t_1}^{t_0} \dots \int_{t_{n-1}}^{t_0}$
- disentangling yields  $1/n!$ , summing over all possible rejections gives

$$\begin{aligned} & f(t) \exp \left\{ - \int_t^{t_0} dt' g(t') \right\} \sum_{n=0}^{\infty} \frac{1}{n!} \left[ \int_t^{t_0} dt' [g(t') - f(t')] \right]^n \\ &= f(t) \exp \left\{ - \int_t^{t_0} dt' f(t') \right\} \quad \checkmark \end{aligned}$$

# Veto algorithm

## What have we achieved?

- we have generated a parton branching (real resolved emission) according to

$$f(t) \Delta(t, t_0) \quad \text{with} \quad \Delta(t, t_0) = \exp \left\{ - \int_t^{t_0} dt' f(t') \right\}$$

$$f(t) \equiv f(t, z) = \frac{\alpha_s}{2\pi t} P(z)$$

- the no-branching probability implies a virtual correction (including unresolved real emissions) of  $\Delta(t_c, t_0)$

**Note:** The Sudakov form factor  $\Delta$  resums logs to all orders

$$d\hat{\sigma}_{\text{NLO}}^{\text{approx}} = d\hat{\sigma}_n \left[ \Delta(t_c, t_0) + \int_{t_c}^{t_0} dt f(t) \Delta(t, t_0) \right]$$

# Veto algorithm

## What have we achieved?

- we have generated a parton branching (real resolved emission) according to

$$f(t) \Delta(t, t_0) \quad \text{with} \quad \Delta(t, t_0) = \exp \left\{ - \int_t^{t_0} dt' f(t') \right\}$$

$$f(t) \equiv f(t, z) = \frac{\alpha_s}{2\pi t} P(z)$$

- the no-branching probability implies a virtual correction (including unresolved real emissions) of  $\Delta(t_c, t_0)$

**Note:** The Sudakov form factor  $\Delta$  resums logs to all orders

$$d\hat{\sigma}_{\text{NLO}}^{\text{approx}} = d\hat{\sigma}_n \left[ \Delta(t_c, t_0) + \int_{t_c}^{t_0} dt f(t) \Delta(t, t_0) \right]$$

# Veto algorithm

## What have we achieved?

- we have generated a parton branching (real resolved emission) according to

$$f(t) \Delta(t, t_0) \quad \text{with} \quad \Delta(t, t_0) = \exp \left\{ - \int_t^{t_0} dt' f(t') \right\}$$

$$f(t) \equiv f(t, z) = \frac{\alpha_s}{2\pi t} P(z)$$

- the no-branching probability implies a virtual correction (including unresolved real emissions) of  $\Delta(t_c, t_0)$

**Note:** The Sudakov form factor  $\Delta$  resums logs to all orders

$$d\hat{\sigma}_{\text{NLO}}^{\text{approx}} = d\hat{\sigma}_n \left[ \Delta(t_c, t_0) + \int_{t_c}^{t_0} dt f(t) \Delta(t, t_0) \right]$$

## Veto algorithm – iteration

- 1) consider a set of  $n$  partons at scale  $t_0$ , which evolve collectively  
Sudakovs factorise, schematically

$$\Delta(t, t_0) = \prod_{i=1}^n \Delta_i(t, t_0), \quad \Delta_i(t, t_0) = \prod_{j=q,g} \Delta_{i \rightarrow j}(t, t_0)$$

- 2) find new scale  $t$  where next branching occurs using veto algorithm
  - generate  $t$  using overestimate  $g_{ab}(t) \propto \alpha_s^{\max} P_{ab}^{\max}(z)$
  - determine “winner” parton  $i$  and select new flavor  $j$
  - accept point with weight  $f_{ab}/g_{ab} = \alpha_s(k_T^2) P_{ab}(z) / \alpha_s^{\max} P_{ab}^{\max}(z)$
- 3) construct splitting kinematics and update event record
- 4) continue until  $t < t_c$ ,  $t_c$  infrared cut-off

## Veto algorithm – iteration

- 1) consider a set of  $n$  partons at scale  $t_0$ , which evolve collectively  
Sudakovs factorise, schematically

$$\Delta(t, t_0) = \prod_{i=1}^n \Delta_i(t, t_0), \quad \Delta_i(t, t_0) = \prod_{j=q,g} \Delta_{i \rightarrow j}(t, t_0)$$

- 2) find new scale  $t$  where next branching occurs using veto algorithm
  - generate  $t$  using overestimate  $g_{ab}(t) \propto \alpha_s^{\max} P_{ab}^{\max}(z)$
  - determine “winner” parton  $i$  and select new flavor  $j$
  - accept point with weight  $f_{ab}/g_{ab} = \alpha_s(k_T^2) P_{ab}(z) / \alpha_s^{\max} P_{ab}^{\max}(z)$
- 3) construct splitting kinematics and update event record
- 4) continue until  $t < t_c$ ,  $t_c$  infrared cut-off

## Veto algorithm – iteration

- 1) consider a set of  $n$  partons at scale  $t_0$ , which evolve collectively  
Sudakovs factorise, schematically

$$\Delta(t, t_0) = \prod_{i=1}^n \Delta_i(t, t_0), \quad \Delta_i(t, t_0) = \prod_{j=q,g} \Delta_{i \rightarrow j}(t, t_0)$$

- 2) find new scale  $t$  where next branching occurs using veto algorithm
  - generate  $t$  using overestimate  $g_{ab}(t) \propto \alpha_s^{\max} P_{ab}^{\max}(z)$
  - determine “winner” parton  $i$  and select new flavor  $j$
  - accept point with weight  $f_{ab}/g_{ab} = \alpha_s(k_T^2) P_{ab}(z) / \alpha_s^{\max} P_{ab}^{\max}(z)$
- 3) construct splitting kinematics and update event record
- 4) continue until  $t < t_c$ ,  $t_c$  infrared cut-off

## Veto algorithm – iteration

- 1) consider a set of  $n$  partons at scale  $t_0$ , which evolve collectively  
Sudakovs factorise, schematically



$$\Delta(t, t_0) = \prod_{i=1}^n \Delta_i(t, t_0), \quad \Delta_i(t, t_0) = \prod_{j=q,g} \Delta_{i \rightarrow j}(t, t_0)$$

- 2) find new scale  $t$  where next branching occurs using veto algorithm
  - generate  $t$  using overestimate  $g_{ab}(t) \propto \alpha_s^{\max} P_{ab}^{\max}(z)$
  - determine “winner” parton  $i$  and select new flavor  $j$
  - accept point with weight  $f_{ab}/g_{ab} = \alpha_s(k_T^2) P_{ab}(z) / \alpha_s^{\max} P_{ab}^{\max}(z)$
- 3) construct splitting kinematics and update event record
- 4) continue until  $t < t_c$ ,  $t_c$  infrared cut-off




## Effects of the parton shower

# Effects of the parton shower




- Thrust and Durham  $2 \rightarrow 3$ -jet rate in  $e^+e^- \rightarrow \text{hadrons}$
- hadronisation region to the right (left) in left (right) plot

# Effects of the parton shower



- Drell-Yan lepton pair production at Tevatron
- if hard cross section computed at leading order, then parton shower is only source of transverse momentum
- starting scale of evolution chosen as  $Q^2 = m_W^2$

# Effects of the parton shower



# Recap

## This lecture:

- parton showers encode approximate higher-order corrections  
→ build upon universal soft-collinear approximation  
(Altarelli-Parisi splitting functions, large- $N_c$ , spin-averaged)
- implemented as a statistical branching process, ordered in evolution variable  $t$  ( $k_T^2$ ,  $\tilde{q}^2$ , etc.)
- produce resolved final state up to scale  $t_{\text{res}} \approx \Lambda_{\text{QCD}}$   
→ further evolution needs hadrons as degrees of freedom

## Next lectures:

- limitations of parton showers and how to overcome them