

## Preferred review

# Clinical applications of antimicrobial host proteins lactoperoxidase, lysozyme and lactoferrin in xerostomia: efficacy and safety

J Tenovuo

*Institute of Dentistry and Turku Immunology Centre, University of Turku, Turku, Finland*

**Innate human salivary defence proteins, lysozyme, lactoferrin and peroxidase, are known to exert a wide antimicrobial activity against a number of bacterial, viral and fungal pathogens *in vitro*. Therefore, these proteins, alone or in combinations, have been incorporated as preservatives in foods and pharmaceuticals as well as in oral health care products to restore salivas' own antimicrobial capacity in patients with dry mouth. These antimicrobials used in oral health care products, such as dentifrices, mouth-rinses, moisturizing gels and chewing gums, have been purified from bovine colostrum. In this review I critically evaluate the clinical efficacy and safety of this kind of preventive approach against various oral diseases and symptoms.**

*Oral Diseases* (2002) 8, 23–29

**Keywords:** lactoperoxidase; lysozyme; lactoferrin; dentifrices; xerostomia; saliva

## Introduction

Adequate flow of saliva is a prerequisite for good oral health. Saliva not only makes oral functions possible but it also protects oral tissues from noxious agents derived from microorganisms, food or drugs. Because the human mouth is a major route of entry for foreign materials into the body, saliva with its protective capacity forms an important part of a first line of defence against exogenous, often harmful agents. A proper flow of saliva is vital not only for oral health but, obviously, also to the general health of humans (Herrera, Lyons and Johnson, 1988; Rhodus and Brown, 1990).

Hyposalivation often leads to many oral problems: xerostomia with all its symptoms, rapid dental decay, increased susceptibility to fungal infections, mucosal infections, etc. All these impair the patients' well-being, particularly because dental and medical professionals still often fail to make proper diagnosis in this interdisciplinary area (Nederfors, 2000). In spite of their known clinical significance the prevalence of hyposalivation and xerostomia is largely unknown. Xerostomia is usually experienced when the salivary flow rate is reduced by about 50% in test subjects with normal salivary flow (Dawes, 1987). However, normal flow is an individual feature and on a population level subjective sensation of oral dryness is only poorly associated with hyposalivation, as quantitated by objective measurements (Hay *et al*, 1998). However, it seems that both hyposalivation and xerostomia increase by age, are more prevalent among women than men, and are often associated with frequent use of drugs (Närhi, 1994; Nederfors, 2000). Depending on how the questions of oral dryness are formulated, the prevalence figures for xerostomia vary in adults and elderly usually between 10 and 40% (Nederfors, 2000).

Saliva protects oral tissues in many ways. A constant flow of saliva eliminates microorganisms from the oral cavity but saliva also contains many innate or acquired defence systems (reviewed, e.g. by Brandtzaeg, 1989; Lagerlöf and Oliveby, 1994; Tenovuo, 1998; Lenander-Lumikari and Loimaranta, 2000). A number of studies have tried to find a clinically significant association between selected salivary agents and caries prevalence or incidence but the results are controversial. This may be explained by several reasons, e.g. various salivary agents interact with each other in a complicated way (Lenander-Lumikari and Loimaranta, 2000). Also inadequate immune defence (usually IgA deficiency) can be compensated by an enhanced non-immunoglobulin activity (Arnold *et al*, 1979; Kirstilä *et al*, 1994b) or by compensatory IgM response (Norhagen Engström *et al*, 1989). These examples of concerted action of salivary

Correspondence: Dr J Tenovuo, Institute of Dentistry, University of Turku, Lemminkäisenkatu 2, FIN-20520 Turku, Finland. Tel.: +358 2 3338340, Fax: +358 2 3338300, E-mail: [jorma.tenovuo@utu.fi](mailto:jorma.tenovuo@utu.fi)  
Received 3 July 2001; revised 7 November 2001; accepted 20 November 2001  
Conflict of interest: none

antimicrobials assure host-derived protection as long as an appropriate amount of saliva is secreted.

### Enhancement of the antimicrobial capacity in saliva

In connection with hyposalivation, any increase in the flow rate inevitably results in an increase in the output of saliva-borne factors and therefore all methods to increase the flow rate are encouraged. This article, however, reviews attempts to enhance or restore saliva's own antimicrobial capacity by commercially available oral health care products. The antimicrobial host proteins most widely used in these products are lysozyme, lactoferrin and lactoperoxidase. To make lactoperoxidase (LP) enzyme antimicrobial, its substrates, thiocyanate ( $\text{SCN}^-$ ) and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), are also included in the products. Hydrogen peroxide is usually generated *in situ* by a glucose-glucose oxidase system.

A large amount of scientific literature exists of how these non-immunoglobulin components work separately and together (for reviews, see Tenovuo, 1998; Lenander-Lumikari and Loimaranta, 2000). The reported antimicrobial mechanisms are listed in Table 1. An interesting point is that all these three antimicrobial agents act in an additive way and even synergistic interactions have been described (Tenovuo *et al*, 1982a; Lassiter *et al*, 1987; Soukka, Lumikari and Tenovuo, 1991a,b; Lenander-Lumikari, Månson-Rahemtulla and Rahemtulla, 1992). Therefore, it seems sensible to combine them also in oral health care products.

### Products with antimicrobial proteins

It is relatively simple to add lysozyme and lactoferrin into oral health care products, such as toothpastes and

**Table 1** Antimicrobial mechanisms reported for lysozyme, lactoferrin and peroxidase enzyme systems. A more detailed description of these mechanisms with references is found in Tenovuo (1989)

#### Lysozyme

- Muramidase activity (lysis of peptidoglycan layer)
- Cationic-dependent activation of bacterial autolysins
- Aggregation of bacteria
- Inhibition of bacterial adhesion to tooth surfaces
- Inhibition of bacterial glucose uptake and acid production
- De-chaining of streptococci

#### Lactoferrin

- Deprivation of iron from bacteria (bacteriostatic action)
- Bactericidal action by direct contact of iron-free form of lactoferrin (apo-LF)
- Inhibition of bacterial adhesion to tooth surfaces
- Aggregation of bacteria
- Activation of phagocytic cells

#### Salivary peroxidase systems

- Inactivation of bacterial glycolytic enzymes
- Inhibition of bacterial glucose uptake and acid production
- Inhibition of bacterial transport of amino acids
- Damage of the bacterial cell wall
- Killing of bacteria (with  $\text{I}^-$  as a substrate instead of  $\text{SCN}^-$ )
- Inhibition of bacterial adhesion to saliva-coated hydroxyapatite

mouth-rinses. These antimicrobial proteins can be purified in a cost-effective way from bovine colostrum and commercial products comprising such bovine lysozyme and lactoferrin preparations include at least Biotene<sup>®</sup>, Oralbalance<sup>®</sup>, BioXtra<sup>®</sup> and Zendium Saliva<sup>®</sup>. The latter two contain 'whey extract' or 'colostrum' and therefore also bovine immunoglobulins, which, however, are not reported by the manufacturer to be specific against human oral pathogens. It is possible that these products contain also colostrum-containing growth factors, such as IGF-1, TGF- $\beta$ 1 and TGF- $\beta$ 2, which are known to be present in many colostrum products (Wei, Loimaranta, Tenovuo *et al*, 2002). Whether these have any effects on mucosal tissues is not known.

Human lactoferrin, or its antimicrobial 25-residue peptide fragment lactoferricin B, has recently been cloned by fermentation of *Aspergillus niger* (Headon, 2000). In future, it is likely that bovine milk derived lactoferrin will be replaced by recombinant human lactoferrin, which is reported to be active and safe for human clinical use (Headon, 2000). The possible difference in the efficacy of iron-free versus iron-saturated lactoferrin in oral health care products is not yet known.

The mimicking of salivary peroxidase systems is somewhat more complicated than lysozyme or lactoferrin. Both salivary peroxidase and myeloperoxidase, the two peroxidase enzymes in human mixed saliva, have been purified but only for research purposes. Large-scale purification from human saliva (salivary peroxidase) or from human polymorphonuclear leukocytes (myeloperoxidase) is difficult and far too expensive for commercial purposes. Therefore, for commercial purposes bovine lactoperoxidase (purified from milk or colostrum) has been used because this enzyme is structurally and catalytically very close to human salivary peroxidase (Månsson-Rahemtulla *et al*, 1988). Thiocyanate ( $\text{SCN}^-$ ) is added as potassium thiocyanate but  $\text{H}_2\text{O}_2$  is generated *in situ* in the mouth by a glucose-glucose oxidase system added to these oral hygiene products. The lactoperoxidase system is incorporated into commercial products like Biotene<sup>®</sup>, BioXtra<sup>®</sup>, Zendium Saliva<sup>®</sup> and Oralbalance<sup>®</sup>. Many of these comprise not only a toothpaste, but also a non-alcoholic mouth-rinse, a xylitol-flavoured chewing gum, moisturizing gels and/or denture adhesives for patients with dry mouth problems. The lactoperoxidase system retains its antibacterial activity for 1–2 years (Bosch, Van Doorne and De Vries, 2000) but data of the possible reduction in lysozyme or lactoferrin functions are not available. All lactoperoxidase system containing products are non-foaming because detergents inactivate the enzymes (Hoogendoorn, 1985).

### Do these products work *in vivo*?

As early as 1983 Månsson-Rahemtulla *et al* showed that it was indeed possible to design a mouth-rinse which *in vivo* elevates the hypothiocyanite levels to bacteriostatic concentrations. The volume of the rinse, its pH and the amount of  $\text{H}_2\text{O}_2$  generated were found to the

critical determinants, but at pH 5.5–6.5 high HOSCN/OSCN<sup>-</sup> concentrations, as high as >200 µM, could be generated *in vivo* (Månsson-Rahemtulla *et al*, 1983). The average whole saliva concentrations of HOSCN/OSCN<sup>-</sup> range from 30 to 70 µM (mean 44 µM) in resting whole saliva with much lower concentrations (mean 5 µM) in paraffin-stimulated whole salivas (Tenovuo, Pruitt and Thomas, 1982b). One of the first products, Biotene<sup>®</sup> toothpaste, was later shown to increase human whole saliva hypothiocyanite levels up to a mean of 200 µM concentration within 20 min after brushing (Lenander-Lumikari, Mikola and Tenovuo, 1993). No such effect was found with a non-lactoperoxidase/non-H<sub>2</sub>O<sub>2</sub> generating toothpaste. Daily use of Biotene<sup>®</sup> toothpaste for 1 month also resulted in a slight increase in the resting levels of salivary hypothiocyanite (Lenander-Lumikari *et al*, 1993).

Although the number of studies actually giving the *in vivo* concentrations of hypothiocyanite is limited, it seems that it is possible to significantly elevate the salivary levels of hypothiocyanite, the antimicrobial agent in saliva. However, no data are available of any increased concentrations of salivary or plaque lysozyme and lactoferrin in connection with the use of these products.

One of the advantages of peroxidase-containing products in patients with dry mouth is that administered lactoperoxidase-enzyme rapidly consumes intraorally generated H<sub>2</sub>O<sub>2</sub> (by aerobic bacteria) whose accumulation in the mouth could be detrimental to oral mucosal cells (Hänström, Johansson and Carlsson, 1983; Tenovuo and Larjava, 1984).

### Antimicrobial effects

There are a number of studies which describe the antimicrobial mechanisms of lysozyme, lactoferrin and HOSCN/OSCN<sup>-</sup> (see review by Tenovuo, 1989 and Table 1). Most of these studies are done *in vitro* using *Streptococcus mutans* as a target bacterium; also, the effects against periodontal pathogens have been published. In most conditions which imitate the *in vivo* situation in oral fluids, the effects are bacteriostatic (Pruitt and Reiter, 1985) (Table 1), but the spectrum against various microbial pathogens is large (Table 2). The observations with the oral health care products with host proteins parallel the above findings *in vitro*. In laboratory conditions Biotene<sup>®</sup> was superior to Zendium<sup>®</sup> to inhibit the growth of *S. mutans*, *S. sobrinus* and *Lactobacillus casei* and the Biotene<sup>®</sup> effect was comparable with that of chlorhexidine (1%)–sodium fluoride (0.2%) gel. No antimicrobial effect was observed with Biotene<sup>®</sup> in which the LP-system was inactivated (Tenovuo, Lumikari and Soukka, 1991).

*In vivo* studies using a commercial Zendium<sup>®</sup> toothpaste have yielded conflicting results in relation to prevention of oral diseases. Human studies have indicated slight positive effects regarding plaque accumulation and prevention of gingivitis and dental caries (Koch, Edlund and Hoogendoorn, 1973; Koch and Strand, 1979; Rotgans and Hoogendoorn, 1979;

**Table 2** Selected oral or orally transmitted pathogens which are susceptible to peroxidase-mediated inhibition or to inhibition by lactoferrin or lysozyme

Gram-positive bacteria
Mutans streptococci
Lactobacilli
Gram-negative bacteria
<i>Actinobacillus actinomycetemcomitans</i>
<i>Porphyromonas gingivalis</i>
<i>Fusobacterium nucleatum</i>
<i>Listeria monocytogenes</i>
<i>Salmonella typhimurium</i>
<i>Escherichia coli</i>
Viruses
Human immunodeficiency virus (HIV)
Herpes simplex type 1
Respiratory syncytial virus (RSV)
Yeasts
<i>Candida albicans</i>

Many of the observations are only from *in vitro* experiments and depend on the concentration of the inhibitory agent. See reviews by Pruitt and Reiter (1985), Tenovuo (1998) and Lenander-Lumikari and Loimaranta (2000).

Schoenfeld *et al*, 1983; Midda and Cooksey, 1986). However, Afseth and Rølla (1983) found no reduction in plaque formation or acidogenicity in subjects using Zendium<sup>®</sup> and Moran and Addy (1984) did not detect any inhibitory effect by Zendium<sup>®</sup> against *S. mutans*. Quite surprisingly, salivary or plaque levels of hypothiocyanite were not reported in any of these studies.

Another toothpaste, Biotene<sup>®</sup>, containing all the peroxidase system components (LP, KSCN, H<sub>2</sub>O<sub>2</sub>-generating system) and lysozyme and lactoferrin has been launched in Europe some 10 years ago after being available in the USA. As stated above, this toothpaste effectively generates hypothiocyanite *in vivo* (Lenander-Lumikari *et al*, 1993) but no Biotene<sup>®</sup>-induced antibacterial effects against total streptococci, mutans streptococci, lactobacilli or total anaerobic flora were detected (Lenander-Lumikari *et al*, 1993; Kirstilä *et al*, 1994a), neither in whole saliva nor in dental plaque. Also plaque pH, acidogenicity or lactic acid production were unaffected by Biotene<sup>®</sup> (Kirstilä *et al*, 1994a) in a double-blind crossover study with 20 volunteers with normal salivary flow rate.

However, one of the leading ideas in designing these products has been the purpose to (partially) restore saliva's own antimicrobial capacity among patients who suffer from dry mouth. This idea seems sound to add physiological salivary antimicrobial agents into the mouth that lacks saliva-mediated protection. Therefore, studies with patients with hyposalivation (and xerostomia) are of greater interest than studies with those who have normal salivation and, consequently, normal outputs of saliva-borne antimicrobials. Kirstilä *et al* (1996) observed during a 4-week clinical trial that out of 20 patients with Sjögren's syndrome, 16 experienced a significant relief of their dry mouth symptoms (xerostomia) although no increase in salivary flow rate or its hypothiocyanite levels were found. No major changes

occurred in salivary microflora either. Other studies with patients having, e.g. radiation-induced hyposalivation and xerostomia have shown Biotene® to improve gingival health via reduced formation of supragingival plaque compared with a control toothpaste (Banoczy *et al.*, 1994; van Steenberghe *et al.*, 1994; Toljanic *et al.*, 1996). Some studies have combined Biotene® toothpaste with Oralbalance® gel with good results in subjective relief of radiation-induced xerostomic symptoms (Epstein *et al.*, 1999; Warde, Kroll, O'Sullivan *et al.*, 2000) but the lack of appropriate controls makes it impossible to rule out the placebo effect.

To facilitate the efficacy and compatibility of these products in a dry mouth, toothpastes are not foaming but lack the mucosa-irritating effect of sodium lauryl sulphate (Herlofson and Barkvoll, 1994). In fact, the early assumption that the antimicrobial proteins in these toothpastes may have inhibited the appearance of aphthous ulcers (Hoogendoorn and Piessens, 1987) may be explained by the absence of ulcer-provoking sodium lauryl sulphate in all these antimicrobial toothpastes (Brokstad-Herlofson and Barkvoll, 1996). This is, however, not the whole truth because in carefully controlled double-blind crossover studies the Zendium® mouth-rinse has been found to reduce the incidence of aphthous ulcers over 50% when compared with sodium lauryl sulphate-free placebo rinse (Fridh and Koch, 1999). The mechanism of inhibition is unclear, as the etiology of aphthous ulcers is not known.

The mouth-rinses with host proteins are non-alcoholic, to be less irritable particularly to dry oral mucosal surfaces. Banoczy *et al.* (1994) reported enhanced keratinization and decreased number of inflammatory cells in both buccal and mucosal smears of xerostomic patients who had used a Biotene/Oralbalance® combination for 1 month. The only brief report of the benefits of Biotene® mouth-rinse is by Mulligan, Navazesh and Slots (1992) who found regular use of this mouth-rinse to notably decrease the total salivary microbial counts *in vivo*. However, Biotene® mouth-rinse has been observed to cause demineralization of a presoftened and also of a sound tooth enamel *in vitro* (Kielbassa, Shohadai and Schulte-Mönting, 2000). This is obviously because of the relatively low pH (5.15), high acid content and non-fluoride content of this mouth-rinse. Therefore, as it is now available, it can hardly be recommended to dentate patients with problems of dry mouth.

To conclude, study designs and small number of subjects in various clinical reports do not allow any direct comparisons with other type of clinical regimens. However, a rough estimate is that in human trials the reduction in plaque accumulation and gingivitis is 10–20%, but as high as 50–90% in the occurrence of aphthous ulcers (Hoogendoorn, 1985; Banoczy *et al.*, 1994; Kirstilä *et al.*, 1996).

### Future aspects

It is likely that these products can be further improved, e.g. by choosing more efficient substrates for peroxid-

ases. The oxidation products of iodide, for example, are much more potent antimicrobials against periodontal pathogens, such as *Actinobacillus actinomycetemcomitans* (Ihalin *et al.*, 1998) as well as *Porphyromonas gingivalis* and *Fusobacterium nucleatum* (Ihalin *et al.*, 2001). Also streptococci and *Candida albicans* are more prone to I<sup>-</sup> oxidation than to hypothiocyanite (Courtois *et al.*, 1995; Bosch *et al.*, 2000; Ihalin *et al.*, 2001). The major problem with iodide in connection with LP-system is that even relatively small concentrations of salivary thiocyanate abolish the bactericidal effect of the LP/I<sup>-</sup>/H<sub>2</sub>O<sub>2</sub> systems as SCN<sup>-</sup> is the preferred substrate for oral peroxidases *in vivo* (van Dalen *et al.*, 1997; Ihalin *et al.*, 2001). An interesting new approach to overcome the blocking effect of SCN<sup>-</sup> is to replace LP in oral hygiene products by horseradish peroxidase, which does not bind or oxidize SCN<sup>-</sup> at neutral pH but readily oxidizes iodide (Ihalin *et al.*, unpublished data). In future, the application of these antimicrobial components may be more effective if encapsulated into reactive liposomes, which have been demonstrated to be good vehicles to exploit horseradish peroxidase, lactoperoxidase, glucose oxidase and lactoferrin against oral streptococci and dental caries (Hill *et al.*, 1997; Martinez-Gomis *et al.*, 1999).

Also interestingly, in a double-blind placebo-controlled study on 140 subjects, positive clinical results against dental plaque and gingivitis have been obtained with a toothpaste supplemented by only SCN<sup>-</sup> and H<sub>2</sub>O<sub>2</sub>, without any peroxidase enzyme (Rosin, Kocher and Kramer, 2001). This is likely because of the enhancement of lysozyme action by SCN<sup>-</sup> (Pollock *et al.*, 1981) and by the positive interaction of SCN<sup>-</sup> with carbamide perhydrate present in these test dentifrices. This compound produces urea, which facilitates SCN<sup>-</sup> penetration into the oral mucosa. Thiocyanate is known to enhance cell proliferation and regeneration, e.g. in wound healing (Kramer and Böhländ, 1996), which might explain its value against gingival inflammation.

It also seems beneficial to combine peroxidase-mediated inhibition with specific IgG antibodies against mutans streptococci (Tenovou *et al.*, 1982a; Loimaranta, Tenovou and Korhonen, 1998). Thus, future products may not only comprise innate host proteins but also specific antibodies against oral pathogens. It should also be stressed that fluoride ions and the peroxidase systems act in an additive manner (Lenander-Lumikari *et al.*, 1997). To summarize, future products obviously comprise an array of various host-related antimicrobials at the expense of synthetic agents, such as chlorhexidine and hexetidine. This approach is also likely to be more physiological, safe and suitable for daily use as no harmful side-effects of host proteins have been described to human oral cell lines (Hänström *et al.*, 1983; Tenovou and Larjava, 1984). Although solely bovine milk derived proteins have been used so far in commercial applications, no short- (e.g. allergy) or long-term (e.g. autoimmunity) adverse effects have been observed. This may result from the immunological similarity, although not complete identity, of these proteins in cows and humans (Månsson-Rahemtulla, Rahemtulla and

Humphreys-Beher, 1990; Headon, 2000). Thus, the described enhancement or restoration of saliva's natural antimicrobial systems could be a useful and safe supplement to maintain good oral health.

## Conclusions

1. Incorporation of bovine milk-derived antimicrobial proteins, particularly lysozyme, lactoferrin and lactoperoxidase (LP), into human oral health care products is a serious and commercially active attempt to enhance and restore saliva's own antimicrobial capacity.
2. There is strong evidence that LP remains active and functional in a dentifrice, producing high concentrations of the antimicrobial oxidation product hypothiocyanite (HOSCN/OSCN<sup>-</sup>) *in vivo*. No such evidence of activity exists for lysozyme or lactoferrin.
3. In clinical trials these antimicrobial toothpastes, mouth-rinses or moisturizing gels relieve the symptoms of xerostomia, but whether it is because of the antimicrobial proteins is not known.
4. There is no clear-cut clinical evidence that these antimicrobial oral health care products could enhance salivas' own defence capacity if salivary flow rate is within normal limits.
5. These products may partly restore the innate immunity of saliva in patients with hyposalivation but their clinical efficacy is still poorly documented.
6. No side-effects have been reported. Because the antimicrobial components are similar, although not completely identical, to those naturally present in human salivary secretions, there seems to be no contraindication for their daily use by patients with dry mouth or by patients suffering from aphthous ulcers. The only exception is mouth-rinse whose low pH and absence of fluoride may cause tooth demineralization, particularly among patients with dry mouth.

## Acknowledgements

Studies in the author's laboratory and clinic have been supported by the Finnish Dental Society, Turku University Central Hospital, the Academy of Finland and the Finnish Technology Development Centre (TEKES).

## References

Afseth J, Rølla G (1983). Clinical experiments with a toothpaste containing amyloglucosidase and glucose oxidase. *Caries Res* **17**: 474–475.

Arnold RR, Pruitt KM, Cole MF, Adamson JM, McGhee JR (1979). Salivary antibacterial mechanisms in immunodeficiency. In: Kleinberg I, Ellison SA, Mandel ID, eds. *Proceedings Saliva and Dental caries*. Spec Suppl. Microbiol Abstract. Information Retrieval: New York, pp 449–462.

Banoczy JB, Dombi C, Czegledly A, Sari K (1994). A clinical study with lactoperoxidase containing gel and toothpaste in patients with dry mouth syndrome. *J Clin Dent* **5**: 65–69.

Bosch EH, Van Doorne H, De Vries S (2000). The lactoperoxidase system: the influence of iodide and the chemical and antimicrobial stability over the period of about 18 months. *J Appl Microbiol* **89**: 215–224.

Brandtzaeg P (1989). Salivary immunoglobulins. In: Tenovuo, J, ed. *Human saliva: clinical chemistry and microbiology*, Vol. II. CRC Press, Inc.: Boca Raton, FL, pp. 1–54.

Brokstad-Herlofson B, Barkvoll P (1996). The effect of two toothpaste detergents on the frequency of recurrent aphthous ulcers. *Acta Odontol Scand* **54**: 150–153.

Courtois P, Vanden Abbeele A, Amrani N, Pourtois M (1995). *Streptococcus sanguis* survival rates in the presence of lactoperoxidase-produced OSCN<sup>-</sup> and OI<sup>-</sup>. *Med Sci Res* **23**: 195–197.

van Dalen CJ, Whitehouse MW, Winterbourn CC, Kettle AJ (1997). Thiocyanate and chloride as competing substrates for myeloperoxidase. *Biochem J* **327**: 487–492.

Dawes C (1987). Physiological factors affecting salivary flow rate, oral sugar clearance, and the sensation of dry mouth in man. *J Dent Res* **66**: 648–653.

Epstein JB, Emerton S, Le ND, Stevenson-Moore P (1999). A double-blind crossover trial of Oral balance gel and Biotene toothpaste versus placebo in patients with xerostomia following radiation therapy. *Oral Oncol* **35**: 132–137.

Fridh G, Koch G (1999). Effect of a mouth-rinse containing amyloglucosidase and glucose oxidase on recurrent aphthous ulcers in children and adolescents. *Swed Dent J* **23**: 49–57.

Hänström L, Johansson A, Carlsson J (1983). Lactoperoxidase and thiocyanate protect cultured mammalian cells against hydrogen peroxide toxicity. *Med Biol* **61**: 268–274.

Hay EM, Thomas E, Pal B, Hajeer A, Chambers H, Silman AJ (1998). Weak association between subjective symptoms of and objective testing of dry eyes and dry mouth: results from a population based study. *Ann Rheum Dis* **57**: 20–24.

Headon DR (2000). Human lactoferrin: production at large scale, characterization and applications. In: Shimazaki, K, Tsuda H, Tomita M, Kuwata T, Perraudin J-P, eds. *Lactoferrin: structure, function and applications*. Elsevier Science B.V.: Amsterdam, pp 415–427.

Herlofson BB, Barkvoll P (1994). Sodium lauryl sulfate and recurrent aphthous ulcers. A preliminary study. *Acta Odontol Scand* **52**: 257–259.

Herrera JL, Lyons MF, Johnson LF (1988). Saliva: its role in health and disease. *J Clin Gastroenterol* **10**: 569–578.

Hill KJ, Kaszuba M, Creeth JE, Jones MN (1997). Reactive liposomes encapsulating a glucose oxidase–peroxidase system with antibacterial activity. *Biochim Biophys Acta* **1326**: 37–46.

Hoogendoorn H (1985). Activation of the salivary peroxidase antimicrobial system: clinical studies. In: Pruitt, KM, Tenovuo, J, eds. *The lactoperoxidase system, chemistry and biological significance*. Marcel Dekker Inc.: New York, pp. 217–227.

Hoogendoorn H, Piessens JP (1987). Treatment of aphthous patients by enhancement of the salivary peroxidase system. *J Oral Pathol* **16**: 425–427.

Ihalin R, Loimaranta V, Lenander-Lumikari M, Tenovuo J (1998). The effect of different (pseudo) halide substrates on peroxidase-mediated killing of *Actinobacillus actinomycetemcomitans*. *J Periodont Res* **33**: 421–427.

Ihalin R, Loimaranta V, Lenander-Lumikari M, Tenovuo J (2001). The sensitivity of *Porphyromonas gingivalis* and *Fusobacterium nucleatum* to different (pseudo) halide-peroxidase combinations compared with mutants streptococci. *J Med Microbiol* **50**: 42–48.

- Kielbassa AM, Shohadai SP, Schulte-Mönting J (2000). Effect of saliva substitutes on mineral content of demineralised and sound dental enamel. *Support Care Cancer* **9**: 40–47.
- Kirstilä V, Lenander-Lumikari M, Söderling E, Tenovuo J (1996). Effects of oral hygiene products containing lactoperoxidase, lysozyme, and lactoferrin on the composition of whole saliva and on subjective oral symptoms in patients with xerostomia. *Acta Odontol Scand* **54**: 391–397.
- Kirstilä V, Lenander-Lumikari M, Tenovuo J (1994a). Effects of a lactoperoxidase system-containing toothpaste on dental plaque and whole saliva *in vivo*. *Acta Odontol Scand* **52**: 346–353.
- Kirstilä V, Tenovuo J, Ruuskanen O, Nikoskelainen J, Irjala K, Vilja P (1994b). Salivary defence factors and oral health in patients with common variable immunodeficiency. *J Clin Immunol* **14**: 229–236.
- Koch G, Edlund K, Hoogendoorn H (1973). Lactoperoxidase in the prevention of plaque accumulation, gingivitis and dental caries. *Odont Revy* **24**: 367–372.
- Koch G, Strand G (1979). Effect of an enzyme dentifrice on caries. A two-year clinical pilot study. *Swed Dent J* **3**: 9–13.
- Kramer A, Böhland H (1996). Biological, medical and chemical aspects of thiocyanate research. *Hygiene Med* **21**: 335–345.
- Lagerlöf F, Oliveby A (1994). Caries-protective factors in saliva. *Adv Dent Res* **8**: 229–238.
- Lassiter MO, Newsome AL, Sams LD, Arnold RR (1987). Characterization of lactoferrin interaction with *Streptococcus mutans*. *J Dent Res* **66**: 480–485.
- Lenander-Lumikari M, Loimaranta V (2000). Saliva and dental caries. *Adv Dent Res* **14**: 40–47.
- Lenander-Lumikari M, Loimaranta V, Hannuksela S, Tenovuo J, Ekstrand J (1997). Combined inhibitory effect of fluoride and hypothiocyanite on the viability and glucose metabolism of *Streptococcus mutans*, serotype c. *Oral Microbiol Immunol* **12**: 231–235.
- Lenander-Lumikari M, Månsson-Rahemtulla B, Rahemtulla F (1992). Lysozyme enhances the inhibitory effects of the peroxidase system on glucose metabolism of *Streptococcus mutans*. *J Dent Res* **71**: 484–490.
- Lenander-Lumikari M, Mikola H, Tenovuo J (1993). Effects of a lactoperoxidase system-containing toothpaste on levels of hypothiocyanite and bacteria in saliva. *Caries Res* **27**: 285–291.
- Loimaranta V, Tenovuo J, Korhonen H (1998). Combined inhibitory effect of bovine immune whey and peroxidase-generated hypothiocyanite against glucose uptake by *Streptococcus mutans*. *Oral Microbiol Immunol* **13**: 378–381.
- Månsson-Rahemtulla B, Pruitt KM, Tenovuo J, Le TM (1983). A mouth-rinse which optimizes *in vivo* generation of hypothiocyanite. *J Dent Res* **62**: 1062–1066.
- Månsson-Rahemtulla B, Rahemtulla F, Baldone DC, Pruitt KM, Hjerpe A (1988). Purification and characterization of human salivary peroxidase. *Biochemistry* **27**: 233–239.
- Månsson-Rahemtulla B, Rahemtulla F, Humphreys-Beher MG (1990). Human salivary peroxidase and bovine lactoperoxidase are cross-reactive. *J Dent Res* **69**: 1839–1846.
- Martinez-Gomis J, Fernandez-Solanas A, Vinas M, Gonzalez P, Planas ME, Sanchez S (1999). Effects of topical application of free and liposome-encapsulated lactoferrin and lactoperoxidase on oral microbiota and dental caries in rats. *Archs Oral Biol* **44**: 901–906.
- Midda M, Cooksey MW (1986). Clinical uses of an enzyme-containing dentifrice. *J Clin Periodontol* **13**: 950–956.
- Moran J, Addy M (1984). The antibacterial properties of some commercially available toothpastes *in vitro*. *Br Dent J* **156**: 175–178.
- Mulligan R, Navazesh M, Slots J (1992). Antibacterial activity of an alcohol-free mouthwash with enzymes. *J Dent Res* **71**: 156 (Abstract).
- Närhi T (1994). Prevalence of subjective feelings of dry mouth in the elderly. *J Dent Res* **73**: 20–25.
- Nederfors T (2000). Xerostomia and hyposalivation. *Adv Dent Res* **14**: 48–56.
- Norhagen Engström G, Engström P-E, Hammarström L, Söder P-Ö, Smith CIE (1989). Immunoglobulin levels in saliva in individuals with selective IgA deficiency: compensatory IgM secretion and its correlation with HLA and susceptibility to infections. *J Clin Immunol* **9**: 279–286.
- Pollock JJ, Katona LI, Goodman H, Cho M-I, Iacono VJ (1981). Bacteriolysis of *Streptococcus mutans* BHT by lysozyme and inorganic anions normally present in human saliva. *Archs Oral Biol* **26**: 711–716.
- Pruitt KM, Reiter B (1985). Biochemistry of peroxidase system: antimicrobial effects. In: Pruitt KM, Tenovuo J, eds. *The Lactoperoxidase system, chemistry and biological significance*. Marcel Dekker Inc.: New York, pp 143–178.
- Rhodus NL, Brown J (1990). The association of xerostomia and inadequate intake in older adults. *J Am Diet Assoc* **90**: 1688–1692.
- Rosin M, Kocher T, Kramer A (2001). Effects of SCN<sup>-</sup>/H<sub>2</sub>O<sub>2</sub> combinations in dentifrices on plaque and gingivitis. *J Clin Periodontol* **28**: 270–276.
- Rotgans J, Hoogendoorn H (1979). The effect of brushing with a toothpaste containing amyloglucosidase and glucose oxidase on dental caries in rats. *Caries Res* **13**: 150–153.
- Schoenfeld S, Stamm JW, Meskin LH, Silverstone LM (1983). The effect of an enzyme-containing dentifrice on plaque and gingivitis. *J Dent Res* **62**: 178 (Abstract).
- Soukka T, Lumikari M, Tenovuo J (1991a). Combined bactericidal effect of human lactoferrin and lysozyme against *Streptococcus mutans* serotype c. *Microb Ecol Health Dis* **4**: 259–264.
- Soukka T, Lumikari M, Tenovuo J (1991b). Combined inhibitory effect of lactoferrin and lactoperoxidase system on the viability of *Streptococcus mutans*, serotype c. *Scand J Dent Res* **99**: 390–396.
- van Steenberghe D, Van den Eynde E, Jacobs R, Quirynen M (1994). Effect of a lactoperoxidase containing toothpaste in radiation-induced xerostomia. *Int Dent J* **44**: 133–138.
- Tenovuo J (1989). Nonimmunoglobulin defence factors in human saliva. In: Tenovuo J, ed. *Human saliva: clinical chemistry and microbiology*, Vol. II. CRC Press, Inc.: Boca Raton, FL, pp 55–91.
- Tenovuo J (1998). Antimicrobial function of human saliva – how important is it for oral health? *Acta Odontol Scand* **56**: 250–256.
- Tenovuo J, Larjava H (1984). The protective effect of peroxidase and thiocyanate against hydrogen peroxide toxicity assessed by the uptake of <sup>3</sup>H-thymidine by human gingival fibroblasts cultured *in vitro*. *Archs Oral Biol* **29**: 445–451.
- Tenovuo J, Lumikari M, Soukka T (1991). Salivary lysozyme, lactoferrin and peroxidases: antibacterial effects of cariogenic bacteria and clinical applications in preventive dentistry. *Proc Finn Dent Soc* **87**: 197–208.
- Tenovuo J, Moldoveanu Z, Mestecky J, Pruitt KM, Månsson-Rahemtulla B (1982a). Interaction of specific and innate factors of immunity: IgA enhances the antimicrobial effect of the lactoperoxidase system against *Streptococcus mutans*. *J Immunol* **128**: 726–731.
- Tenovuo J, Pruitt KM, Thomas EL (1982b). Peroxidase antimicrobial system of human saliva: hypothiocyanite

- levels in resting and stimulated saliva. *J Dent Res* **61**: 982–985.
- Toljanic JA, Siddiqui AA, Pattersson GL, Irwin ME (1996). An evaluation of a dentifrice containing salivary peroxidase elements for the control of gingival disease in patients with irradiated head and neck cancer. *J Prosthet Dent* **76**: 292–296.
- Warde P, Kroll B, O’Sullivan *et al* (2000). A phase II study of Biotene in the treatment of postradiation xerostomia in patients with head and neck cancer. *Support Care Cancer* **8**: 203–208.
- Wei H, Loimaranta V, Tenovuo *et al* (2002). Stability and activity of specific antibodies against *Streptococcus mutans* and ***Streptococcus sobrinus*** in bovine milk fermented with ***Lactobacillus rhamnosus*** strain GG or treated at ultra-high temperature. *Oral Microbiol Immunol*, in press.