

a
Lineage
Primary disease
Subtype
14
7
3
11
20
27
7
37
64
0
30
60
90
Cancer types
enough lines
both groups, too few lines
one Y group missing
b
Lineage
Primary disease
Subtype
Y-positive lines
Y-negative lines
Lymphoid
CNS
Skin
Lung
PNS
Bone
H&N
Soft tissue
Myeloid
Kidney
Pleura
Liver
Bowel
Bladder/UT
Eso/stomach
Biliary
Pancreas
Prostate
ATDG
MEL
NB
B-cell
HNSCC
AML
ES
MESO
LNET
NSCLC
BLCA
RCC
CRC
HCC
EGA
PAAD
ESCC
GBM
SCLC
OCSCC
AST-IDHmut
COAD
LUAD
LUSC
Cancer types 1 3
c
Lung (44 / 28)
Head and Neck (23 / 19)
CNS/Brain (12 / 43)
Lymphoid (10 / 43)
Bowel (18 / 11)
Esophagus/Stomach (36 / 7)
Skin (7 / 31)
Kidney (9 / 14)
Liver (9 / 11)
Bladder/Urinary Tract (8 / 10)
Bone (5 / 27)
Soft Tissue (6 / 14)
Myeloid (5 / 20)
Biliary Tract (6 / 7)
Lineage
Non-Small Cell Lung Cancer (36 / 15)
Head and Neck Squamous Cell Carcinoma (23 / 18)
Adult-Type Diffuse Glioma (9 / 33)
Colorectal Adenocarcinoma (18 / 11)
Mature B-Cell Neoplasms (8 / 23)
Esophagogastric Adenocarcinoma (23 / 6)
Renal Cell Carcinoma (8 / 11)
Lung Neuroendocrine Tumor (7 / 12)
Melanoma (5 / 29)
Hepatocellular Carcinoma (8 / 8)
Bladder Urothelial Carcinoma (7 / 9)
Primary disease
Oral Cavity Squamous Cell Carcinoma (17 / 12)
Colon Adenocarcinoma (12 / 11)
Lung Adenocarcinoma (16 / 8)
Glioblastoma, IDH-Wildtype (7 / 20)
Small Cell Lung Cancer (7 / 12)
Subtype
0.1
0.15
0.2
0.25
0.3
0.4
0.5
0.75
1
1.5
2
any size
Shift in LOY lines (Chronos units)
Probability of detection
0.0
0.5
1.0
d
Δ = +0.670
P = 5.45e-06
FDR = 0.161
Δ = +0.025
P = 0.788
INSM1 gene effect (Chronos)
Y+
n = 7
LOY
n = 13
Y+
n = 21
LOY
n = 31
MYC-altered
MYC-unaltered
Lung cancer
e
Δ = +0.531
P = 8.03e-06
FDR = 0.196
ATRX gene effect (Chronos)
No models
Y+
n = 20
LOY
n = 5
Y+
n = 9
LOY
n = 0
CDKN2A-altered
CDKN2A-unaltered
Melanoma