

a
PC1 ~ passage: R² = 0.93
PC2 ~ condition: R² = 0.66
PC2 (18.4%)
PC1 (42.3%)
sgCTRL
Y-depleted
P12
P13
P14
b
FDR < 0.05
FDR < 0.05, CN-concordant
FDR <0.05, |log2FC| ≥0.5, consistent direction
147 at FDR < 0.05
114 meet additional thresholds (88 copy-number stable)
−log10 P
log2FC (Y-depleted / sgCTRL)
TACSTD2
AP1M2
THBD
CLDN7
GRHL2
SCEL
EBF4
TAF9
F13A1
COL4A2
CREB3L1
SLC16A2
ST14
SPOCK2
c
Higher in some passages, lower in others
RSPO3
ADAM12
TAF9
IFITM1
S100A14
Higher in all 3 passages
SFTPA1
LINC02967
PXDN
CCR9
GRHL2
Each dot: one gene
Passages: P12–P14
Lower in all 3 passages
Highest change across passages (log2 CPM, Y-depleted − sgCTRL)
Lowest change across passages (log2 CPM, Y-depleted − sgCTRL)
Stable
Discordant
Concordant
d
Pearson r
gene 0.10
10 Mb 0.69
arm 0.89
chr10p
chrXp
chrXq
chr15q
chr20p
RNA arm mean log2 CPM difference
lpWGS arm median log2 difference
e
Genes
39
107
25
88
1
1
FDR < 0.05 (147)
|log2FC| ≥0.5 + consistency (114)
f
FDR = 7.3e-04
log2FC = -3.95
GRHL2 (CPM)
sgCTRL
Y-depleted
g
FDR = 3.9e-04
log2FC = 1.32
TAF9 (log2 CPM)
sgCTRL
Y-depleted
h
FDR = 1.8e-02
log2FC = -2.03
PXDN (log2 CPM)
sgCTRL
Y-depleted