

a
ATAC (each passage)
RNA
D0 lpWGS
Y-depleted − sgCTRL (log2, chromosome median)
Lost
Gained
33 of 95 gained
% regions FDR < 0.05
Chromosome
b
ρ = 0.60; P = 5.3e-05
RNA difference (log2)
lpWGS difference (log2)
18p
18q
13q
ρ = -0.05; P = 0.774
ATAC difference (log2)
39 autosomal arms; ρ, Spearman; r, Pearson
ρ = 0.30; P = 0.067
r = 0.42; P = 0.008
20p
RNA, combined difference (log2)
ρ = 0.29; P = 0.074
r = 0.22; P = 0.188
RNA, August difference (log2)
ρ = 0.26; P = 0.110
r = 0.45; P = 0.004
RNA, May difference (log2)
c
ATAC, 5 Mb median
lpWGS Δ
Region
Gained / lost
chr18 (gained 33; lost 4)
log2 (Y-depleted / sgCTRL)
chr13 (gained 1; lost 36)
Position (Mb, hg38)
d
All targets
Chromosome exclusion
TFAP2
ZIC
TEAD
GATA
RUNX
EBF
SOX6
Mantel–Haenszel OR vs matched controls (95% CI)
e
Lower-expression DE genes
58/228
40/150
Non-DE genes
101/12,983
68/9,111
All promoters
lpWGS |Δ| < 0.2
% with a lost promoter peak (ATAC FDR < 0.10)