

a Lower expression in Y-depleted HT1080

D01 Interferon-response genes




n = 11; annotation q = 1.3e-08

D02 Integrin–collagen–matrix module




n = 15; annotation q = 1.1e-20

D03 Ankyrin–spectrin scaffold




n = 6; annotation q = 1.5e-12

D04 Chemokines, growth factors and contractile genes




n = 17; annotation q = 4.7e-07

D08 AKR1C enzymes and DHDH




n = 4; annotation q = 5.9e-06

D10 Cadherin–junction module




n = 5; annotation q = 2.3e-07

D11 PI3K–PLC signalling




n = 5; annotation q = 4.8e-05

D15 UNC5–FLRT–latrophilin module



n = 4; annotation q = 5.9e-03


D18 Innate-response signalling



n = 3; annotation q = 1.2e-03


b Higher expression in Y-depleted HT1080

U07 Eicosanoid metabolism, glutathione enzymes and cytokines




n = 11; annotation q = 3.0e-05

U03 Chemokine receptors and immune co-signalling




n = 7; annotation q = 5.7e-07

U16 TREM receptors and CKLF/CMTM family




n = 6; annotation q = 6.3e-05

U05 Hedgehog-associated module




n = 4; annotation q = 2.4e-06

U06 Eph–ephrin module




n = 4; annotation q = 6.3e-07

U08 WNT ligands / modulators




n = 4; annotation q = 1.9e-04

U09 Synaptic adhesion-associated genes




n = 4; annotation q = 6.5e-07

U11 BMP ligands / antagonist



n = 3; annotation q = 7.5e-04

U18 Netrin–guidance module



n = 4; annotation q = 7.4e-05

Physical support ≥ 0.7 — Other STRING association

Node area scales with |mean $\log_2$ RNA difference|