
a
HT1080
A549
DU145
A549 LOY clones
A549 clones, multiome
MB49 cells (mouse)
MB49 tumours (mouse)
HT1080 spheroids
A549 (Tahoe)
J82 (Tahoe)
SK-MEL-2 (Tahoe)
HCT15 (Tahoe, 6 wells)
Lymphocytes
iPSC trophoblast
iPSC neural crest
Fetal tissues
Fibroblasts
Lymphoblastoid lines
hESC UTY-KO
hESC UTX/UTY-DKO
XYY vs XY (flipped)
-0.93
-1.03
-2.01
-6.24 (FDR 0.014), clipped
-6.21 (FDR 0.017), clipped
+0.56 (FDR 0.19)
+0.28 (FDR 0.51)
not tested
-0.72 (FDR 0.048)
not tested
not tested
not tested
+0.42 (FDR 0.79)
+0.54 (FDR 0.18)
-0.11 (FDR 0.33)
-0.35 (FDR 0.11)
-0.06 (FDR 0.92)
+1.38 (FDR 0.67)
-0.06 (FDR 0.8)
+0.07 (FDR 0.59)
-0.02
Thesis lines
Somatic LOY
Natural Y-mosaic, single-cell
45,X vs 46,XY
Not LOY
−4
−3
−2
−1
0
1
2
PXDN effect, Y-loss polarity (log2; clipped at ±3)
b
DDX3Y -8.40
DDX3Y -8.37
DDX3Y -4.29
Y genes absent; contrast label
TXLNGY -14.31
Y genes absent; contrast label
Y genes absent; contrast label
DDX3Y -8.80
DDX3Y -7.59
DDX3Y -6.70
DDX3Y -7.34
DDX3Y -7.87
Y genes absent; contrast label
Y genes absent; contrast label
Y genes absent; contrast label
Y genes absent; contrast label
DDX3Y -8.68
DDX3Y -10.26
Y genes absent; contrast label
Y genes absent; contrast label
DDX3Y -1.10