

a
chrX, 3 windows
chr1 150–160 Mb
chr11 110–140 Mb
chr20 0–30 Mb
r = 0.69, slope 0.62
296 windows
RNA 10-Mb mean log2 CPM difference
lpWGS 10-Mb median log2 difference
b
r = 0.10
17,443 genes
4 points outside axes
RNA mean log2 CPM difference
lpWGS gene-level log2 difference
c
chr20p
chr11q
chr10q
chr1q
chr9q
chr19q
chr6p
chr11p
20p deletion
Genes at FDR < 0.05
d
chr20q
chr15q
chrXq
chr18p
chrXp
chr20p
DMSO-LOY-1X minus sgCTRL
LOY-2X minus sgCTRL (arm median log2)
e
Effect sizes agree closely (Pearson r = 0.954);
no gene reaches FDR < 0.05 without passage blocking
Paired FDR < 0.05
FDR <0.05, |log2FC| ≥0.5,
consistent direction
Unpaired model log2FC
Paired model log2FC (~ passage + condition)
f
Passage: R² = 0.93
PC1 score
P12
P13
P14
g
Y-depleted ÷
sgCTRL = 0.28
MSY reads
(% assigned)
sgCTRL P12
sgCTRL P13
sgCTRL P14
Y-dep. P12
Y-dep. P13
Y-dep. P14
h
C13
Y13
Y12
C12
Y14
C14
r
C, sgCTRL; Y, Y-depleted