

a
Genes per quadrant at |log2FC| > 0.25 (> 0.5); 13,300 genes
DOWN-OPEN
208 (7)
UP-OPEN
391 (60)
DOWN-CLOSED
573 (269)
UP-CLOSED
170 (45)
Promoter ATAC log2FC (median of tested peaks)
RNA log2FC (Y-depleted / sgCTRL)
b
DOWN-CLOSED (573 genes; top 4 sets per collection)
inflammatory response
epithelial mesenchymal transition
kras signaling up
kras signaling dn
g protein coupled receptor signaling pathway
synapse organization
regulation of membrane potential
action potential
benporath es with h3k27me3
benporath suz12 targets
nikolsky breast cancer 19q13.1 amplicon
servitja islet hnf1a targets up
Hallmark
GO:BP
CGP
Fold enrichment vs all genes
−log10 BH FDR
Genes
FDR < 0.05 vs RNA-direction background
c
ES H3K27me3 targets (% of genes)
14.1% 24/170
5.3% 84/1,576
4.3% 17/391
10.8% 40/371
2.4% 176/7,403
1.6% 19/1,159
10.8% 62/573
3.8% 55/1,449
5.8% 12/208
RNA (|log2FC| > 0.25)
Promoter ATAC (|log2FC| > 0.25)
UP-OPEN (391 genes; top 4 sets per collection)
response to xenobiotic stimulus
organic hydroxy compound transport
cellular response to xenobiotic stimulus
sterol transport
liu prostate cancer dn
ohguchi liver hnf4a targets dn
meissner npc hcp with h3k4me2
mebarki hcc progenitor fzd8crd dn
d
ES H3K27me3 targets
PRC2 targets
BH FDR ≥ 0.05
K = 1 matching
Closed genes, unmatched
Closed genes (> 0.25), matched
RNA-flat genes only
Closed genes (> 0.5), matched
Open genes (> 0.25), matched
Lost peaks, FDR < 0.05
Lost peaks, FDR < 0.10
Gained peaks, FDR < 0.10
n = 1,114
n = 644
n = 236
n = 326
n = 1,597
n = 128
n = 247
n = 67
Odds ratio (Mantel–Haenszel, 95% CI)
Matched on baseline logCPM × GC × CpG o/e (K = 3 controls; bars show K = 1)