Matched CPM + 1 contrasts at D0 and endpoint

Guide $\log_2$ fold change (LOY / sgCTRL)

Chr 1 (n = 7,915) | Chr 2 (n = 4,799) | Chr 3 (n = 4,150) | Chr 4 (n = 2,918)

Chr 5 (n = 3,362) | Chr 6 (n = 4,002) | Chr 7 (n = 3,511) | Chr 8 (n = 2,578)

Chr 9 (n = 3,007) | Chr 10 (n = 2,845) | Chr 11 (n = 4,999) | Chr 12 (n = 3,962)

Chr 13 (n = 1,234) | Chr 14 (n = 2,328) | Chr 15 (n = 2,238) | Chr 16 (n = 3,183)

Chr 17 (n = 4,526) | Chr 18 (n = 1,051) | Chr 19 (n = 5,447) | Chr 20 (n = 2,094)

Chr 21 (n = 847) | Chr 22 (n = 1,682) | Chr X (n = 3,224) | Chr Y (n = 186)




Grey: all observed guide pairs (2.5% opacity). Black: stage-wise medians. Density reflects guide count; no downsampling.

Both stages: $\log_2[(\text{LOY CPM} + 1)/(\text{sgCTRL CPM} + 1)]$. Endpoint CPM is the within-genotype mean of R1/R2.

All panels share limits and retain all finite values. Guides absent from both genotypes at either stage are omitted.

Chromosome-level summaries using matched CPM + 1 contrasts




ΔLFC = endpoint − D0 for each guide. Panel b subtracts the median NTC ΔLFC within each guide set.

Grey intervals: interquartile ranges of guide shifts, not confidence intervals. Diamonds: D0 read-threshold sensitivity.

NTC median ΔLFC: -0.108 (n = 998); ≥50 subset: -0.107.

Descriptive chromosome associations; no significance tests. Guide counts are not independent biological replicates.

Original display: D0 CPM + 1 contrast to saved MAGeCK endpoint LFC



Grey: all observed guide pairs (2.5% opacity). Black: stage-wise medians. Density reflects guide count; no downsampling.

The two estimators use different normalisation. This companion view is not used to quantify chromosome shifts.

All panels share limits and retain all finite values. Guides absent from both genotypes at either stage are omitted.