
a
b
Oncogene addiction
Paralog or synthetic lethal
Collateral
Lost dependency
Y paralog
Observed
Lung (44 / 28)
Head and Neck (23 / 19)
CNS/Brain (12 / 43)
Lymphoid (10 / 43)
Bowel (18 / 11)
Esophagus/Stomach (36 / 7)
Skin (7 / 31)
Kidney (9 / 14)
Liver (9 / 11)
Bladder/Urinary Tract (8 / 10)
Bone (5 / 27)
Soft Tissue (6 / 14)
Myeloid (5 / 20)
Biliary Tract (6 / 7)
Lineage
Non-Small Cell Lung Cancer (36 / 15)
Head and Neck Squamous Cell Carcinoma (23 / 18)
Adult-Type Diffuse Glioma (9 / 33)
Colorectal Adenocarcinoma (18 / 11)
Mature B-Cell Neoplasms (8 / 23)
Esophagogastric Adenocarcinoma (23 / 6)
Renal Cell Carcinoma (8 / 11)
Lung Neuroendocrine Tumor (7 / 12)
Melanoma (5 / 29)
Hepatocellular Carcinoma (8 / 8)
Bladder Urothelial Carcinoma (7 / 9)
Primary disease
Oral Cavity Squamous Cell Carcinoma (17 / 12)
Colon Adenocarcinoma (12 / 11)
Lung Adenocarcinoma (16 / 8)
Glioblastoma, IDH-Wildtype (7 / 20)
Small Cell Lung Cancer (7 / 12)
Subtype
NRAS in NRAS Q61 (-1.35)
KRAS in KRAS-driver (-1.18)
BRAF in BRAF V600E (-0.94)
CTNNB1 in APC loss (-0.88)
CDK4 in CDK4-amplified (-0.77)
ERBB2 in ERBB2-amplified (-0.75)
MDM2 in MDM2-amplified (-0.49)
PIK3CA in PIK3CA-driver (-0.48)
NFE2L2 in NFE2L2-driver (-0.40)
MYCN in MYCN-amplified (-0.33)
STAG1 in STAG2-mutant (-0.72)
WRN in MSI (-0.54)
CREBBP in EP300-mutant (-0.38)
SMARCA2 in SMARCA4-mutant (-0.28)
ARID1B in ARID1A-mutant (-0.17)
PIK3CB in PTEN loss (-0.15)
PRMT5 in MTAP-deleted (-0.28)
MAT2A in MTAP-deleted (-0.13)
MDM2 in TP53-altered (+0.78)
CDK6 in RB1 loss (+0.46)
CDK4 in RB1 loss (+0.37)
EIF1AX in Y-negative (-1.23)
DDX3X in Y-negative (-0.75)
RPS4X in Y-negative (-0.60)
ZFX in Y-negative (-0.27)
EIF1AX
DDX3X
RPS4X
ZFX
Probability of detection
0.0
0.5
1.0
Observed
p of 0.05 or more
p below 0.05 only
genome-wide, this cancer type alone
genome-wide, all types of its level