

DETSI Wide Bay Report

EP Consulting and Dumelle Consulting

Table of contents

1	Executive Summary	1
2	Background	2
3	The Data	6
4	Water Quality and Geographic Modeling	7
4.1	Model Fit	10
4.2	Block Kriging	13
4.3	Alternative Approaches	14
5	Resilience Modeling	19
5.1	Model Fit	21
5.2	Alternative Approaches	22
6	Limitations and Future Suggestions	23
7	Revisiting the Research Questions	24
8	Discussion	25
	References	28

1 Executive Summary

Seagrass meadows are a critical component of the Wide Bay ecosystem, providing habitat structure for a wide array of important marine species like dugongs and green turtles. This report presents a spatially explicit statistical analysis of total seagrass cover in Wide Bay across three financial years (2021–2022, 2022–2023, and 2023–2024), a period following two

successive flood events that impacted seagrass ecosystems. Using water quality data from the eReefs monitoring platform alongside field observations and geographic data, we address three management-relevant research questions: what water quality conditions trigger concern for seagrass cover, whether and where cover has recovered since the flood, and what site-level characteristics make seagrass meadows more or less resilient to disturbance.

Our results indicate that temperature is a water quality driver of total seagrass cover in Wide Bay, with a one-degree increase in 90-day mean temperature associated with a 0.5 percentage point decrease in cover. Turbidity and flood turbidity showed little statistical evidence of association with total cover after accounting for other variables, likely reflecting the absence of pre-flood baseline data against which post-flood change can be assessed. Bay-wide, block kriging predictions show no evidence of declining total cover since the floods, with some zone classes and tidal ranges showing signs of modest improvement. At the site level, species richness and dominant species identity were the strongest predictors of resilience, with higher species richness and sites dominated by HS tending to be the most resilient. Distance to coral habitat also emerged as an important, nonlinear predictor of resilience, whereby sites far away from coral habitat were most resilient. Together, these findings can help inform where seagrass conservation investment in Wide Bay is likely to be most effective.

2 Background

Accurately characterizing drivers and resilience of seagrass cover is essential to improving the ecological health of the Wide Bay ecosystem. Over the past few decades, various scientific efforts have informed our current understanding of trends, behaviors, and flood responses of seagrass cover in Wide Bay [Preen et al., 1995, McMahon et al., 2005, York et al., 2022] as well as Australia more widely [Kilminster et al., 2015, Gilby et al., 2018, O'Brien et al., 2018]. Remote sensing has offered new capabilities for seagrass cover mapping [Roelfsema and Phinn, 2009], while innovative statistical approaches have offered novel methods for studying resilience at site-specific scales [Gilby et al., 2023]. Our work contributes to this body of literature by applying spatially explicit statistical models [Cressie, 1993] to study seagrass cover drivers, resilience, and recovery in Wide Bay. Specifically, we provide and contextualize answers to four specific research questions, applicable to Wide Bay:

1. Can we identify water quality covariates that act as triggers for seagrass cover management responses? Are these triggers different based on whether they result from a specific event (e.g., flood) or are ambient?
2. Did seagrass cover improve after the flood events? If so, how and where?
3. Can we identify site-specific resilience indicators, which describe where recovery is likely to be enhanced or impeded?
4. What factors contribute to seagrass resilience?

The spatially explicit statistical models (i.e., spatial models) we used to answer these research questions are related to commonly used linear and generalized linear models but incorporate *spatial dependence*. Spatial dependence measures how similar two sites are based on their distance. Formally, we can build spatial dependence directly into spatial models by leveraging Tobler’s First Law of Geography [Tobler, 1970], which states succinctly that “nearby sites tend to be more similar than distant sites”. Not only is this intuitive from a “first principles” perspective, but spatial models also provide many advantages for modeling spatial data compared to models that assume independence among observations (i.e., nonspatial models). First, nonspatial models tend to underestimate uncertainty, yielding covariate standard errors and p-values for covariates that are too small and corresponding confidence intervals that are too narrow. Second, spatial models improve prediction at unsampled locations by borrowing strength from nearby observations, yielding more precise predictions and corresponding standard errors that vary by location. Third, spatial models decompose model error into separate spatial and nonspatial components, informing residual and resilience structure in ways that nonspatial models cannot. Fourth, spatial models can provide ecologically meaningful insight into the spatial dependence structure of the process, informing relationships among measured and unmeasured variables as well as future sampling plans. Zimmerman and Ver Hoef [2024] provide a thorough review of spatial models for ecological and environmental data, providing more detail and justification for the claims made above.

Spatial models offer many benefits over commonly used machine learning algorithms like random forests [Breiman, 2001, Cutler et al., 2007, James et al., 2017, Kuhn and Silge, 2022]. First, traditional machine learning algorithms do not incorporate spatial dependence and, by consequence, often perform worse than spatial models at characterizing important covariates and predicting at unobserved locations [Fox et al., 2020]. Second, machine learning algorithms struggle to quantify uncertainty in a principled manner, something spatial models excel at. Third, covariate inference is more straightforward using spatial models, as they naturally provide p-values that enable structured testing of hypotheses. Fourth, spatial models provide a rich set of diagnostics like leverage and Cook’s distances [Montgomery et al., 2021], quantities not generally defined for machine learning algorithms. This is not to say that machine learning algorithms cannot be very useful in ecological data analyses, but rather than spatial models are sometimes a more effective tool for modeling spatial data.

Though there are many benefits of spatial models over machine learning algorithms, the reverse is also true. Machine learning algorithms much more simply capture complex nonlinearities “out-of-the-box”, bypassing the tuning often required for spatial models that involve nonlinear components like splines. This can be especially useful for ecological data, which are often full of nonlinear relationships [Fox et al., 2017]. Machine learning algorithms also typically rely on few, if any, distributional assumptions and can be more robust against unusual observations like outliers [Hastie et al., 2009]. Machine learning algorithms also tend to be much more computationally efficient than spatial models. Importantly, machine learning algorithms and spatial models can be complementary tools that help shed insight into complicated ecological processes (something we reiterate in the upcoming analyses). An exciting and developing area

of statistical research involves incorporating spatial modeling strategies directly into machine learning algorithms [Saha et al., 2023, Heaton et al., 2025].

The first spatially explicit model we define is the spatial linear model. The spatial linear model is a *mixed-effects model* [Pinheiro and Bates, 2000, Bolker, 2015] written as

$$y_i = \beta_0 + \beta_1 x_{1,i} + \beta_2 x_{2,i} + \dots + \beta_p x_{p,i} + \tau_i + \epsilon_i, \quad (1)$$

where $i = 1, 2, \dots, n$ indexes the n observations. The quantity y_i is the response variable (for observation i), the β parameters are slope parameters controlling the covariates, $x_{j,i}$ (for $j = 1, 2, \dots, p$). The random error τ_i is a spatial random error, and the random error ϵ_i is the nonspatial (i.e., independent) random error. The spatial random error is spatially structured, capturing spatial residual effects like environmental gradients we do not have data to fully represent. The nonspatial random error is not spatially structured and captures nonspatial residual effects like measurement error and microscale variation. The difference between a traditional (i.e., nonspatial) linear model and a spatial linear model is the inclusion of τ_i (in the spatial linear model). The spatial random error implied by τ_i is governed by a *spatial covariance function*. The spatial covariance function allows the model to incorporate spatial dependence, yielding the improved model performance previously described. Figure 1 shows three different spatial covariance functions that describe different ways the spatial covariance decays with distance. Model parameters and standard errors are estimated from data using likelihood-based [Patterson and Thompson, 1971, Harville, 1977, Wolfinger et al., 1994] or semivariogram-based [Cressie, 1985, Curriero and Lele, 1999] estimation methods. Often, the spatial linear model in Equation 1 is written more compactly using *matrix notation*:

$$\mathbf{y} = \mathbf{X}\beta + \tau + \epsilon, \quad (2)$$

where $\mathbf{X}$ is a matrix that holds the j covariates for each of n observations, and $\mathbf{y}$, β , τ , and ϵ are vectors that contain each of the n or j quantities from Equation 1. The spatial linear model can be adjusted to account for nonspatial random effects just as one would add random effects to a nonspatial linear mixed model:

$$\mathbf{y} = \mathbf{X}\beta + \mathbf{Z}\mathbf{u} + \tau + \epsilon, \quad (3)$$

where $\mathbf{Z}$ is a random effect design matrix (dimension $n \times k$) that indexes each of the k random effects in $\mathbf{u}$. For more information about specific forms of various spatial covariance functions, see Schabenberger and Gotway [2017] and Zimmerman and Ver Hoef [2024].

Like the traditional linear model, the spatial linear model can capture complex nonlinearities in the data through the use of interaction effects [Faraway, 2002, Lenth and Piaskowski, 2025], splines [Chambers and Hastie, 1992, Ruppert et al., 2003], additive models [Wood, 2017, Pedersen et al., 2019], and penalized regression models like the ridge and lasso [Tibshirani, 1996]. Generally, this is accomplished by leveraging *basis functions* that break down nonlinear behavior into a set of additive linear components that may include penalty terms which guard against overfitting. Furthermore, there is no assumption that y_i itself is normally distributed; rather,




Figure 1: Three spatial covariance functions that represent different spatial dependence behavior.

distributional assumptions are placed on the random errors after incorporating the slope parameters. This clarification implies the spatial (and nonspatial) linear models can be quite effective at modeling a range of data that are not normally distributed. For highly nonnormal data, tools like generalized linear models, which we discuss later, can be helpful. Finally, under general conditions, the slope parameters of the spatial (and nonspatial) linear models are approximately normally distributed with a large sample size (see [Casella and Berger \[2024\]](#) for a general justification via the Central Limit Theorem and [Zhang and Zimmerman \[2005\]](#) for nuances regarding spatial data). This is beneficial because it justifies valid hypothesis testing on the slope coefficients (often of primary ecological interest), no matter the distribution of the data (given certain assumptions are satisfied).

One of the most useful applications of the spatial linear model is predicting the response variable at unobserved locations. A very useful spatial prediction method called Kriging has a rich theoretical justification [[Cressie, 1990](#), [Stein, 1999](#)]. Kriging is the spatial analogue to best linear unbiased prediction (BLUP) for mixed models [[Henderson, 1975](#)]. Via Kriging, the spatial linear model produces BLUPs at each unobserved location of interest, alongside prediction intervals to characterize uncertainty. While useful, sometimes the goal is to create a BLUP for an entire region of interest. This technique, which builds upon the theoretical foundations of Kriging, is called block Kriging [[Ver Hoef, 2002](#), [Cressie, 2006](#)]. The intuition is to make point predictions at a very dense grid in the study region and then average these predictions, appropriately characterizing uncertainty. Together, Kriging and block Kriging provide a complete set of tools for spatial prediction, given the fit of a spatial linear model.

Spatial generalized linear models are an extension of spatial linear models to highly nonnormal

data [Christensen and Waagepetersen, 2002, Zhang, 2002, Diggle and Ribeiro Jr, 2007, Bonat and Ribeiro Jr, 2016]. Ver Hoef et al. [2024] proposed a novel application of the Laplace approximation to enable restricted maximum likelihood estimation of spatial generalized linear models. Their model formulation builds upon Equation 1 and is given by

$$g(\mu_i) = \beta_0 + \beta_1 x_{1,i} + \beta_2 x_{2,i} + \dots + \beta_p x_{p,i} + \tau_i + \epsilon_i, \quad (4)$$

where $g(\mu_i)$ is a “link function” that links the distribution of y_i to a linear function of its mean, μ_i . Common link functions include the log link for count and skewed data and the logit link for binomial and proportion data (Table 1). Using the same argument as for spatial linear models, nonspatial random effects can be added to spatial generalized linear models.

Table 1: Common generalized linear model families, link functions, and appropriate data types.

Family	Link Function	Link Name	Data Type
Binomial	$f(\mu) = \log(\mu/(1 - \mu))$	Logit	Binary; Binary Count
Poisson	$f(\mu) = \log(\mu)$	Log	Count
Negative Binomial	$f(\mu) = \log(\mu)$	Log	Count
Beta	$f(\mu) = \log(\mu/(1 - \mu))$	Logit	Proportion
Gamma	$f(\mu) = \log(\mu)$	Log	Skewed
Inverse Gaussian	$f(\mu) = \log(\mu)$	Log	Skewed

Spatial models are complicated, both theoretically and computationally. This has contributed to their relative lack of use in the ecological community. However, recent advances in open-source software have made spatial models more accessible to ecologists who have strong subject matter expertise but may lack sufficient statistical background or computational skills to implement them from scratch. One such advancement is the **spmodel** **R** package [Dumelle et al., 2023], which emulates base-**R** functions like `lm()` and `glm()` to account for spatial dependence via `splm()` and `spglm()`, respectively. Familiar **R** functions like `summary()` and `predict()` can be used directly on models fit via **spmodel**. Importantly, **spmodel** also has tools for spatial random forest residual modeling [Fox et al., 2020], applications to large data of several thousand observations via spatial indexing [Ver Hoef et al., 2023], and cross validation [Stone, 1974, Roberts et al., 2017].

3 The Data

We started with nearly 4,000 seagrass cover observations in Wide Bay across three financial years (2021-2022, 2022-2023, and 2023-2024). Some data (approximately 1,000 observations) were available from 1998-2002 but were omitted as they were far away temporally and did not have corresponding eReefs data [Steven et al., 2019]. In total, eReefs data with modeled

water quality covariates related to turbidity and salinity were available at 2,663 observations. Because one of our research questions involved connecting water quality covariates to management actions, we proceed with this subset of 2,663 observations for modeling. Of the 2,663 observations, 1,073 were in 2021-2022, 524 were in 2022-2023, and 1,066 were in 2023-2024 (Table 2).

Table 2: Total seagrass observations in Wide Bay by financial year.

Financial Year	Sample Size	Percent (of Total)
2021-2022	1,073	40.3%
2022-2023	524	19.7%
2023-2024	1,066	40.0%
Total	2,663	100.0%

4 Water Quality and Geographic Modeling

Total cover is a percentage that ranges from 0% (no seagrass) to 68.3%. Figure 2 shows total cover by financial year throughout the Bay. Total cover tends to be largest along the eastern shoreline of the Bay and lowest in the middle of the Bay. Figure 2 also reveals that 2022-2023 lacked sampling in much of the Great Sandy Strait.

We considered for modeling several water quality and geographic covariates. Table 3 summarizes the water quality variables, which included both ambient and flood-related metrics related to turbidity, salinity, and temperature. The “Variable” column indicates the water quality variable turbidity, salinity, or temperature. The “Flood” variable indicates whether the measurement corresponds to a flood variable or not. The “Period” variable indicates the prior period (in days) relative to sampling or indicates whether the variable corresponds to a flood event. The “Type” column is the type of statistic, either a mean, minimum, maximum, number of days above (or below) a cutoff during “Period”, or a maximum spell (i.e., duration) above (or below) a cutoff during “Period”. The “Included” variable indicates whether the variable was included as a covariate in the final spatial model. We included only a few variables in the final model: mean turbidity in the 90 days prior to sampling, average mean turbidity during both flood events, and mean temperature in the 90 days prior to sampling. We only included a single turbidity metric prior to sampling because these variables were all quite correlated with one another, making it hard for the model to tease apart signals. We believed 90 day mean turbidity prior to sampling to be the most accurate “running average” of turbidity prior to sampling. We used a similar rationale to select only a single turbidity metric from the flood and a single temperature metric prior to sampling. We omitted a flood salinity metric primarily because we lacked data for salinity prior to sampling, thus it would be challenging to establish a “baseline” for salinity in order to tease apart prior to sampling




Figure 2: Total seagrass cover in Wide Bay by financial year.

effects from flood effects (moreover, they did not seem strongly related to total cover). We provide more detail regarding the statistical modeling in Section 4.1.

Table 3: Wide Bay water quality variables considered for modeling.

Variable	Flood	Period	Type	Cutoff	Included
Turbidity (NTU)	No	7	Mean	NA	No
Turbidity (NTU)	No	30	Mean	NA	No
Turbidity (NTU)	No	90	Mean	NA	Yes
Turbidity (NTU)	No	7	Above	3	No
Turbidity (NTU)	No	30	Above	3	No
Turbidity (NTU)	No	90	Above	3	No
Turbidity (NTU)	No	7	Above	5	No
Turbidity (NTU)	No	30	Above	5	No
Turbidity (NTU)	No	90	Above	5	No
Turbidity (NTU)	No	7	Above	6	No
Turbidity (NTU)	No	30	Above	6	No
Turbidity (NTU)	No	90	Above	6	No
Turbidity (NTU)	Yes	Flood	Above	3	No
Turbidity (NTU)	Yes	Flood	Above	5	No
Turbidity (NTU)	Yes	Flood	Above	6	No
Turbidity (NTU)	Yes	Flood	Duration	3	No
Turbidity (NTU)	Yes	Flood	Duration	5	No
Turbidity (NTU)	Yes	Flood	Duration	6	No
Turbidity (NTU)	Yes	Flood	Mean	NA	Yes
Turbidity (NTU)	Yes	Flood	Max	NA	No
Salinity (PSU)	No	7	Below	20	No
Salinity (PSU)	No	30	Below	20	No
Salinity (PSU)	No	90	Below	20	No
Salinity (PSU)	No	7	Below	25	No
Salinity (PSU)	No	30	Below	25	No
Salinity (PSU)	No	90	Below	25	No
Salinity (PSU)	No	7	Below	30	No
Salinity (PSU)	No	30	Below	30	No
Salinity (PSU)	No	90	Below	30	No
Salinity (PSU)	Yes	Flood	Below	20	No
Salinity (PSU)	Yes	Flood	Below	25	No
Salinity (PSU)	Yes	Flood	Below	30	No
Salinity (PSU)	Yes	Flood	Duration	20	No
Salinity (PSU)	Yes	Flood	Duration	25	No
Salinity (PSU)	Yes	Flood	Duration	30	No
Salinity (PSU)	Yes	Flood	Mean	NA	No

Table 3: Wide Bay water quality variables considered for modeling.

Variable	Flood	Period	Type	Cutoff	Included
Salinity (PSU)	Yes	Flood	Min	NA	No
Temperature (deg C)	No	7	Mean	NA	No
Temperature (deg C)	No	30	Mean	NA	No
Temperature (deg C)	No	90	Mean	NA	Yes

Table 4 summarizes the geographic variables, which have the following structure:

- The “Sediment Type” variable has five levels: Sand, Sandy Mud, Muddy Sand, Mud, and Rock.
- The “Tidal Range” variable has three levels: Shallow Subtidal, Intertidal, and Deep Subtidal.
- The “Years Since Flood (One)” is numeric from 0.096 (May, 2022) to 1.65 (November, 2023).
- The “Years Since Flood (Two)” is numeric from 0 (Pre-flood) to 1.40 (November, 2023).
- The “Algae Presence” variable has two levels: Algae present and algae not present
- The “Algae Cover” variable is numeric from zero to one.
- The “Growing Season” variable has two levels: Growing (September - December) and Not Growing (January - August).
- The “Zone Class” variable has six levels: Central, East, Great Sandy Strait, Hervey Bay, Southern, and Southwest (Figure 3).

Table 4: Wide Bay geographic variables considered for modeling.

Variable	Effect	Included
Sediment Type	Fixed	Yes
Tidal Range	Fixed	Yes
Years Since Flood One	Fixed	Yes
Years Since Flood Two	Fixed	No
Algae Presence	Fixed	No
Algae Cover	Fixed	No
Growing Season	Fixed	No
Zone Class	Random	Yes

From Table 4, we kept the sediment type, tidal range, and years since flood (one) variables as covariates in the model. We only included time since the first flood, as nearly all the data aside from the observations in May 2022 were after both floods, making it possible to study cumulative flood effects but harder to tease apart effects from each individual flood. The algae variables were removed, as they did not seem related to total cover. We removed growing




Figure 3: Observations and their zone classes in Wide Bay.

season from the model after determining that it likely is not ecologically useful, as a simple month proxy likely does not adequately account for the complexities of the more nuanced aspects of seagrass growth with time. We used zone class as a random effect to account for within-class correlation that may impact the relationships among total cover and the remaining variables. We provide more detail regarding the modeling in Section 4.1.

4.1 Model Fit

Table 5: Variables in the final Wide Bay total cover model, their type (numeric or categorical), and if categorical, the number of unique levels.

Variable	Type	No. Levels
90 Day Temperature	Numeric	NA
90 Day Turbidity	Numeric	NA
Mean Floods Turbidity/Years Since Flood One ²	Numeric	NA
Sediment Type	Categorical	5
Tidal Range	Categorical	3

To describe total seagrass cover in Wide Bay, our final model was a spatial linear model (Equation 1) with the following explanatory variables: tidal range, sediment type, 90 day mean temperature (prior to sampling), 90 day turbidity (prior to sampling), and mean turbidity during the flood periods scaled by time since the first flood squared (Table 5). We scaled mean turbidity during the first flood period so that the effect of turbidity during the flood event was allowed to decay with time since the flood. Intuitively, this means that with time since the flood, the impact of turbidity during the flood event is lessened. We also considered scaling by time since flood (not squared), which produced similar results. Furthermore, we used an exponential spatial covariance function and included a random effect for zone class. We fit the model using restricted maximum likelihood via the `sp1m()` function in `spmodel`.

Table 6 shows the results from an analysis of variance applied to the fitted model. The sediment type and temperature covariates had small p -values (p -value < 0.01) indicating significant evidence of a relationship with total cover, while other covariates had large p -values (p -value > 0.1) indicating little evidence of an association with total cover; here are some relevant takeaways:

- Sediment Type: The estimated (marginal) average total cover is largest in mud (4.3%), followed by sand (3.6%), muddy sand (3.2%), rock (3.0%), and sandy mud (2.8%).
- 90 Day Temperature: A one-degree (C) increase in 90 day temperature is associated with a decrease in average total cover by 0.5% points.

Together, these results can inform triggers for management responses in various conditions. For example, given the decrease in total cover with temperature, one could propose a specific




Figure 4: Average total cover as a function of 90 day mean temperature. Estimates (and confidence intervals) were evaluated at the means of all other numeric variables, and for the intertidal range and Sand sediment type (the same trends occur for both variables among the remaining species and subzones).

temperature threshold above which management intervention is warranted. Note that the estimated marginal means for tidal range and sediment type were calculated using the `emmeans` R package [Lenth and Piaskowski, 2025].

Table 6: ANOVA table of covariates in the Wide Bay water quality and geographic model.

Effects	df	Chi-sq	p.value
90 Day Temperature	1	54.926	0.000
Sediment Type	4	16.638	0.002
Tidal Range	2	3.870	0.144
90 Day Turbidity	1	1.880	0.170
Mean Flood Turbidity/Years Since Flood One ²	1	0.003	0.960

We may inspect model diagnostics to better understand the overall model fit. Here we discuss how to partition the model into different components of variability, infer decay behavior of spatial autocorrelation, analyze standardized residuals to assess model assumptions, and identify influential observations that have a sizable impact on model fit. From Table 7, we see that of the total model variability, 5.4% is attributable to the explanatory variables (this quantity is sometimes called the Pseudo R-squared), 54.2% is attributable to the spatial random error, 5.8% is attributable to the zone class, and the remaining variability is independent random error. The estimated exponential spatial range (ϕ) is 14,757 meters, which suggests two total cover observations are approximately (spatially) uncorrelated at a distance of $3\phi \approx 45\text{km}$, or forty-five kilometers. While the raw residuals will inform our resilience modeling in Section 5, we can check certain model assumptions using the standardized residuals, which have been decorrelated. Generally, the standardized residuals are centered around zero, though there

is a slight right skew suggesting some observed total cover values are larger than expected (according to the model). Observations are called “influential” if they highly influence model fit. Influence is often measured via Cook’s distance, and no observations exceed the often-used Cook’s distance threshold value of one [Cook and Weisberg, 1982].

Table 7: Variance components in the Wide Bay water quality and geographic model.

Variance Component	Percent (of Total Variability)	Parameter Value
Covariates	5.4%	NA
Spatial Effect	54.2%	28.8
Independent Error	34.6%	18.4
Zone Class	5.8%	14757.0

4.2 Block Kriging

A tremendous benefit of using the spatial linear model is that block Kriging can be directly applied to predict the mean total seagrass cover bay-wide for any combination of explanatory variables and geographic regions, even if these variables were not directly used in modeling. Figure 5 and Figure 6 show that generally, there is no evidence total cover in the zone classes or conservation zones is decreasing since the floods (in financial year 2021-2022). While zone class was included in the model as a random effect, conservation zone type was not – via block Kriging, we can still quantify broad trends in total seagrass cover. Figure 7 shows that total seagrass cover is improving in the deep subtidal and intertidal classes since the floods, while Figure 8 shows that total seagrass cover is recovering bay-wide since the floods. Another useful application of block Kriging is scenario modeling, which can be used to characterize the impact of various future changes to broad-scale climatic patterns.

4.3 Alternative Approaches

In addition to the spatial linear model, we considered several different modeling approaches (all using the same explanatory variables): a nonspatial linear model, a spatial beta regression model, a spatial gamma regression model, a random forest model, and a spatial binomial model. The nonspatial linear model provided a useful baseline to compare the spatial linear model against, elucidating the impact of spatial dependence on model fit. The spatial beta regression model is a generalized linear model used to characterize proportion data between zero and one. To adhere to this restriction, total cover was 1) scaled to a proportion and 2) if exactly zero (one), set to 0.01 (0.99). The spatial gamma regression model is a generalized linear model used to characterize skewed positive data. To adhere to this restriction, if total cover was exactly zero, it was set to 0.01. The random forest provided a useful comparison to a machine learning approach, quite different than the aforementioned statistical approaches. We characterized the directionality of random forest effects using partial dependence plots




Figure 5: Average total cover predictions by zone class. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.




Figure 6: Average total cover predictions by conservation zone. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.




Figure 7: Average total cover predictions by tidal range. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.




Figure 8: Average total cover predictions in Wide Bay. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.

(Figure 9). Finally, the binomial model is a generalized linear model used to characterize presence/absence (success/failure) data. To adhere to this restriction, total cover was 1) given a value of zero if there was no seagrass or 2) given a value of one if there was any seagrass.

All six approaches found notable, negative relationships between temperature and total cover as well as notable to moderate relationships between sediment type and total cover as well as tidal range and total cover. The nonspatial linear model, random forest, and spatial beta regression model found notable, generally negative relationships between turbidity and total cover, although quite small in magnitude. From Table 8, the spatial linear model and random forest had the lowest out-of-sample root-mean-squared-prediction error, followed by the generalized linear models and then by the nonspatial linear model. Random forest partial dependence plots [Greenwell, 2017], which characterize the marginal behavior of a variable in a random forest model, suggested linear effects of the model variables to be reasonable. Given this context, we chose the spatial linear model as the final model, although it is certainly reasonable to justify other modeling choices that enforce specific, more realistic restrictions on the response distribution like the spatial beta regression model (enforcing proportions).





Figure 9: Random forest partial dependence for total cover as a function of 90 day mean temperature.

Table 8: Wide Bay out of sample prediction metrics for the various approaches. Mean bias (MBias) tended to be small relative to root-mean-squared-prediction error (RMSPE).

Approach	MBias	RMSPE
RF	-0.03	4.54
SPLM	0.00	4.70
SPGLM-Gamma	1.18	5.12
SPGLM-Beta	-0.16	5.31
LM	0.00	5.45

5 Resilience Modeling



ter quality and geographic drivers on total seagrass cover. It does not, however, quantify resilience on a site-specific scale. [Gilby et al. \[2023\]](#) provide a framework to quantify resilience on a site-specific scale. They suggest identifying hotspots (i.e., brightspots), sites that are performing better than expected (according to some model fit), as well as coldspots, sites that are performing worse than expected (according to some model fit). They identify hotspots and coldspots based on deviations exceeding expected bounds (based on confidence intervals) based on the model fit. They argue these hotspots and coldspots provide significant utility, both in better understanding the drivers of site-specific resilience as well as informing subsequent management action.

Table 9: Wide Bay resilience variables considered for modeling.

Variable	Effect	Included	Nonlinearity Degree
Coral Habitat Total Area	Fixed	Yes	2
Seagrass Habitat Total Area	Fixed	Yes	2
Mangrove Habitat Total Area	Fixed	Yes	2
Gastropod Habitat Total Area	Fixed	No	NA
Distance to Coral Habitat	Fixed	Yes	3
Distance to Seagrass Habitat	Fixed	Yes	2
Distance to Mangrove Habitat	Fixed	Yes	2
Distance to Gastropod Habitat	Fixed	Yes	2
Species Richness	Fixed	Yes	1
Habitat Count	Fixed	Yes	2
Dominant Species	Fixed	Yes	NA
Bathymetry Mean Depth	Fixed	Yes	2
Bathymetry Mean Curvature	Fixed	No	NA
Bathymetry Mean Slope	Fixed	Yes	2
Zone Class	Random	Yes	NA

Building from [Gilby et al. \[2023\]](#), we use the residuals from the model in Section 4 as the basis for studying site-specific resilience, linking the values of these residuals to site-specific variables like species habitat, species richness, and bathymetry via a separate spatial linear model, with a random effect for zone class. These models capture varying nonlinear behavior in the residuals as a function of the resilience variables using B-spline basis functions [[de Boor, 1978](#), [Eilers and Marx, 1996](#)] whose degree is informed by random forest partial dependence plots (built from a random forest model on the residuals). While [Gilby et al. \[2023\]](#) focused more on specific hotspots, linking environmental variables to those sites, we used our framework to provide a different set of information – the characterizing of the most important drivers of resilience, on average. Because we were interested in hypothesis testing, not parsimony, we included all potential variables as fixed effects in the model aside from “Gastropod Habitat Total Area” and “Bathymetry Mean Curvature” because there were some challenges processing these data which prohibited their use. These included area of and distance to coral, seagrass, mangrove,

and gastropod habitat, species richness and dominant species, habitat count, and bathymetry metrics (Table 9), which have the following structure:

- The “Coral Habitat Total Area” is numeric from 0 msq to 367,906 msq.
- The “Seagrass Habitat Total Area” is numeric from 0 msq to 773,161 msq.
- The “Mangrove Habitat Total Area” is numeric from 0 msq to 241,057 msq.
- The “Distance to Coral Habitat” is numeric from 0 m to 30,563 m.
- The “Distance to Seagrass Habitat” is numeric from 0 m to 7,049 m.
- The “Distance to Mangrove Habitat” is numeric from 0 m to 38,867 m.
- The “Distance to Gastropod Habitat” is numeric from 0 m to 121,609 m.
- The “Species Richness” is numeric (integer) from 0 unique species to 3 unique species.
- The “Habitat Count” is numeric (integer) from 0 unique habitats to 3 unique habitats.
- The “Dominant Species” has five levels: “Absent”, “HD”, “HO”, “HS”, and “ZMHU”.
- The “Bathymetry Mean Depth” is numeric from -30.58 m to 0.25 m.
- The “Bathymetry Mean Slope” is numeric from 0.004 degrees to 3.93 degrees.
- The “Zone Class” variable has six levels: Central, East, Great Sandy Strait, Hervey Bay, Southern, Southwest (Figure 3).

5.1 Model Fit

Table 10: ANOVA table of covariates in the Wide Bay resilience model.

Effects	df	Chi-sq	p.value
Species Richness	1	238.769	0.000
Dominant Species	4	68.448	0.000
Distance to Coral Habitat	2	51.856	0.000
Distance to Gastropod Habitat	2	3.196	0.202
Distance to Mangrove Habitat	2	2.913	0.233
Seagrass Habitat Total Area	2	2.834	0.242
Mangrove Habitat Total Area	2	1.161	0.560
Bathymetry Mean Slope	2	0.699	0.705
Distance to Seagrass Habitat	2	0.551	0.759
Bathymetry Mean Depth	2	0.435	0.804
Coral Habitat Total Area	2	0.094	0.954
Habitat Count	2	0.042	0.979

From Table 10, we find strong evidence (p -value < 0.001) that species richness, dominant species, and distance to coral habitat are associated with resilience:

- Species Richness: An increase of species richness by one species was associated with an average increase in resilience by approximately five points (Figure 11).

- **Dominant Species:** The estimated (marginal) average resilience score is largest for HS (-0.17), followed by HD (-2.9), HO (-3.2), and ZMHU (-3.9). The overall ranking is most important, as the negative average scores are in part likely due to the heavy skew in the data.
- **Distance from Coral Habitat:** From zero to around 10,000 m from coral habitat, there is a decreasing trend with resilience (i.e., increasing distance decreases resilience). Upwards of 10,000 m, there is an increasing trend with resilience (i.e., increasing distance increases resilience).

We found little evidence ($p\text{-value} > 0.1$) the remaining variables were associated with resilience (after controlling for other modeled variables).




Figure 11: Average species richness (left) and distance to coral habitat (right) effects on resilience. Predictions (and confidence intervals) were evaluated at the means of all other numeric variables and for the ZMHU species in the Central zone (similar trends occur for both variables among the remaining species and zones).

5.2 Alternative Approaches

Another way we studied resilience was using a random forest (fit to the same residuals). Largely, we found similar results as the spatial linear model. Notably, for the random forest:

- Species richness was most important, with a general increasing trend in resilience with richness.
- Distance from mangrove habitat was second-most important, with a general increasing trend in resilience with distance from mangroves.
- Dominant species was third-most important, with HS being the most resilient followed by HO, ZMHU, and HD.
- Bathymetry mean depth was fourth-most important, with a general increasing trend in resilience with depth.
- Distance from coral habitat was fifth-most important, with a similar quadratic trend (Figure 12) as in the spatial linear model.
- The remaining variables (gastropod distance, seagrass area, bathymetry slope, seagrass distance, zone class, habitat count, mangrove area, and coral area) were least important.

Other alternatives to explore involve using the residuals from water quality and geographic models fit using the spatial generalized linear models and random forests. These residuals could be analyzed using spatial linear models (recall that the residuals are unconstrained) and random forests. We expect these alternative approaches to yield generally similar results, but there could be nontrivial differences given the heavy skew of the data, however.




Figure 12: Random forest partial dependence for total cover as a function of distance from coral habitat.

6 Limitations and Future Suggestions

While the spatial linear model was useful for characterizing broad patterns related to tidal range, sediment type, temperature, and turbidity and site-specific patterns capturing resilience, all while accounting for spatial dependence, there are some limitations. First, the eReefs water quality data are modeled, not directly observed in the field, potentially propagating modeling error. Second, there are many correlated water quality variables, making it hard (for any model) to isolate the effects of each component of water quality pre- and post-flood. Third,

there is a lack of baseline mean salinity in the 90 days prior to sampling, which makes it challenging to directly study the effect of mean salinity during the flood period. Fourth, there is a lack of data immediately after each flood – nearly all of the data are after both floods, making it challenging to isolate the contribution from each flood separately. Fifth, there was no pre-flood data, making it challenging to compare against a pre-flood baseline. Sixth, there is a lack of additional data sources that could help quantify the impact of various explanatory variables on total cover pre- and post-flood, potentially varying across both space and time (e.g., light attenuation, biological interactions, etc.). Seventh, the locations at which data were collected were not randomly sampled. Randomly sampling sites helps to guard against sampling preferentially [Diggle et al., 2010] and provides more robust and representative data sources [Dumelle et al., 2022]. Eighth, the data have heavy skew and many observations without seagrass. While we explored several approaches and ultimately preferred the spatial linear model because of its interpretability and utility with block Kriging, the corresponding assumptions are less reliable than if the data had a more dense set of nonzero seagrass cover observations. No matter the approach ultimately used to analyze the data, highly skewed data with many zeros complicate modeling.

7 Revisiting the Research Questions

Research Question 1: Can we identify water quality covariates that act as triggers for seagrass cover management responses? Are these triggers different based on whether they result from a specific event (e.g., flood) or are ambient?

Temperature was the primary water quality variable associated with total seagrass cover in Wide Bay, with a one-degree increase in 90-day mean temperature associated with a 0.5 percentage point decrease in cover. This ambient temperature signal was statistically significant and ecologically interpretable, as elevated temperatures may compound the effects of other stressors on seagrass. In contrast, neither ambient turbidity nor flood turbidity showed strong statistical evidence of association with total cover after accounting for other variables. This could result in part due to the absence of pre-flood baseline data, which made it difficult to isolate the contribution of flood-driven turbidity from background conditions. It could also be in part due to the challenges associated with analyzing highly skewed data with many zeros. Across tidal ranges, estimated average total cover was modest throughout, and sediment type was a significant predictor, whereby muddy substrates supported slightly higher average cover than sand, sandy mud, or rock.

Research Question 2: Did seagrass cover improve after the flood event(s)? If so, how and where?

Block kriging predictions show no evidence of declining total cover across zone classes or conservation zones since the 2021–2022 financial year. In fact, total cover seems to be notably improving in Hervey Bay as well as the General Use Zone and Marine National Park Zone in particular. Deep subtidal and intertidal tidal ranges showed signs of modest improvement

over the monitoring period, while bay-wide predictions suggest a gradual positive trend since the floods. However, these findings should be interpreted with context, as the absence of pre-flood baseline data means it is not possible to assess whether the observed cover levels represent recovery relative to a pre-disturbance state or simply reflect the current trajectory of the ecosystem.

Research Question 3: Can we identify site-specific resilience indicators, which describe where recovery is likely to be enhanced or impeded? What factors contribute to seagrass resilience?

We identified several site-specific indicators of seagrass resilience in Wide Bay. Species richness and dominant species identity were the strongest indicators of resilience, with each additional species associated with an average increase in resilience of approximately five points, and sites dominated by HS tending to be most resilient, followed by HD, HO, and ZMHU. Distance to coral habitat was also a strong resilience indicator, exhibiting a nonlinear pattern — resilience declined with increasing distance from coral up to approximately 10,000 m, then increased beyond that threshold, a pattern consistent with coral and seagrass occupying different but adjacent positions along Wide Bay’s water quality and substrate gradient. These findings were broadly corroborated by the random forest, which ranked species richness as the most important predictor, followed by distance to mangrove habitat, dominant species, bathymetry mean depth, and distance to coral habitat. Together, these results suggest that management actions protecting structurally diverse, species-rich meadows are likely to be most effective at improving resilience, while sites with low species richness (or no seagrass) may require more active intervention to support recovery.

8 Discussion

This report applies a rigorous, spatially explicit statistical framework to characterize the drivers, recovery, and resilience of seagrass cover in Wide Bay across a three-year monitoring period that included two successive flood events. The analytical approach taken here is well-suited to the structure and scale of the data, and the findings have direct implications for adaptive management of the bay’s seagrass ecosystem.

The use of spatial linear models as the primary analytical framework is well-justified from both statistical and ecological perspectives. Seagrass cover observations collected across a bay are inherently spatially structured, as nearby sites share similar environmental conditions, water quality and disturbance histories, and ecological communities. Models that ignore this dependence will likely provide insufficient results. By explicitly modeling spatial dependence via a spatial covariance function, the analyses presented here produce more reliable estimates of covariate effects and more honest characterizations of uncertainty than a conventional non-spatial model would provide. These models are typically also more interpretable and characterize uncertainty more effectively than machine learning models. The inclusion of zone class as a random effect further accounts for structured variation among management-relevant regions

of the bay, ensuring that covariate estimates reflect within-class patterns rather than being confounded by between-class differences in baseline cover. The block Kriging helps understand bay-wide trends and can immediately characterize bay-wide predictions on the total seagrass cover scale (and not a transformed scale), providing a flexible platform for scenario modeling under future conditions.

The two-stage modeling framework — first modeling total cover as a function of water quality and geographic drivers, then modeling the residuals from that model as a function of site-level resilience variables — is a principled approach to separating system-wide environmental forcing from site-specific ecological capacity, building directly on the framework of [Gilby et al. \[2023\]](#). The extension to average driver characterization, rather than discrete hotspot identification, provides the more generalizable insight needed to inform system-wide conservation strategy in a bay of this scale. The robustness of the findings is supported by the consistency of results across multiple modeling approaches — the spatial linear model, spatial beta and gamma regression models, and random forest all identified temperature and sediment type as important predictors in the water quality model, and species richness and dominant species as dominant predictors in the resilience model. Agreement between parametric and nonparametric approaches under these circumstances provides meaningful corroboration that the identified relationships reflect genuine ecological signal rather than model-specific artifacts.

The two-stage modeling framework – first modeling total cover as a function of water quality and geographic drivers, then modeling the residuals from that model as a function of site-level resilience variables – is a structured approach to separating bay-wide environmental patterns from site-specific ecological resilience capacity. This builds directly on the framework proposed by [Gilby et al. \[2023\]](#) to help understand drivers of average resilience and provide useful insight needed to inform broader conservation strategies. The robustness of the findings is supported by the general similarity of results across multiple modeling approaches – for example, all water quality and geographic approaches found identifying temperature and sediment type as important while both resilience approaches found species richness, dominant species, and distance to coral habitat as important in the resilience model. Together, these results provide evidence that the relationships identified are genuine rather than resulting solely from specific modeling response frameworks.

Several limitations bear directly on the interpretation of results. Most critically, the absence of pre-flood baseline data makes it difficult, if not impossible, to assess whether observed cover levels represent genuine recovery or simply the current trend of the ecosystem. The near-complete overlap of post-flood-one and post-flood-two sampling also prevents isolation of individual contributions from each flood, an important gap given that the two events may have different effects on different areas of the bay. The heavy skew of the cover data, with many zero observations, present additional modeling challenges. While the spatial linear model is a useful tool here that can capture complex nonlinearities and spatial dependence simultaneously, perhaps future work could further explore alternative approaches such as the spatial beta regression model which enforces biologically realistic bounds on the response, or a machine learning algorithm like random forest. One downside of an alternative approach such as

these, however, is that block Kriging is not readily available on the response scale, often the scale of interest (spatial generalized linear models), or uncertainty quantification is challenging (machine learning algorithms).

Overall, the results of this report provide a quantitative, spatially structured foundation for adaptive management of Wide Bay seagrass. The understanding of important water quality and geographic variables, average predictions via block Kriging, and the study of site-level resilience indicators provides evidence that can inform targeted, context-specific intervention through the bay.

References

- Benjamin M. Bolker. Linear and generalized linear mixed models. In Gordon A. Fox, Simoneta Negrete-Yankelevich, and Vinicio J. Sosa, editors, *Ecological Statistics: Contemporary Theory and Application*, pages 309–333. Oxford University Press, Oxford, 2015. doi: 10.1093/acprof:oso/9780199672547.003.0014.
- Wagner Hugo Bonat and Paulo Justiniano Ribeiro Jr. Practical likelihood analysis for spatial generalized linear mixed models. *Environmetrics*, 27(2):83–89, 2016. doi: 10.1002/env.2375.
- Leo Breiman. Random forests. *Machine Learning*, 45(1):5–32, 2001. doi: 10.1023/A:1010933404324.
- George Casella and Roger L. Berger. *Statistical Inference*. Chapman and Hall/CRC, 2nd edition, 2024.
- John M. Chambers and Trevor J. Hastie. *Statistical Models in S*. Routledge, 1992.
- Ole F. Christensen and Rasmus Waagepetersen. Bayesian prediction of spatial count data using generalized linear mixed models. *Biometrics*, 58(2):280–286, 2002. doi: 10.1111/j.0006-341X.2002.00280.x.
- R. Dennis Cook and Sanford Weisberg. *Residuals and Influence in Regression*. Chapman and Hall, New York, 1982.
- Noel Cressie. Fitting variogram models by weighted least squares. *Journal of the International Association for Mathematical Geology*, 17(5):563–586, 1985. doi: 10.1007/BF01032109.
- Noel Cressie. The origins of kriging. *Mathematical Geology*, 22(3):239–252, 1990. doi: 10.1007/BF00889887.
- Noel Cressie. *Statistics for Spatial Data*. Wiley, Hoboken, NJ, 1993. doi: 10.1002/9781119115151.
- Noel Cressie. Block kriging for lognormal spatial processes. *Mathematical Geology*, 38(4):413–443, 2006. doi: 10.1007/s11004-005-9022-8.
- Frank C. Curriero and Subhash Lele. A composite likelihood approach to semivariogram estimation. *Journal of Agricultural, Biological, and Environmental Statistics*, 4(1):9–28, 1999.
- D. Richard Cutler, Thomas C. Edwards, Karen H. Beard, Adele Cutler, Kyle T. Hess, Jacob Gibson, and Joshua J. Lawler. Random forests for classification in ecology. *Ecology*, 88(11):2783–2792, 2007. doi: 10.1890/07-0539.1.
- Carl de Boor. *A Practical Guide to Splines*. Springer-Verlag, New York, 1978.

- Peter J. Diggle and Paulo Justiniano Ribeiro Jr. *Model-Based Geostatistics*. Springer Series in Statistics. Springer, New York, 2007. ISBN 9780387329072. doi: 10.1007/978-0-387-48536-2.
- Peter J. Diggle, Raquel Menezes, and Ting-li Su. Geostatistical inference under preferential sampling. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 59(2): 191–232, 2010. doi: 10.1111/j.1467-9876.2009.00701.x.
- Michael Dumelle, Matt Higham, Jay M. Ver Hoef, Anthony R. Olsen, and Lisa Madsen. A comparison of design-based and model-based approaches for finite population spatial sampling and inference. *Methods in Ecology and Evolution*, 13(9):2018–2029, 2022. doi: 10.1111/2041-210X.13919.
- Michael Dumelle, Matt Higham, and Jay M. Ver Hoef. spmodel: Spatial statistical modeling and prediction in R. *PLOS ONE*, 18(3):e0282524, 2023. doi: 10.1371/journal.pone.0282524.
- Paul H. C. Eilers and Brian D. Marx. Flexible smoothing with B-splines and penalties. *Statistical Science*, 11(2):89–121, 1996. doi: 10.1214/ss/1038425655.
- Julian James Faraway. *Practical Regression and ANOVA Using R*. University of Bath, Bath, 2002. URL <https://cran.r-project.org/doc/contrib/Faraway-PRA.pdf>.
- Eric W. Fox, Ryan A. Hill, Scott G. Leibowitz, Anthony R. Olsen, Darren J. Thornbrugh, and Marc H. Weber. Assessing the accuracy and stability of variable selection methods for random forest modeling in ecology. *Environmental Monitoring and Assessment*, 189(7):316, 2017. doi: 10.1007/s10661-017-6025-0.
- Eric W. Fox, Jay M. Ver Hoef, and Anthony R. Olsen. Comparing spatial regression to random forests for large environmental data sets. *PLOS ONE*, 15(3):e0229509, 2020. doi: 10.1371/journal.pone.0229509.
- Ben L. Gilby, Andrew D. Olds, Rod M. Connolly, Paul S. Maxwell, Christopher J. Henderson, and Thomas A. Schlacher. Seagrass meadows shape fish assemblages across estuarine seascapes. *Marine Ecology Progress Series*, 588:179–189, 2018. doi: 10.3354/meps12394.
- Ben L. Gilby, Lucy A. Goodridge Gaines, Hayden P. Borland, Christopher J. Henderson, Jesse D. Mosman, Andrew D. Olds, and Hannah J. Perry. Drivers of ecological condition identify bright spots and sites for management across coastal seascapes. *Estuaries and Coasts*, 46(4):906–924, 2023. doi: 10.1007/s12237-022-01164-z.
- Brandon M. Greenwell. pdp: An R package for constructing partial dependence plots. *The R Journal*, 9(1):421–436, 2017. doi: 10.32614/RJ-2017-016. URL <https://journal.r-project.org/archive/2017/RJ-2017-016/index.html>.
- David A. Harville. Maximum likelihood approaches to variance component estimation and to related problems. *Journal of the American Statistical Association*, 72(358):320–338, 1977. doi: 10.1080/01621459.1977.10480998.

- Trevor Hastie, Robert Tibshirani, and Jerome Friedman. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. Springer Series in Statistics. Springer, 2nd edition, 2009. doi: 10.1007/978-0-387-84858-7. URL <https://hastie.su.domains/ElemStatLearn/>.
- Matthew J. Heaton, Andrew Millane, and Jake S. Rhodes. A scalable spatial decorrelation preprocessing approach for machine and deep learning. *Journal of Data Science*, pages 1–15, 2025.
- Charles R. Henderson. Best linear unbiased estimation and prediction under a selection model. *Biometrics*, 31(2):423–447, 1975. doi: 10.2307/2529430.
- Gareth James, Daniela Witten, Trevor Hastie, and Robert Tibshirani. *An Introduction to Statistical Learning with Applications in R*. Springer, 2017. doi: 10.1007/978-1-4614-7138-7.
- Kieryn Kilminster, Kathryn McMahon, Michelle Waycott, Gary A. Kendrick, Peter Scanes, Len McKenzie, Katherine R. O’Brien, Mitchell Lyons, Angus Ferguson, Paul Maxwell, et al. Unravelling complexity in seagrass systems for management: Australia as a microcosm. *Science of the Total Environment*, 534:97–109, 2015. doi: 10.1016/j.scitotenv.2015.04.061.
- Max Kuhn and Julia Silge. *Tidy Modeling with R: A Framework for Modeling in the Tidyverse*. O’Reilly Media, 2022. URL <https://www.tmw.org>.
- Russell V. Lenth and Julia Piaskowski. *emmeans: Estimated Marginal Means, aka Least-Squares Means*, 2025. URL <https://CRAN.R-project.org/package=emmeans>. R package version 2.0.1.
- Kathryn McMahon, Susan Bengtson Nash, Geoff Eaglesham, Jochen F. Müller, Norman C. Duke, and Steve Winderlich. Herbicide contamination and the potential impact to seagrass meadows in Hervey Bay, Queensland, Australia. *Marine Pollution Bulletin*, 51(1–4):325–334, 2005. doi: 10.1016/j.marpolbul.2004.10.048.
- Douglas C. Montgomery, Elizabeth A. Peck, and G. Geoffrey Vining. *Introduction to Linear Regression Analysis*. John Wiley & Sons, 6th edition, 2021.
- Katherine R. O’Brien, Michelle Waycott, Paul Maxwell, Gary A. Kendrick, James W. Udy, Angus J. P. Ferguson, Kieryn Kilminster, Peter Scanes, Len J. McKenzie, Kathryn McMahon, et al. Seagrass ecosystem trajectory depends on the relative timescales of resistance, recovery and disturbance. *Marine Pollution Bulletin*, 134:166–176, 2018. doi: 10.1016/j.marpolbul.2017.09.006.
- H. Desmond Patterson and Robin Thompson. Recovery of inter-block information when block sizes are unequal. *Biometrika*, 58(3):545–554, 1971. doi: 10.1093/biomet/58.3.545.
- Eric J. Pedersen, David L. Miller, Gavin L. Simpson, and Noam Ross. Hierarchical generalized additive models in ecology: an introduction with mgcv. *PeerJ*, 7:e6876, 2019. doi: 10.7717/peerj.6876.

- José C. Pinheiro and Douglas M. Bates. *Mixed-Effects Models in S and S-PLUS*. Springer, 2000. doi: 10.1007/b98882.
- A. R. Preen, W. J. Lee Long, and R. G. Coles. Flood and cyclone related loss, and partial recovery, of more than 1000 km² of seagrass in Hervey Bay, Queensland, Australia. *Aquatic Botany*, 52(1–2):3–17, 1995. doi: 10.1016/0304-3770(95)00491-H.
- David R. Roberts, Volker Bahn, Simone Ciuti, Mark S. Boyce, Jane Elith, Gurutzeta Guillera-Arroita, Severin Hauenstein, José J. Lahoz-Monfort, Boris Schröder, Wilfried Thuiller, et al. Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography*, 40(8):913–929, 2017. doi: 10.1111/ecog.02881.
- C. M. Roelfsema and S. R. Phinn. An integrated field and remote sensing approach for mapping seagrass cover, Moreton Bay, Australia. *Journal of Spatial Science*, 54(1):45–62, 2009. doi: 10.1080/14498596.2009.9635166.
- David Ruppert, Matt P. Wand, and Raymond J. Carroll. *Semiparametric Regression*. Cambridge University Press, Cambridge, 2003. doi: 10.1017/CBO9780511755453.
- Arkajyoti Saha, Sumanta Basu, and Abhirup Datta. Random forests for spatially dependent data. *Journal of the American Statistical Association*, 118(541):665–683, 2023. doi: 10.1080/01621459.2021.1950003.
- Oliver Schabenberger and Carol A. Gotway. *Statistical Methods for Spatial Data Analysis*. Chapman and Hall/CRC, 2017. doi: 10.1201/9781315275086.
- Michael L. Stein. *Interpolation of Spatial Data: Some Theory for Kriging*. Springer, 1999. doi: 10.1007/978-1-4612-1494-6.
- Andrew D. L. Steven, Mark E. Baird, Richard Brinkman, Nicholas J. Car, Simon J. Cox, Mike Herzfeld, Jonathan Hodge, Emlyn Jones, Edward King, Nugzar Margvelashvili, et al. eReefs: An operational information system for managing the Great Barrier Reef. *Journal of Operational Oceanography*, 12(sup2):S12–S28, 2019. doi: 10.1080/1755876X.2019.1656720.
- Mervyn Stone. Cross-validatory choice and assessment of statistical predictions. *Journal of the Royal Statistical Society: Series B (Methodological)*, 36(2):111–133, 1974. doi: 10.1111/j.2517-6161.1974.tb00994.x.
- Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1):267–288, 1996. doi: 10.1111/j.2517-6161.1996.tb02080.x.
- Waldo R. Tobler. A computer movie simulating urban growth in the Detroit region. *Economic Geography*, 46(sup1):234–240, 1970. doi: 10.2307/143141.
- Jay Ver Hoef. Sampling and geostatistics for spatial data. *Ecoscience*, 9(2):152–161, 2002. doi: 10.1080/11956860.2002.11682701.

- Jay M. Ver Hoef, Michael Dumelle, Matt Higham, Erin E. Peterson, and Daniel J. Isaak. Indexing and partitioning the spatial linear model for large data sets. *PLOS ONE*, 18(11): e0291906, 2023. doi: 10.1371/journal.pone.0291906.
- Jay M. Ver Hoef, Eryn Blagg, Michael Dumelle, Philip M. Dixon, Dale L. Zimmerman, and Paul Conn. Marginal inference for hierarchical generalized linear mixed models with patterned covariance matrices using the Laplace approximation. *Environmetrics*, 35(7):e2872, 2024. doi: 10.1002/env.2872.
- Russ Wolfinger, Randy Tobias, and John Sall. Computing Gaussian likelihoods and their derivatives for general linear mixed models. *SIAM Journal on Scientific Computing*, 15(6): 1294–1310, 1994. doi: 10.1137/0915079.
- Simon N. Wood. *Generalized Additive Models: An Introduction with R*. Chapman and Hall/CRC, 2nd edition, 2017. doi: 10.1201/9781315370279.
- Paul H. York, Catherine V. Bryant, and Michael Rasheed. *Post-Flood Seagrass Monitoring in Hervey Bay May 2022*. Department of Environment and Science, 2022.
- Hao Zhang. On estimation and prediction for spatial generalized linear mixed models. *Biometrics*, 58(1):129–136, 2002. doi: 10.1111/j.0006-341X.2002.00129.x.
- Hao Zhang and Dale L. Zimmerman. Towards reconciling two asymptotic frameworks in spatial statistics. *Biometrika*, 92(4):921–936, 2005. doi: 10.1093/biomet/92.4.921.
- Dale L. Zimmerman and Jay M. Ver Hoef. *Spatial Linear Models for Environmental Data*. Chapman and Hall/CRC, 2024. doi: 10.1201/9781003044680.