

Moreton Bay Seagrass Resilience Modelling

DRFA 21/22 Biodiversity Conservation Marine Final Report Project

Michael Dumelle and Erin E. Peterson

Table of contents

1	Introduction	1
2	Seagrass Cover Model	2
2.1	Methods	2
2.1.1	Data	2
2.1.2	Exploratory Analyses	5
2.1.3	Statistical Modeling	5
2.1.4	Block Kriging	10
3	Resilience Model	10
3.1	Data	14
3.2	Spatial Linear Model	16
3.3	Random forest model	17
4	Limitations and Future Suggestions	19
5	Revisiting the Research Questions	20
6	Summary	21
	References	22
7	Acknowledgements	26

1 Introduction

Over the past few decades, various scientific studies have informed our current understanding of trends, behaviors, and flood responses of seagrass cover in Moreton Bay [[Abal and Dennison](#),

1996, Udy and Dennison, 1997, Samper-Villarreal et al., 2016, Ovsyanikova et al., 2026] as well as Australia more widely [Kilminster et al., 2015, Gilby et al., 2018, O’Brien et al., 2018]. Remote sensing has offered new capabilities for seagrass cover mapping [Roelfsema and Phinn, 2009], while innovative statistical approaches have offered novel methods for studying resilience at site-specific scales [Gilby et al., 2023]. Our work adds to this body of literature by applying spatially explicit statistical models [Cressie, 1993] to study total percent seagrass cover (i.e., seagrass cover) drivers, resilience, and recovery in Moreton Bay and Wide Bay (Dumelle and Peterson 2026b). Specifically, we provide and contextualize answers to four specific research questions, applicable to Moreton Bay:

1. Can we identify water quality covariates that act as triggers for seagrass cover management responses? Are these triggers different based on whether they result from a specific event (e.g., flood) or ambient conditions?
2. Did seagrass cover improve after the flood event(s)? If so, how and where?
3. Can we identify site-specific resilience indicators, which describe where recovery is likely to be enhanced or impeded? If so, what factors contribute to seagrass resilience?

2 Seagrass Cover Model

2.1 Methods

2.1.1 Data

The seagrass cover dataset was provided by Science Under Sail Australia (SUSA; <https://www.scienceundersail.com.au/>). It included 7657 observations across three financial years (2020-2021, 2022-2023, and 2023-2024), where water quality and geographic covariates were also available. Of the 7657 observations, 1995 were collected in 2020-2021 (pre-flood), 3329 in 2022-2023 (1-year post-flood), and 2333 in 2023-2024 (2-years post-flood) (Table 1).

Table 1: The sample size and percent of total sample size for total percent seagrass cover observations in Moreton Bay, by financial year.

Financial Year	Sample Size	Percent of Total
2020-2021	1,995	26.1%
2022-2023	3,329	43.5%
2023-2024	2,333	30.5%
Total	7,657	100.0%

Seagrass cover is a percentage that ranges from 0 (no seagrass) to 100 (complete seagrass cover). shows total cover by financial year throughout the Bay. Total cover tends to be highest near

the eastern shoreline of the Bay and lowest in the middle of the Bay (Figure 1). Note that 2022-2023 (post-flood) had the the largest number of seagrass cover samples, as well as the only samples collected in the middle of the Bay.




Figure 1: Total percent seagrass cover in Moreton Bay by financial year.

EcoFutures Consulting used eReefs data [Steven et al., 2019] to generate a broad suite of ambient and flood-related turbidity, salinity, and temperature variables at biologically and ecologically relevant temporal windows (Table 2). Ambient water quality variables were based on the mean, minimum, and maximum values for three periods: 7, 30, and 90 days prior to sampling. Ambient variables were also calculated representing the frequency of days water quality thresholds were exceeded (above or below) and the maximum spell (i.e. duration) of days those thresholds were continuously exceeded during the time periods. Similar water quality variables representing flood-event conditions were also calculated for a 30-day flood period.

Table 2: Moreton Bay water quality variables considered in the exploratory analyses, including the variable name and measurement unit (in parentheses), the type of water-quality variable (Ambient, Flood), the time period in days, statistic calculated, and the threshold value used (if any).

Variable	Type	Period	Statistic	Threshold
Turbidity (NTU)	Ambient	30, 7, 90	Mean	
Turbidity (NTU)	Ambient	30, 7, 90	Frequency	> 3
Turbidity (NTU)	Ambient	30, 7, 90	Frequency	> 5
Turbidity (NTU)	Ambient	30, 7, 90	Frequency	> 6
Salinity (PSU)	Ambient	30, 7, 90	Frequency	< 20
Salinity (PSU)	Ambient	30, 7, 90	Frequency	< 25
Salinity (PSU)	Ambient	30, 7, 90	Frequency	< 30
Temperature (deg C)	Ambient	30, 7, 90	Mean	
Turbidity (NTU)	Flood	30	Mean	
Turbidity (NTU)	Flood	30	Max	
Turbidity (NTU)	Flood	30	Frequency	> 3
Turbidity (NTU)	Flood	30	Frequency	> 5
Turbidity (NTU)	Flood	30	Frequency	> 6
Turbidity (NTU)	Flood	30	Duration	> 3
Turbidity (NTU)	Flood	30	Duration	> 5
Turbidity (NTU)	Flood	30	Duration	> 6
Salinity (PSU)	Flood	30	Min	
Salinity (PSU)	Flood	30	Mean	
Salinity (PSU)	Flood	30	Frequency	< 20
Salinity (PSU)	Flood	30	Frequency	< 25
Salinity (PSU)	Flood	30	Frequency	< 30
Salinity (PSU)	Flood	30	Duration	< 20
Salinity (PSU)	Flood	30	Duration	< 25
Salinity (PSU)	Flood	30	Duration	< 30

Additional covariates were also provided by EcoFutures Consulting, which are used to represent geographic features, proximity to rivers, and depth, as well as the years since the flood and the seagrass growing season. These variables have the following structure:

- Sediment Type includes six categorical levels: Sand, Sandy Mud, Muddy Sand, Mud, Rock, and Other.
- Tidal Range has three categorical levels: Shallow Subtidal, Intertidal, and Deep Subtidal.
- Years Since Flood is numeric ranging from zero (pre flood) to 1.93 years post-flood (February, 2024).
- Nearest River includes six categorical levels: Brisbane River, Caboolture River, Logan River, Ningi Creek, Pimpama Coomera and Nerang Rivers, and Pine River.

- Distance to (Nearest River) Outlet is numeric and ranges from 0.24 to 34.97 kilometers.
- Growing Season is a categorical variable with two levels: Growing (September - December) and Not Growing (January - August).
- EHMP subzone has eleven categorical levels: Bramble Bay, Broadwater, Central Bay, Deception Bay, Eastern Banks, Moreton Bay Open Coastal Waters, North-Eastern Moreton Bay, Northern Banks Open Coastal Waters, South-Eastern Moreton Bay, Southern Bay, and Waterloo Bay (Figure 2).

2.1.2 Exploratory Analyses

Extensive exploratory analyses were undertaken to 1) understand the data structure, 2) identify outliers/errors, 3) examine correlations between potential covariates and the response (i.e. seagrass cover), as well as multicollinearity in the covariates, 4) evaluate model assumptions, and 5) visualise spatial and temporal patterns in the data.

Results revealed substantial collinearity among ambient and flood-related water-quality variables, which was expected given their nested temporal windows and threshold values. We retained only one variable from each correlated group. We also removed variables showing no meaningful association with seagrass cover or where the relationship lacked biological plausibility. Geographic variables exhibited less collinearity than water quality variables; however, we removed the distance to the nearest river outlet and the nearest river variables due to weak associations with seagrass cover. Growing season was excluded because a simple monthly proxy inadequately captures seasonal seagrass growth dynamics.

The final set of potential covariates considered in the modelling included mean turbidity in the 90 days prior to sampling, mean turbidity during the flood event, mean temperature in the 90 days prior to sampling, sediment type, tidal range, years since flood, and EHMP zone.

2.1.3 Statistical Modeling

We considered multiple modeling approaches in this study, which each accounted for the data distribution, nonlinear relationships between seagrass cover and covariates, and spatial dependence inherent in biological data collected through space and time to different degrees (Appendix 1). We used the full set of potential covariates to fit a nonspatial linear model, spatial linear model (i.e., spatial model), spatial beta regression model, spatial gamma regression model, and a random forest model.

There were several reasons for fitting these initial models. The nonspatial linear model provided a useful baseline to compare the spatial linear model against, elucidating the impact of spatial dependence on model fit. The spatial beta regression model is a generalized linear model used to characterize proportion data ranging between zero and one. To adhere to this restriction, total seagrass cover was 1) scaled to a proportion and 2) if exactly zero or one, set to 0.01 or 0.99, respectively. The spatial gamma regression model is a generalized linear




Figure 2: Observations and their EHMP Subzones in Moreton Bay.

model used to characterize skewed positive data. To adhere to this restriction, a value of 0.01 was added to seagrass cover observations that were equal to zero. Finally, the random forest model provided a useful assessment of a machine learning approach, which requires fewer distributional assumptions than the aforementioned statistical approaches and more naturally describes nonlinear relationships between variables (Appendix 1). We also characterized random forest effects using partial dependence plots [Greenwell, 2017], which isolate the marginal contribution of a variable while holding other variables at their mean values. Leave-one-out cross-validation predictions were then used to calculate the mean bias (MBias) and root mean square prediction error (RMSPE) for each model.

The results of this initial model comparison showed that the spatial linear model had the lowest RMSPE, followed by the random forest, generalized linear models, and the nonspatial linear model (Table 3). The nonspatial and spatial linear models also had much lower mean bias values than the random forest model, while the two generalized linear models produced unacceptably biased predictions. Nevertheless, all five approaches identified similar relationships between seagrass cover and the covariates, in the directions expected. Random forest partial dependence plots [Greenwell, 2017] suggested that linear effects of the model variables was reasonable. Given these results, we chose to use a spatial linear modeling framework for the Moreton Bay seagrass cover model.

Table 3: Moreton Bay out of sample prediction metrics for the various approaches. Mean bias (MBias) tended to be small relative to root-mean-squared-prediction error (RMSPE).

Approach	MBias	RMSPE
SPLM	-0.02	18.41
RF	-0.11	18.84
SPGLM-Beta	-2.76	19.60
SPGLM-Gamma	5.42	20.07
LM	0.00	22.54

A suite of spatial linear models (Appendix 1, Equation 1), with exponential spatial covariance functions, were fit using the `splm()` function in `spmodel`. Parameters were estimated using restricted maximum likelihood. Models were compared using XXX.

The final total seagrass cover model for Moreton Bay included the following covariates: tidal range, sediment type, 90-day mean temperature (prior to sampling), pre- versus post-flood, years since the flood, 90 day turbidity (prior to sampling), and mean turbidity during the flood period scaled by time since the flood squared (Table 4). Notice that we scaled mean turbidity during the flood period so that the effect of turbidity during the flood event was allowed to decay with time since the flood. Intuitively, this means that as the time since the flood increases, the impact of turbidity during the flood event decreases. We also considered scaling by time since flood (not squared), which produced similar results. A random effect for EHMP subzone was also included in the final model.




Figure 3: Random forest partial dependence for seagrass cover as a function of mean turbidity in the 90 days prior to sampling.

Table 4: Covariates in the final Moreton Bay total cover model, their type (numeric or categorical), and if categorical, the number of unique levels.

Variable	Type	No. Levels
90 Day Temperature	Numeric	NA
90 Day Turbidity	Numeric	NA
Mean Flood Turbidity/Years Since Flood ²	Numeric	NA
Sediment Type	Categorical	6
Tidal Range	Categorical	3

Table 5: ANOVA table of covariates in the Moreton Bay water quality and geographic model.

Effects	df	Chi-sq	p.value
Tidal Range	2	608.900	0.000
Sediment Type	5	113.720	0.000
90 Day Temperature	1	16.967	0.000
Mean Flood Turbidity/Years Since Flood ²	1	11.966	0.001
90 Day Turbidity	1	7.854	0.005

Table 5 shows the results from an analysis of variance applied to the fitted model. All covariates had large test statistics and small p -values (p -value < 0.01), suggesting that they had a significant relationship with total seagrass cover; here are some relevant takeaways:

- Tidal Range: The estimated (marginal) average total cover is largest in the intertidal (26.89%), followed by shallow subtidal (16.51%) and deep subtidal (4.88%).

- Sediment Type: The estimated (marginal) average total cover is largest in the other category (25.6%), followed by muddy sand (19.1%), sandy mud (17.8%), mud (14.3%), sand (14.2%), and rock (5.62%).
- 90 Day Temperature: A one-degree (C) increase in 90-day water temperature prior to sampling is associated with a decrease in average seagrass cover by 0.88%.
- Mean Flood Turbidity/(Years Since Flood)²: A one-unit increase in mean turbidity during the flood is associated with a decrease in average seagrass cover of 2.05%/(Years Since Flood)² (Figure 4).
- 90 Day Turbidity: A one-unit increase in 90-day turbidity (prior to sampling) is associated with a decrease in average seagrass cover by 7.18% (Figure 4).

Note that the estimated marginal means for tidal range and sediment type were calculated using the `emmeans` R package [Lenth and Piaskowski, 2025].




Figure 4: Average total seagrass percent cover as a function of 90-day mean turbidity (left) and flood turbidity (right). Estimates (and confidence intervals) were evaluated at one year post flood, at the means of all other numeric variables, and for the intertidal depth range and sand sediment type (the same trends occur for both variables among the remaining species and subzones).

We examined model diagnostics to better understand the overall model fit. First, we partitioned the model into different components of variability (Table 6) and found that 11.2% is attributable to the covariates (this quantity is sometimes called the Pseudo R-squared), 31.6% to the spatial random error, 7.3% to the EHMP subzone, and the remaining variability is independent random error. Second, we examined the decay behaviour of spatial autocorrelation.

The estimated exponential spatial range (ϕ) is 745.5 meters, which suggests two total seagrass cover observations are approximately (spatially) uncorrelated at a distance of $3\phi \approx 2\text{km}$, or two kilometers. Third, we analyzed the standardized model residuals, which have been decorrelated, to assess model assumptions and identify influential observations that had a sizable impact on model fit. Generally, the standardized residuals are centered around zero, though there is a slight right skew suggesting some observed total cover values are larger than expected (according to the model). Observations are referred to as “influential” if they highly influence model fit. Influence is often assessed a Cook’s distance threshold value of one [Cook and Weisberg, 1982]. No observations exceed this threshold.

Table 6: Variance components in the Moreton Bay seagrass model.

Variance Component	Percent (of Total Variability)	Parameter Value
Covariates	11.2%	NA
Spatial Effect	31.6%	182.1
Independent Error	49.9%	287.5
EHMP Subzone	7.3%	745.5

2.1.4 Block Kriging

A tremendous benefit of using the spatial linear model is that block kriging [Ver Hoef, 2002, Cressie, 2006] can be used to predict the mean total seagrass cover bay-wide for any combination of covariates and geographic regions, even if these variables were not directly used in the model (see Appendix 1 for additional details). Block kriging was undertaken, by financial year, for EHMP subzone, Marine Park zone, tidal range and for the whole of Moreton Bay.

Figure 5 and Figure 6 show there are several EHMP subzones and Marine Park zones that have recovered to pre-flood levels (e.g., Deception Bay, Marine National Park Zone), while others have not (e.g., Southern Bay, General Use Zone). When we examine recovery by tidal range (Figure 7), we see that seagrass cover is recovering less effectively in the intertidal range than the shallow or deep subtidal ranges. When we examine recovery for the whole of Moreton Bay (Figure 8), the model suggests that seagrass cover has improved since the 2022 flood.

3 Resilience Model

The seagrass cover model quantifies bay-wide effects of influential drivers on seagrass cover and enables assessment of recovery trajectories at management-relevant scales. However, it does not quantify site-specific resilience. Gilby et al. [2023] propose identifying hotspots (i.e. brightspots, sites performing better than expected) and coldspots (sites performing worse than expected) based on model residuals exceeding confidence interval bounds. They argue that hotspots and




Figure 5: Average total cover predictions by EHMP subzone. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.




Figure 6: Average total cover predictions by conservation zone. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.




Figure 7: Average total cover predictions by tidal range. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.




Figure 8: Average total cover predictions in Moreton Bay. Red dots indicate the mean from the raw data. Black dots indicate the predicted mean in the region alongside a prediction interval.

coldspots offer significant utility for understanding drivers of site-specific resilience and informing management decisions. [Gilby et al. \[2023\]](#) primarily focussed on specific hotspots, linking environmental variables to those sites. We build on that work by using the seagrass model residuals as a proxy for site-specific resilience, and then characterizing the most important drivers of that resilience, on average.

3.1 Data

We used the raw seagrass model residuals (Figure 9) as a the response variable in a second spatial linear model used to study site-specific resilience in Moreton Bay. Model covariates included variables describing seagrass diversity and dominant species, and bathymetry, as well as proximity to and area of coral, seagrass, mangrove, and gastropod habitat (Table 8). EHMP subzone was again included as a random effect. Because we were interested in hypotheses testing rather than parsimony, we included all potential covariates as fixed effects in the model. Potential covariates have the following structure:

- Coral Habitat Total Area is numeric from 0 msq to 616,180 meters squared (msq).
- Seagrass Habitat Total Area is numeric from 0 msq to 773,161 msq.
- Mangrove Habitat Total Area is numeric from 0 msq to 561,317 msq.
- Gastropod Habitat Total Area is numeric from 0 msq to 114,795 msq.
- Distance to Coral Habitat is numeric from 0 m to 35,387 m.
- Distance to Seagrass Habitat is numeric from 0 m to 11,382 m.

- Distance to Mangrove Habitat is numeric from 0 m to 27,617 m.
- Distance to Gastropod Habitat is numeric from 0 m to 32,034 m.
- Species Richness is numeric (integer) from 0 to 4 unique species.
- Habitat Count is numeric (integer) from 0 to 4 unique habitats.
- Dominant Species represents the dominant seagrass species and has seven levels: “Absent”, “CS”, “HD”, “HO”, “HS”, “SI”, and “ZMHU”.
- Bathymetry Mean Depth is numeric from -35.95 m to 0.79 m.
- Bathymetry Mean Curvature is numeric from -0.002 degrees to 0.0001 degrees.
- Bathymetry Mean Slope is numeric from 0.009 degrees to 4.27 degrees.
- EHMP subzone is a categorical variable with eleven levels: Bramble Bay, Broadwater, Central Bay, Deception Bay, Eastern Banks, Moreton Bay Open Coastal Waters, North-Eastern Moreton Bay, Northern Banks Open Coastal Waters, South-Eastern Moreton Bay, Southern Bay, and Waterloo Bay (Figure 2).

We expected there to be nonlinear relationships between the response and covariates and so we used B-spline basis functions [de Boor, 1978, Eilers and Marx, 1996] to describe these relationships. The degree of these functions was informed by partial dependence plots derived from a random forest model fitted to the seagrass model residuals (Table 7).




Figure 9: Seagrass model residuals for resilience analysis in Moreton Bay, by financial year.

Table 7: Covariates included in the resilience model, including whether they were a fixed or random effect, and the nonlinearity degree of the B-splines basis function.

Variable	Effect	Degree
Coral Habitat Total Area	Fixed	2
Seagrass Habitat Total Area	Fixed	2
Mangrove Habitat Total Area	Fixed	2
Gastropod Habitat Total Area	Fixed	2
Distance to Coral Habitat	Fixed	3
Distance to Seagrass Habitat	Fixed	2
Distance to Mangrove Habitat	Fixed	2
Distance to Gastropod Habitat	Fixed	2
Species Richness	Fixed	1
Habitat Count	Fixed	2
Dominant Species	Fixed	NA
Bathymetry Mean Depth	Fixed	2
Bathymetry Mean Curvature	Fixed	2
Bathymetry Mean Slope	Fixed	2
EHMP Subzone	Random	NA

3.2 Spatial Linear Model

Table 8: ANOVA table of covariates in the Moreton Bay resilience model.

Effects	df	Chi-sq	p.value
Dominant Species	6	1637.726	0.000
Species Richness	1	22.188	0.000
Bathymetry Mean Depth	2	21.876	0.000
Bathymetry Mean Curvature	2	8.999	0.011
Gastropod Habitat Total Area	2	5.155	0.076
Bathymetry Mean Slope	2	4.768	0.092
Coral Habitat Total Area	2	2.845	0.241
Seagrass Habitat Total Area	2	1.971	0.373
Distance to Mangrove Habitat	2	1.871	0.392
Distance to Seagrass Habitat	2	1.836	0.399
Habitat Count	2	0.722	0.697
Distance to Gastropod Habitat	2	0.613	0.736
Distance to Mangrove Habitat	2	0.536	0.765
Distance to Coral Habitat	3	0.718	0.869

The ANVOVA table for the final resilience model (Table 8) shows strong evidence (p -value < 0.001) that dominant species, species richness, and bathymetry mean depth are associated with resilience:

- Dominant Species: The estimated (marginal) average resilience score is largest for CS (38.4), followed by ZMHU (18.8), HS (10.8), SI (10.7), HO (3.7), and HD (-3.1).
- Species Richness: An increase of species richness by one species was associated with an average increase in resilience by approximately two units (Figure 10).
- Bathymetry Mean Depth: Deeper sites tended to be more resilient than shallower sites, on average (Figure 10).

We found moderate evidence (p -value < 0.1) that bathymetry mean curvature, gastropod habitat total area, and bathymetry mean slope are associated with resilience:

- Bathymetry Mean Curvature: Sites with more curvature tended to be less resilient than sites with less curvature.
- Gastropod Habitat Total Area: Sites with more gastropod habitat area tended to be more resilient than sites with less gastropod habitat area.
- Bathymetry Mean Slope: Sites with less extreme slopes (between zero and one) tended to have increasing resilience with slope, while sites with more extreme slopes (greater than one) tended to have decreasing resilience with slope.

We found little evidence (p -value > 0.1) the remaining variables were associated with resilience, after controlling for other model covariates.

3.3 Random forest model

As mentioned previously, we also fit a random forest model to the seagrass model residuals. Largely, we found similar results as the spatial linear model. Notably, for the random forest:

- Dominant species was most important, with a similar ranking of resilient species as for the spatial linear model approach (e.g., CS, ZMHU most resilient and Absent least resilient).
- Species richness was second-most important, with a general increasing trend in resilience with richness.
- Distance to gastropod habitat was third-most important, with a decreasing trend at relatively short distances ($< 15,000$ m) and an increasing trend at large distances ($> 15,000$ m).
- Bathymetry mean depth was fourth-most important variable, with a general decreasing trend with decreasing depth (Figure 11), which is similar to the spatial linear model results.




Figure 10: Average species richness (left) and bathymetry mean depth (right) effects on resilience. Predictions (and confidence intervals) were evaluated at the means of all other numeric variables and for the ZMHU species in the Southern Bay (similar trends occur for both variables among the remaining species and subzones).

- Distance to gastropod habitat, distance to mangrove habitat, distance to coral habitat, and seagrass area had similar importance as the remaining bathymetry variables.
- The remaining variables (distance to seagrass habitat, mangrove area, EHMP subzone, habitat count, coral area, and gastropod area) were least important.

The consistency between the two resilience models and their variable importance rankings reinforce the spatial linear model’s identification of dominant species, species richness, and bathymetry mean depth as key resilience drivers.




Figure 11: Random forest partial dependence for resilience as a function of bathymetry mean depth.

4 Limitations and Future Suggestions

While the spatial linear model was useful for characterizing broad patterns in seagrass cover related to tidal range, sediment type, temperature, and turbidity and site-specific patterns capturing resilience, all while accounting for spatial dependence, there are some limitations. First, the eReefs water quality covariates are based on modeled data that are not directly observed in the field, potentially propagating modeling error. Second, there are many correlated water-quality variables, making it difficult (for any model) to isolate the effects of each component of water quality pre- and post-flood. Third, the lack of a covariate describing mean salinity in the 90 days prior to sampling makes it challenging to directly study the effect of mean salinity during the flood period. Fourth, seagrass cover data was not collected in the first four months following the flood. It is therefore unclear whether the seagrass data collected in the 2022-2023 financial year fully describe flood impacts. Finally, covariates representing local-scale drivers of seagrass cover (e.g., light attenuation, biological interactions, etc.) were not available, which might explain some additional variability in the response through space and time.

5 Revisiting the Research Questions

Research Question 1: Can we identify water quality covariates that act as triggers for seagrass cover management responses? Are these triggers different based on whether they result from a specific event (e.g., flood) or are ambient?

We identified ambient 90-day turbidity, turbidity during the flood event, and ambient 90-day water temperature as significant water quality covariates in the seagrass cover model. Our results suggest that ambient and flood-related turbidity act through distinct mechanisms and at different magnitudes; a one-unit increase in 90-day mean turbidity was associated with a 7.18 percentage point decrease in total cover, while the effect of flood turbidity decayed with time since the flood. Across tidal ranges, intertidal sites supported the highest average total cover (26.89%), followed by shallow subtidal (16.51%) and deep subtidal (4.88%), suggesting that management thresholds may need to be calibrated differently depending on depth zone, as deeper sites with inherently lower cover may be disproportionately sensitive to turbidity increases. Similarly, sediment type affected baseline cover levels, as sites with muddy sand and sandy mud substrates supported higher average seagrass cover than sand or rock sites. Together, these results suggest that management triggers should differ depending on whether the driver is chronic ambient turbidity or an acute flood event, and should further account for the tidal and sediment context of the sites in question.

Research Question 2: Did seagrass cover improve after the flood event(s)? If so, how and where?

Total seagrass cover improved throughout Moreton Bay between 2022–2023 and 2023–2024, suggesting the system has been recovering since the flood. However, recovery was spatially heterogeneous; Deception Bay and the Marine National Park zone returned to pre-flood cover levels, while Southern Bay and the General Use Zone lagged considerably behind. Recovery was also relatively slow compared to the magnitude of the impact in the intertidal tidal range compared to shallow and deep subtidal ranges, suggesting that intertidal sites may warrant prioritized management attention immediately following floods.

Research Question 3: Can we identify site-specific resilience indicators, which describe where recovery is likely to be enhanced or impeded? What factors contribute to seagrass resilience?

We identified several site-specific indicators of seagrass resilience. Dominant species, species richness, and bathymetry mean depth were the strongest predictors of seagrass cover resilience, with sites dominated by CS or ZMHU species, higher species richness, and greater water depth tending to outperform model expectations and hence, infer greater resilience. Gastropod habitat area, bathymetric curvature, and slope provided moderate additional explanatory power. These findings may suggest that deeper, species-rich meadows dominated by CS and ZMHU exhibit a high level of resilience, while sites with low species richness and/or dominated by other species may require more active intervention to support recovery.

6 Summary

We applied a rigorous, spatially explicit statistical framework to characterize the drivers, recovery, and resilience of seagrass cover in Moreton Bay across a three-year monitoring period that included the 2022 Moreton Bay flood event. The analytical approach taken here is well-suited to the structure and scale of the data, and the findings have direct implications for adaptive management of the bay’s seagrass ecosystem.

The use of spatial linear models as the primary analytical framework is well-justified from both statistical and ecological perspectives. Total percent seagrass cover observations collected across the bay are inherently spatially structured, as nearby sites share similar environmental conditions, water quality and disturbance histories, as well as ecological communities. Models that ignore this dependence may well provide misleading results and inference. By explicitly modeling spatial dependence via a spatial covariance function, the analyses presented here produce more reliable estimates of covariate effects and more honest characterizations of uncertainty than a conventional non-spatial model provides. These models are typically also more interpretable and characterize uncertainty more effectively than machine learning models. The inclusion of EHMP subzone as a random effect further accounts for structured variation among management-relevant regions of the bay, ensuring that covariate estimates reflect within-subzone patterns rather than being confounded by between-subzone differences in baseline cover. The block kriging helps understand bay-wide trends and can immediately characterize bay-wide predictions on the original total seagrass cover scale (0-100), rather than a transformed scale required for generalized linear models; providing a flexible framework for scenario modeling under future flood conditions and climate scenarios.

The use of a two-stage modeling framework – first modeling total seagrass cover as a function of water quality and geographic drivers, then modeling the seagrass model residuals as a function of site-level resilience covariates – is a structured approach to separating bay-wide environmental patterns from site-specific ecological resilience capacity. This builds directly on the framework proposed by [Gilby et al. \[2023\]](#) to help understand drivers of average resilience and provide useful insight needed to inform broader conservation strategies. The robustness of the findings is further supported by the consistency of results across multiple modeling approaches (e.g., spatial beta regression model, random forest, etc.), which provides evidence that the relationships identified are genuine rather than resulting solely from specific modeling frameworks.

Overall, the results described in this report provide a quantitative, spatially structured foundation for adaptive decision making. The understanding of important water quality and geographic variables, average predictions via block Kriging, and the study of site-level resilience indicators provides data-driven evidence that can inform temporally targeted, context-specific interventions throughout the bay.

References

- E. G. Abal and W. C. Dennison. Seagrass depth range and water quality in southern Moreton Bay, Queensland, Australia. *Marine and Freshwater Research*, 47(6):763–771, 1996. doi: 10.1071/MF9960763.
- Benjamin M. Bolker. Linear and generalized linear mixed models. In Gordon A. Fox, Simoneta Negrete-Yankelevich, and Vinicio J. Sosa, editors, *Ecological Statistics: Contemporary Theory and Application*, pages 309–333. Oxford University Press, Oxford, 2015. doi: 10.1093/acprof:oso/9780199672547.003.0014.
- Wagner Hugo Bonat and Paulo Justiniano Ribeiro Jr. Practical likelihood analysis for spatial generalized linear mixed models. *Environmetrics*, 27(2):83–89, 2016. doi: 10.1002/env.2375.
- Leo Breiman. Random forests. *Machine Learning*, 45(1):5–32, 2001. doi: 10.1023/A:1010933404324.
- George Casella and Roger L. Berger. *Statistical Inference*. Chapman and Hall/CRC, 2nd edition, 2024.
- John M. Chambers and Trevor J. Hastie. *Statistical Models in S*. Routledge, 1992.
- Ole F. Christensen and Rasmus Waagepetersen. Bayesian prediction of spatial count data using generalized linear mixed models. *Biometrics*, 58(2):280–286, 2002. doi: 10.1111/j.0006-341X.2002.00280.x.
- R. Dennis Cook and Sanford Weisberg. *Residuals and Influence in Regression*. Chapman and Hall, New York, 1982.
- Noel Cressie. Fitting variogram models by weighted least squares. *Journal of the International Association for Mathematical Geology*, 17(5):563–586, 1985. doi: 10.1007/BF01032109.
- Noel Cressie. The origins of kriging. *Mathematical Geology*, 22(3):239–252, 1990. doi: 10.1007/BF00889887.
- Noel Cressie. *Statistics for Spatial Data*. Wiley, Hoboken, NJ, 1993. doi: 10.1002/9781119115151.
- Noel Cressie. Block kriging for lognormal spatial processes. *Mathematical Geology*, 38(4): 413–443, 2006. doi: 10.1007/s11004-005-9022-8.
- Frank C. Curriero and Subhash Lele. A composite likelihood approach to semivariogram estimation. *Journal of Agricultural, Biological, and Environmental Statistics*, 4(1):9–28, 1999.
- D. Richard Cutler, Thomas C. Edwards, Karen H. Beard, Adele Cutler, Kyle T. Hess, Jacob Gibson, and Joshua J. Lawler. Random forests for classification in ecology. *Ecology*, 88(11): 2783–2792, 2007. doi: 10.1890/07-0539.1.

- Carl de Boor. *A Practical Guide to Splines*. Springer-Verlag, New York, 1978.
- Peter J. Diggle and Paulo Justiniano Ribeiro Jr. *Model-Based Geostatistics*. Springer Series in Statistics. Springer, New York, 2007. ISBN 9780387329072. doi: 10.1007/978-0-387-48536-2.
- Michael Dumelle, Matt Higham, and Jay M. Ver Hoef. spmodel: Spatial statistical modeling and prediction in R. *PLOS ONE*, 18(3):e0282524, 2023. doi: 10.1371/journal.pone.0282524.
- Paul H. C. Eilers and Brian D. Marx. Flexible smoothing with B-splines and penalties. *Statistical Science*, 11(2):89–121, 1996. doi: 10.1214/ss/1038425655.
- Julian James Faraway. *Practical Regression and ANOVA Using R*. University of Bath, Bath, 2002. URL <https://cran.r-project.org/doc/contrib/Faraway-PRA.pdf>.
- Eric W. Fox, Ryan A. Hill, Scott G. Leibowitz, Anthony R. Olsen, Darren J. Thornbrugh, and Marc H. Weber. Assessing the accuracy and stability of variable selection methods for random forest modeling in ecology. *Environmental Monitoring and Assessment*, 189(7):316, 2017. doi: 10.1007/s10661-017-6025-0.
- Eric W. Fox, Jay M. Ver Hoef, and Anthony R. Olsen. Comparing spatial regression to random forests for large environmental data sets. *PLOS ONE*, 15(3):e0229509, 2020. doi: 10.1371/journal.pone.0229509.
- Ben L. Gilby, Andrew D. Olds, Rod M. Connolly, Paul S. Maxwell, Christopher J. Henderson, and Thomas A. Schlacher. Seagrass meadows shape fish assemblages across estuarine seascapes. *Marine Ecology Progress Series*, 588:179–189, 2018. doi: 10.3354/meps12394.
- Ben L. Gilby, Lucy A. Goodridge Gaines, Hayden P. Borland, Christopher J. Henderson, Jesse D. Mosman, Andrew D. Olds, and Hannah J. Perry. Drivers of ecological condition identify bright spots and sites for management across coastal seascapes. *Estuaries and Coasts*, 46(4):906–924, 2023. doi: 10.1007/s12237-022-01164-z.
- Brandon M. Greenwell. pdp: An R package for constructing partial dependence plots. *The R Journal*, 9(1):421–436, 2017. doi: 10.32614/RJ-2017-016. URL <https://journal.r-project.org/archive/2017/RJ-2017-016/index.html>.
- David A. Harville. Maximum likelihood approaches to variance component estimation and to related problems. *Journal of the American Statistical Association*, 72(358):320–338, 1977. doi: 10.1080/01621459.1977.10480998.
- Trevor Hastie, Robert Tibshirani, and Jerome Friedman. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. Springer Series in Statistics. Springer, 2nd edition, 2009. doi: 10.1007/978-0-387-84858-7. URL <https://hastie.su.domains/ElemStatLearn/>.
- Matthew J. Heaton, Andrew Millane, and Jake S. Rhodes. A scalable spatial decorrelation preprocessing approach for machine and deep learning. *Journal of Data Science*, pages 1–15, 2025.

- Charles R. Henderson. Best linear unbiased estimation and prediction under a selection model. *Biometrics*, 31(2):423–447, 1975. doi: 10.2307/2529430.
- Gareth James, Daniela Witten, Trevor Hastie, and Robert Tibshirani. *An Introduction to Statistical Learning with Applications in R*. Springer, 2017. doi: 10.1007/978-1-4614-7138-7.
- Kieryn Kilminster, Kathryn McMahon, Michelle Waycott, Gary A. Kendrick, Peter Scanes, Len McKenzie, Katherine R. O’Brien, Mitchell Lyons, Angus Ferguson, Paul Maxwell, et al. Unravelling complexity in seagrass systems for management: Australia as a microcosm. *Science of the Total Environment*, 534:97–109, 2015. doi: 10.1016/j.scitotenv.2015.04.061.
- Max Kuhn and Julia Silge. *Tidy Modeling with R: A Framework for Modeling in the Tidyverse*. O’Reilly Media, 2022. URL <https://www.tmwr.org>.
- Russell V. Lenth and Julia Piaskowski. *emmeans: Estimated Marginal Means, aka Least-Squares Means*, 2025. URL <https://CRAN.R-project.org/package=emmeans>. R package version 2.0.1.
- Douglas C. Montgomery, Elizabeth A. Peck, and G. Geoffrey Vining. *Introduction to Linear Regression Analysis*. John Wiley & Sons, 6th edition, 2021.
- Katherine R. O’Brien, Michelle Waycott, Paul Maxwell, Gary A. Kendrick, James W. Udy, Angus J. P. Ferguson, Kieryn Kilminster, Peter Scanes, Len J. McKenzie, Kathryn McMahon, et al. Seagrass ecosystem trajectory depends on the relative timescales of resistance, recovery and disturbance. *Marine Pollution Bulletin*, 134:166–176, 2018. doi: 10.1016/j.marpolbul.2017.09.006.
- E. Ovsyanikova, W. N. Venables, and J. W. Udy. Spatial and temporal modelling of seagrass distribution in Moreton Bay, Australia, based on long-term citizen science data. *Estuarine, Coastal and Shelf Science*, 328:109610, 2026. doi: 10.1016/j.ecss.2025.109610.
- H. Desmond Patterson and Robin Thompson. Recovery of inter-block information when block sizes are unequal. *Biometrika*, 58(3):545–554, 1971. doi: 10.1093/biomet/58.3.545.
- Eric J. Pedersen, David L. Miller, Gavin L. Simpson, and Noam Ross. Hierarchical generalized additive models in ecology: an introduction with mgcv. *PeerJ*, 7:e6876, 2019. doi: 10.7717/peerj.6876.
- José C. Pinheiro and Douglas M. Bates. *Mixed-Effects Models in S and S-PLUS*. Springer, 2000. doi: 10.1007/b98882.
- David R. Roberts, Volker Bahn, Simone Ciuti, Mark S. Boyce, Jane Elith, Gurutzeta Guillera-Arroita, Severin Hauenstein, José J. Lahoz-Monfort, Boris Schröder, Wilfried Thuiller, et al. Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography*, 40(8):913–929, 2017. doi: 10.1111/ecog.02881.

- C. M. Roelfsema and S. R. Phinn. An integrated field and remote sensing approach for mapping seagrass cover, Moreton Bay, Australia. *Journal of Spatial Science*, 54(1):45–62, 2009. doi: 10.1080/14498596.2009.9635166.
- David Ruppert, Matt P. Wand, and Raymond J. Carroll. *Semiparametric Regression*. Cambridge University Press, Cambridge, 2003. doi: 10.1017/CBO9780511755453.
- Arkajyoti Saha, Sumanta Basu, and Abhirup Datta. Random forests for spatially dependent data. *Journal of the American Statistical Association*, 118(541):665–683, 2023. doi: 10.1080/01621459.2021.1950003.
- Jimena Samper-Villarreal, Catherine E. Lovelock, Megan I. Saunders, Chris Roelfsema, and Peter J. Mumby. Organic carbon in seagrass sediments is influenced by seagrass canopy complexity, turbidity, wave height, and water depth. *Limnology and Oceanography*, 61(3): 938–952, 2016. doi: 10.1002/lno.10262.
- Oliver Schabenberger and Carol A. Gotway. *Statistical Methods for Spatial Data Analysis*. Chapman and Hall/CRC, 2017. doi: 10.1201/9781315275086.
- Michael L. Stein. *Interpolation of Spatial Data: Some Theory for Kriging*. Springer, 1999. doi: 10.1007/978-1-4612-1494-6.
- Andrew D. L. Steven, Mark E. Baird, Richard Brinkman, Nicholas J. Car, Simon J. Cox, Mike Herzfeld, Jonathan Hodge, Emlyn Jones, Edward King, Nugzar Margvelashvili, et al. eReefs: An operational information system for managing the Great Barrier Reef. *Journal of Operational Oceanography*, 12(sup2):S12–S28, 2019. doi: 10.1080/1755876X.2019.1656720.
- Mervyn Stone. Cross-validatory choice and assessment of statistical predictions. *Journal of the Royal Statistical Society: Series B (Methodological)*, 36(2):111–133, 1974. doi: 10.1111/j.2517-6161.1974.tb00994.x.
- Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1):267–288, 1996. doi: 10.1111/j.2517-6161.1996.tb02080.x.
- Waldo R. Tobler. A computer movie simulating urban growth in the Detroit region. *Economic Geography*, 46(sup1):234–240, 1970. doi: 10.2307/143141.
- James W. Udy and William C. Dennison. Growth and physiological responses of three seagrass species to elevated sediment nutrients in Moreton Bay, Australia. *Journal of Experimental Marine Biology and Ecology*, 217(2):253–277, 1997. doi: 10.1016/S0022-0981(97)00060-3.
- Jay Ver Hoef. Sampling and geostatistics for spatial data. *Ecoscience*, 9(2):152–161, 2002. doi: 10.1080/11956860.2002.11682701.
- Jay M. Ver Hoef, Michael Dumelle, Matt Higham, Erin E. Peterson, and Daniel J. Isaak. Indexing and partitioning the spatial linear model for large data sets. *PLOS ONE*, 18(11): e0291906, 2023. doi: 10.1371/journal.pone.0291906.

- Jay M. Ver Hoef, Eryn Blagg, Michael Dumelle, Philip M. Dixon, Dale L. Zimmerman, and Paul Conn. Marginal inference for hierarchical generalized linear mixed models with patterned covariance matrices using the Laplace approximation. *Environmetrics*, 35(7):e2872, 2024. doi: 10.1002/env.2872.
- Russ Wolfinger, Randy Tobias, and John Sall. Computing Gaussian likelihoods and their derivatives for general linear mixed models. *SIAM Journal on Scientific Computing*, 15(6): 1294–1310, 1994. doi: 10.1137/0915079.
- Simon N. Wood. *Generalized Additive Models: An Introduction with R*. Chapman and Hall/CRC, 2nd edition, 2017. doi: 10.1201/9781315370279.
- Hao Zhang. On estimation and prediction for spatial generalized linear mixed models. *Biometrics*, 58(1):129–136, 2002. doi: 10.1111/j.0006-341X.2002.00129.x.
- Hao Zhang and Dale L. Zimmerman. Towards reconciling two asymptotic frameworks in spatial statistics. *Biometrika*, 92(4):921–936, 2005. doi: 10.1093/biomet/92.4.921.
- Dale L. Zimmerman and Jay M. Ver Hoef. *Spatial Linear Models for Environmental Data*. Chapman and Hall/CRC, 2024. doi: 10.1201/9781003044680.

7 Acknowledgements

The eReefs datasets (model simulations, satellite or in-situ observations) and software were produced as part of the eReefs project (ereefs.org.au), which is a collaboration between Australia’s national science agency CSIRO, the Australian Institute of Marine Science (AIMS) and the Queensland Government, with observations obtained through the Integrated Marine Observing System (IMOS) and the Great Barrier Reef Marine Park Authority (GBRMPA). eReefs is funded by the Australian Government’s Reef Trust.

#Appendix 1. Spatial Linear Models and Random Forest Models {.unnumbered} The spatially explicit statistical models (i.e., spatial models) we used to answer these research questions are related to commonly used linear and generalized linear models but incorporate *spatial dependence*. Spatial dependence measures how similar two sites are based on their distance. Formally, we can build spatial dependence directly into spatial models by leveraging Tobler’s First Law of Geography [Tobler, 1970], which states succinctly that “nearby sites tend to be more similar than distant sites”. Not only is this intuitive from a “first principles” perspective, but spatial models also provide many advantages for modeling spatial data compared to models that assume independence among observations (i.e., nonspatial models). First, nonspatial models tend to underestimate uncertainty, yielding covariate standard errors and p-values for covariates that are too small and corresponding confidence intervals that are too narrow. Second, spatial models improve prediction at unsampled locations by borrowing strength from nearby observations, yielding more precise predictions and corresponding standard errors that

vary by location. Third, spatial models decompose model error into separate spatial and nonspatial components, informing residual and resilience structure in ways that nonspatial models cannot. Fourth, spatial models can provide ecologically meaningful insight into the spatial dependence structure of the process, informing relationships among measured and unmeasured variables as well as future sampling plans. Zimmerman and Ver Hoef [2024] provide a thorough review of spatial models for ecological and environmental data, providing more detail and justification for the claims made above.

Spatial models also offer many benefits over commonly used machine learning algorithms like random forests [Breiman, 2001, Cutler et al., 2007, James et al., 2017, Kuhn and Silge, 2022]. First, traditional machine learning algorithms do not incorporate spatial dependence and, by consequence, often perform worse than spatial models at characterizing important covariates and predicting at unobserved locations [Fox et al., 2020]. Second, machine learning algorithms struggle to quantify uncertainty in a principled manner, something spatial models excel at. Third, covariate inference is more straightforward using spatial models, as they naturally provide p-values that enable structured testing of hypotheses. Fourth, spatial models provide a rich set of diagnostics like leverage and Cook’s distances [Montgomery et al., 2021], quantities not generally defined for machine learning algorithms. This is not to say that machine learning algorithms cannot be very useful in ecological data analyses, but rather that spatial models are sometimes a more effective tool for modeling spatial data.

Though there are many benefits of spatial models over machine learning algorithms, the reverse is also true. It is much simpler to capture complex nonlinearities using “out-of-the-box” machine learning algorithms, bypassing the tuning often required for spatial models that involve nonlinear components like splines. This can be especially useful for ecological data, which often exhibit nonlinear relationships [Fox et al., 2017]. Machine learning algorithms also typically rely on few, if any, distributional assumptions and can be more robust against unusual observations like outliers [Hastie et al., 2009]. Machine learning algorithms also tend to be much more computationally efficient than spatial models. Importantly, machine learning algorithms and spatial models can be complementary tools that help shed insight into complicated ecological processes (something we reiterate in the upcoming analyses). An exciting and developing area of statistical research involves incorporating spatial modeling strategies directly into machine learning algorithms [Saha et al., 2023, Heaton et al., 2025].

The first spatially explicit model we define is the spatial linear model. The spatial linear model is a *mixed-effects model* [Pinheiro and Bates, 2000, Bolker, 2015] written as

$$y_i = \beta_0 + \beta_1 x_{1,i} + \beta_2 x_{2,i} + \dots + \beta_p x_{p,i} + \tau_i + \epsilon_i, \quad (1)$$

where $i = 1, 2, \dots, n$ indexes the n observations. The quantity y_i is the response variable (for observation i), the β parameters are slope parameters controlling the covariates, $x_{j,i}$ (for $j = 1, 2, \dots, p$). The random error τ_i is a spatial random error, and the random error ϵ_i is the nonspatial (i.e., independent) random error. The spatial random error is spatially structured, capturing spatial residual effects like environmental gradients we do not have data to fully

represent. The nonspatial random error is not spatially structured and captures nonspatial residual effects like measurement error and microscale variation. The difference between a traditional (i.e., nonspatial) linear model and a spatial linear model is the inclusion of τ_i (in the spatial linear model). The spatial random error implied by τ_i is governed by a *spatial covariance function*. The spatial covariance function allows the model to incorporate spatial dependence, yielding the improved model performance previously described. Figure 12 shows three different spatial covariance functions that describe different ways the spatial covariance decays with distance. Model parameters and standard errors are estimated from data using likelihood-based [Patterson and Thompson, 1971, Harville, 1977, Wolfinger et al., 1994] or semivariogram-based [Cressie, 1985, Curriero and Lele, 1999] estimation methods. Often, the spatial linear model in Equation 1 is written more compactly using *matrix notation*:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\tau} + \boldsymbol{\epsilon}, \quad (2)$$

where $\mathbf{X}$ is a matrix that holds the j covariates for each of n observations, and $\mathbf{y}$, $\boldsymbol{\beta}$, $\boldsymbol{\tau}$, and $\boldsymbol{\epsilon}$ are vectors that contain each of the n or j quantities from Equation 1. The spatial linear model can be adjusted to account for nonspatial random effects just as one would add random effects to a nonspatial linear mixed model:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\mathbf{u} + \boldsymbol{\tau} + \boldsymbol{\epsilon}, \quad (3)$$

where $\mathbf{Z}$ is a random effect design matrix (dimension $n \times k$) that indexes each of the k random effects in $\mathbf{u}$. For more information about specific forms of various spatial covariance functions, see Schabenberger and Gotway [2017] and Zimmerman and Ver Hoef [2024].




Figure 12: Three spatial covariance functions that represent different spatial dependence behavior.

Like the traditional linear model, the spatial linear model can capture complex nonlinearities in the data through the use of interaction effects [Faraway, 2002, Lenth and Piaskowski, 2025], splines [Chambers and Hastie, 1992, Ruppert et al., 2003], additive models [Wood, 2017, Pedersen et al., 2019], and penalized regression models like the ridge and lasso [Tibshirani, 1996]. Generally, this is accomplished by leveraging *basis functions* that break down nonlinear behavior into a set of additive linear components that may include penalty terms which guard against overfitting. Furthermore, there is no assumption that y_i itself is normally distributed; rather, distributional assumptions are placed on the random errors after incorporating the slope parameters. This clarification implies the spatial (and nonspatial) linear models can be quite effective at modeling a range of data that are not normally distributed. For highly nonnormal data, tools like generalized linear models, which we discuss later, can be helpful. Finally, under general conditions, the slope parameters of the spatial (and nonspatial) linear models are approximately normally distributed with a large sample size (see Casella and Berger [2024] for a general justification via the Central Limit Theorem and Zhang and Zimmerman [2005] for nuances regarding spatial data). This is beneficial because it justifies valid hypothesis testing on the slope coefficients (often of primary ecological interest), no matter the distribution of the data (given certain assumptions are satisfied).

One of the most useful applications of the spatial linear model is predicting the response variable at unobserved locations. A very useful spatial prediction method called Kriging has a rich theoretical justification [Cressie, 1990, Stein, 1999]. Kriging is the spatial analogue to best linear unbiased prediction (BLUP) for mixed models [Henderson, 1975]. Via Kriging, the spatial linear model produces BLUPs at each unobserved location of interest, alongside prediction intervals to characterize uncertainty. While useful, sometimes the goal is to create a BLUP for an entire region of interest. This technique, which builds upon the theoretical foundations of Kriging, is called block Kriging [Ver Hoef, 2002, Cressie, 2006]. The intuition is to make point predictions at a very dense grid in the study region and then average these predictions, appropriately characterizing uncertainty. Together, Kriging and block Kriging provide a complete set of tools for spatial prediction, given the fit of a spatial linear model.

Spatial generalized linear models are an extension of spatial linear models to highly nonnormal data [Christensen and Waagepetersen, 2002, Zhang, 2002, Diggle and Ribeiro Jr, 2007, Bonat and Ribeiro Jr, 2016]. Ver Hoef et al. [2024] proposed a novel application of the Laplace approximation to enable restricted maximum likelihood estimation of spatial generalized linear models. Their model formulation builds upon Equation 1 and is given by

$$g(\mu_i) = \beta_0 + \beta_1 x_{1,i} + \beta_2 x_{2,i} + \dots + \beta_p x_{p,i} + \tau_i + \epsilon_i, \quad (4)$$

where $g(\mu_i)$ is a “link function” that links the distribution of y_i to a linear function of its mean, μ_i . Common link functions include the log link for count and skewed data and the logit link for binomial and proportion data (Table 9). Using the same argument as for spatial linear models, nonspatial random effects can be added to spatial generalized linear models.

Table 9: Common generalized linear model families, link functions, and appropriate data types.

Family	Link Function	Link Name	Data Type
Binomial	$f(\mu) = \log(\mu/(1 - \mu))$	Logit	Binary; Binary Count
Poisson	$f(\mu) = \log(\mu)$	Log	Count
Negative Binomial	$f(\mu) = \log(\mu)$	Log	Count
Beta	$f(\mu) = \log(\mu/(1 - \mu))$	Logit	Proportion
Gamma	$f(\mu) = \log(\mu)$	Log	Skewed
Inverse Gaussian	$f(\mu) = \log(\mu)$	Log	Skewed

Spatial models are complicated, both theoretically and computationally. This has contributed to their relative lack of use in the ecological community. However, recent advances in open-source software have made spatial models more accessible to ecologists who have strong subject matter expertise but may lack sufficient statistical background or computational skills to implement them from scratch. One such advancement is the `spmodel` **R** package [Dumelle et al., 2023], which emulates base-**R** functions like `lm()` and `glm()` to account for spatial dependence via `splm()` and `splgm()`, respectively. Familiar **R** functions like `summary()` and `predict()` can be used directly on models fit via `spmodel`. Importantly, `spmodel` also has tools for spatial random forest residual modeling [Fox et al., 2020], applications to large data of several thousand observations via spatial indexing [Ver Hoef et al., 2023], and cross validation [Stone, 1974, Roberts et al., 2017].

#Appendix 2 {.unnumbered}

This is an appendix