

SCHUR-WEYL DUALITY FOR THE BRAUER ALGEBRA AND THE ORTHO-SYMPLECTIC LIE SUPERALGEBRA

MICHAEL EHRIG AND CATHARINA STROPPEL

ABSTRACT. We give a proof of a Schur-Weyl duality statement between the Brauer algebra and the ortho-symplectic Lie superalgebra $\mathfrak{osp}(V)$ over the complex numbers, and for the Brauer algebra and classical orthogonal respectively symplectic Lie algebras in arbitrary characteristic.

1. INTRODUCTION

Fix as ground field the complex numbers \mathbb{C} . In this paper we describe the centralizer of the action of the ortho-symplectic Lie superalgebra $\mathfrak{osp}(V)$ on the tensor powers of its defining representation V . Here V is a vector superspace of superdimension $\text{sdim} V$ equal to $2m|2n$ or $2m+1|2n$ equipped with a supersymmetric bilinear form, and $\mathfrak{osp}(V)$ denotes the Lie superalgebra of all endomorphisms preserving this form. In particular, the extremal cases $n=0$ respectively $m=0$ give the classical orthogonal respectively symplectic simple Lie algebras. Our main result is the following generalization of Brauer's centralizer theorem in the classical case, see e.g. [GW], to a Lie superalgebra version.

Theorem A. *Let m and n be nonnegative integers and let V be as above. Let $\delta = 2m - 2n$ respectively $\delta = 2m + 1 - 2n$ be the supertrace of V . Assume that one of the following assumptions holds*

- $\text{sdim} V \neq 2m|0$ and $d \leq m + n$ or
- $\text{sdim} V = 2m|0$ with $m > 0$ and $d < m$.

Then there is a canonical isomorphism of algebras

$$\text{End}_{\mathfrak{osp}(V)}(V^{\otimes d}) \cong \text{Br}_d(\delta). \quad (1.1)$$

Here $\text{Br}_d(\delta)$ denotes the Brauer algebra on d strands with parameter δ which was originally introduced by Brauer in [Br], see Definition 3.6. Note that in case $m=0$ or $n=0$ we obtain Brauer's classical centralizer theorem explained in modern language for instance in [GW]. In contrast to these classical cases, the endomorphism algebras of finite dimensional representations of a Lie superalgebra are in general not semisimple. Hence, our theorem covers also the (most interesting) cases in which $\text{Br}_d(\delta)$ is not semisimple.

M.E. was financed by the DFG Priority program 1388, C.S. thanks the MPI in Bonn for excellent working conditions and financial support.

These non-semisimple Brauer algebras also appear as idempotent truncations of endomorphism rings of modules in category \mathcal{O} for the classical Lie algebras, see [ES1], [ES2]. In [ES1] it was shown that the Brauer algebra can be equipped with a nonnegative \mathbb{Z} -grading (which is even Koszul in case $\delta \neq 0$), see [ES1], [ES2], [Li]. As a consequence of our theorem, the endomorphism rings (1.1) inherit a grading. The existence of such a grading on Brauer algebras is rather unexpected and up to now this grading cannot be described intrinsically in terms of the representation theory of the Lie superalgebra.

The choice of \mathbb{C} as ground field allowed us not having to pass to a version of Kostant's \mathbb{Z} -form of the universal enveloping algebra for Lie superalgebras. Since we do not want to get into the technicalities of these \mathbb{Z} -forms here, we restrict ourselves for general ground fields to (ordinary) Lie algebras: Denote by \mathbb{k} an arbitrary field. Assume that either $m = 0$ or $n = 0$, i.e., $\mathfrak{osp}(V)$ is either the classical orthogonal or classical symplectic Lie algebra. In these cases denote by $\mathcal{U}_{\mathbb{Z}}(\mathfrak{osp}(V))$ Kostant's \mathbb{Z} -form of $\mathcal{U}(\mathfrak{osp}(V))$, see [H, Section 26], and by $\mathcal{U}_{\mathbb{k}} = \mathcal{U}_{\mathbb{Z}}(\mathfrak{osp}(V)) \otimes_{\mathbb{Z}} \mathbb{k}$ its specialization to \mathbb{k} . It acts on the \mathbb{k} -form $V_{\mathbb{k}}$, of the vector representation. Let $\mathrm{Br}_d(\delta)_{\mathbb{k}}$ denote the Brauer algebra defined over the field \mathbb{k} .

Theorem B. *Let \mathbb{k} be an arbitrary field. Assume $m = 0$ or $n = 0$ and let δ be the supertrace of $V_{\mathbb{k}}$. Assume $d < m$ in the even uthogonal case and $d \leq m + n$ otherwise, then there is a canonical isomorphism of algebras*

$$\mathrm{End}_{\mathcal{U}_{\mathbb{k}}}(V_{\mathbb{k}}^{\otimes d}) \cong \mathrm{Br}_d(\delta)_{\mathbb{k}}.$$

The existence of a ring homomorphism (1.1) from the Brauer algebra to the endomorphism ring of $V^{\otimes d}$ is not new. It is for instance a special case of the Brauer algebra actions defined in [BCHLLS] and also follows from the more abstract theory of Deligne categories, [De]. However, the isomorphism theorem is, as far as we know, new. The crucial point here is that in general (and in particular in contrast to the classical case), the tensor space is not completely reducible and then unfortunately the methods from [GW] and [BCHLLS] do not apply. As far as we know there is not much known about the indecomposable summands appearing in this tensor space and even less is known about the general structure of the category of finite dimensional representations of $\mathfrak{osp}(V)$, see [GS], [Sea], and also [Co] for some partial results. This is very much in contrast to the case of the general Lie superalgebra $\mathfrak{gl}(m|n)$, where the corresponding (mixed) tensor spaces were recently studied and described in detail, [CW], [BS2]. There, the role of the Brauer algebra is played by the *walled* Brauer algebra respectively *walled* Brauer category, sometimes also called *oriented* Brauer category, see [BS2], [BCNR], [SS]. As our main tool we construct an embedding of the Brauer algebra into an additive closure of the *walled* Brauer category and then use the above mentioned results for the general linear case to deduce Theorem A in the ortho-symplectic case.

More precisely we restrict the action of $\mathfrak{osp}(V)$ to a suitably embedded $\mathfrak{gl}(m|n)$ and decompose the occurring representation $V^{\otimes d}$ as a direct sum of mixed tensor products on which the walled Brauer category acts naturally. One of the crucial steps is the following generalization from [BS2]:

Theorem C. *Let $m, n \in \mathbb{Z}_{\geq 0}$, $d > 0$. The corresponding oriented Brauer category $\mathcal{OB}_d(m-n)$ acts on $V^{\otimes d}$ and its action commutes with the action of $\mathfrak{gl}(m|n)$.*

We then embed the Brauer algebra into the endomorphism ring of $V^{\otimes d}$ viewed as an object inside the additive closure of the oriented Brauer category. The injectivity of Theorem A is then deduced from the corresponding injectivity result [BS2, Theorem 7.1] for walled Brauer algebras. Finally we show (by very elementary arguments) that any $\mathfrak{gl}(m|n)$ endomorphism that commutes also with the action of $\mathfrak{osp}(V)$ is already contained in the image of the embedding and gives the desired isomorphism.

Similar results in the spirit of Schur-Weyl duality for the ortho-symplectic supergroup were independently also obtained by Lehrer and Zhang in [LZ1] and [LZ2] using methods from graded-commutative algebraic geometry. Although their results go further, since they for instance also describe the kernel of the action for arbitrary values of d, m, n , our proof of the isomorphism theorem is considerably more elementary. The proof of the main theorem only involves methods from (super) linear algebra and the (elementary) theorem of mixed Schur-Weyl duality from [BS2]. Using however the non-elementary results of [BS2], it is possible to describe also the kernel of our action in general, but it requires to work with a graded version of the Brauer algebra as defined in [ES1] and will appear in a separate article where this graded Brauer algebra is studied. We expect that our approach generalizes to the quantized (super) case using the quantised walled Brauer algebras, see [DDS], in a straightforward manner.

Acknowledgement We would like to thank Gus Lehrer for many useful comments as well as pointing out an error in a previous version of this paper. Furthermore we would like to thank Kevin Coulembier and Daniel Tubbenhauer for various comments on a preliminary version.

2. THE LIE SUPERALGEBRA

We fix as ground field the complex numbers \mathbb{C} . By a superspace we mean a $\mathbb{Z}/2\mathbb{Z}$ -graded vector space $V = V_0 \oplus V_1$; for any homogeneous element $v \in V$ we denote by $|v| \in \{0, 1\}$ its parity. The integer $\dim V_0 - \dim V_1$ is called the *supertrace* of V and we denote by $\text{sdim } V = \dim V_0 | \dim V_1$ its *superdimension*. Given a superspace V let $\mathfrak{gl}(V)$ be the corresponding general Lie superalgebra, i.e. the superspace $\text{End}_{\mathbb{C}}(V)$ of all endomorphism with the superbracket

$$[X, Y] = X \circ Y - (-1)^{|X||Y|} Y \circ X.$$

For the whole paper we fix now $n \in \mathbb{Z}_{\geq 0}$ and a finite dimensional super-space $V = V_0 \oplus V_1$ with $\dim V_1 = 2n$, and equipped with a non-degenerate supersymmetric bilinear form $\langle -, - \rangle$, i.e. a bilinear form $V \times V \rightarrow \mathbb{C}$ which is symmetric when restricted to $V_0 \times V_0$, skew-symmetric on $V_1 \times V_1$ and zero on mixed products. It will sometimes be convenient to work with a fixed homogeneous basis $v_i, i \in I$ of V , for a suitable indexing set I , and right dual basis v_i^* , i.e. $\langle v_i, v_j^* \rangle = \delta_{i,j}$. Attached to this data we have the following:

Definition 2.4. *The ortho-symplectic Lie superalgebra $\mathfrak{osp}(V)$ is the Lie supersubalgebra of $\mathfrak{gl}(V)$ consisting of all endomorphisms which respect the supersymmetric bilinear form. Explicitly, a homogeneous element $X \in \mathfrak{osp}(V)$ has to satisfy*

$$\langle Xv, w \rangle + (-1)^{|X||v|} \langle v, Xw \rangle = 0, \quad (2.2)$$

for homogeneous $v \in V$.

From now on we make the convention that, whenever the parity of an element appears in a formula, the element is assumed to be homogeneous.

Remark 2.5. If $X \in \mathfrak{osp}(V)$, then $\langle Xv, w \rangle \neq 0$ implies $|X| = |v| + |w|$.

3. THE BRAUER ALGEBRA

The following algebra was originally defined by Brauer [Br] in his study of the orthogonal group.

Definition 3.6. *Let $d \in \mathbb{Z}_{\geq 0}$ and $\delta \in \mathbb{C}$. The Brauer algebra $\text{Br}_d(\delta)$ is the associative unital \mathbb{C} -algebra generated by the elements*

$$s_i, e_i \text{ for } 1 \leq i \leq d-1,$$

subject to the relations

$$\begin{aligned} s_i^2 &= 1, & s_i s_j &= s_j s_i, & s_k s_{k+1} s_k &= s_{k+1} s_k s_{k+1}, \\ e_i^2 &= \delta e_i, & e_i e_j &= e_j e_i, & e_k e_{k+1} e_k &= e_{k+1}, & e_{k+1} e_k e_{k+1} &= e_k, \\ s_i e_i &= e_i = e_i s_i, & s_k e_{k+1} e_k &= s_{k+1} e_k, & s_{k+1} e_k e_{k+1} &= s_k e_{k+1}, \end{aligned} \quad (3.3)$$

for $1 \leq i, j \leq d-1$, $|i-j| > 1$, and $1 \leq k \leq d-2$.

Following Brauer we realize $\text{Br}_d(\delta)$ as a diagram algebra: A *Brauer diagram* on $2d$ vertices is a partitioning b of the set $\{1, 2, \dots, d, 1^*, 2^*, \dots, d^*\}$ into d subsets of cardinality 2. Let $\mathcal{B}[d]$ be the set of such Brauer diagrams. An element can be displayed graphically by arranging $2d$ vertices in two rows $1, 2, \dots, d$ and $1^*, 2^*, \dots, d^*$, with each vertex linked to precisely one other vertex. Two such diagrams are considered to be the same or equivalent if they link the same d pairs of points.

Special Brauer diagrams are the "unit" $1 = \{\{1, 1^*\}, \{2, 2^*\}, \dots, \{d, d^*\}\}$ connecting always j with j^* for all $1 \leq j \leq d$, and for $1 \leq i \leq d-1$ the s_i (respectively e_i) which connects j with j^* except of the new pairs $\{i, (i +$

$1)^*\}, \{i+1, i^*\}$ (respectively $\{i, i+1\}, \{i^*, (i+1)^*\}$) involving the numbers i and $i+1$;

$$s_i \mid \cdots \mid \overset{i \quad i+1}{\times} \mid \cdots \mid e_i \mid \cdots \mid \overset{i \quad i+1}{\cup} \mid \cdots \mid \quad (3.4)$$

Given two Brauer diagrams b and b' , their concatenation $b' \circ b$ is obtained by putting b on top of b' identifying vertex i^* in b with vertex i in b' and removing all the internal loops. Let $c(b, b')$ be the number of loops removed. Then we have the following fact, see e.g. [GW, Section 9 and 10] for details:

Lemma 3.7. *The Brauer algebra $\text{Br}_d(\delta)$ is, via the assignment (3.4) on generators, canonically isomorphic to the \mathbb{C} -algebra with basis $\mathcal{B}[d]$ and multiplication $bb' = \delta^{c(b, b')} b \circ b'$ for $b, b' \in \mathcal{B}[d]$.*

Remark 3.8. Generically, the algebra $\text{Br}_d(\delta)$ is semi-simple, but not semi-simple for specific integral values for δ (depending on d), see e.g. [ES1, 2.2] for a detailed description of the structure of these algebras.

We now define the two actions on the tensor space $V^{\otimes d}$ from Theorem A. The action of $\mathfrak{osp}(V)$ on $V^{\otimes d}$ is given by the comultiplication

$$\Delta(X) = X \otimes 1 + 1 \otimes X$$

for $X \in \mathfrak{osp}(V)$, keeping in mind the tensor product rule

$$(X \otimes Y)(v \otimes w) = (-1)^{|Y| \cdot |v|} (Xv \otimes Yw).$$

Explicitly we have for homogeneous elements $w_i \in V$

$$X \cdot (w_1 \otimes \dots \otimes w_r) = \sum_{i=1}^r (-1)^{(|w_1| + \dots + |w_{i-1}|) |X|} w_1 \otimes \dots \otimes w_{i-1} \otimes Xw_i \otimes w_{i+1} \otimes \dots \otimes w_r.$$

Definition 3.9. *Define the following linear endomorphisms*

$$\begin{aligned} \sigma : V \otimes V &\longrightarrow V \otimes V, \text{ via } \sigma(v \otimes w) := (-1)^{|v| \cdot |w|} w \otimes v \text{ and} \\ \tau : V \otimes V &\longrightarrow V \otimes V, \text{ via } \tau(v \otimes w) := \langle v, w \rangle \sum_{i \in I} (-1)^{|v_i|} v_i \otimes v_i^*. \end{aligned}$$

of $V \otimes V$ and for each fixed natural number $d \geq 2$ the endomorphisms

$$s_i = \text{id}^{\otimes(i-1)} \otimes \sigma \otimes \text{id}^{\otimes(d-i-1)} \quad \text{and} \quad e_i = \text{id}^{\otimes(i-1)} \otimes \tau \otimes \text{id}^{\otimes(d-i-1)}$$

of $V^{\otimes d}$ for $1 \leq i < d$.

The following proposition is straight-forward to check.

Proposition 3.10. *The maps σ, τ and s_i, e_i are $\mathfrak{osp}(V)$ -equivariant.*

Theorem 3.11. *Let $\delta = \dim V_0 - \dim V_1$ be the supertrace of V . Then there is a right action of $\text{Br}_d(\delta)$ on $V^{\otimes d}$ given by*

$$v \cdot s_i := s_i(v) \quad \text{and} \quad v \cdot e_i = e_i(v)$$

for $v \in V^{\otimes d}$ and $1 \leq i \leq d-1$. It commutes with the action of $\mathfrak{osp}(V)$.

Remark 3.12. Restricting only to the s_i gives us an action of the symmetric group S_d on $V^{\otimes d}$, which clearly commutes with the action of $\mathfrak{gl}(V)$. This is the super Schur-Weyl duality, [BS2, Theorem 7.5], originally proved by Sergeev [Sev] and Berele-Regev, [BR].

Proof of Theorem 3.11. By Proposition 3.10 all involved morphisms commute with the action of $\mathfrak{osp}(V)$. Hence it remains to show that it defines an action of the Brauer algebra. It is obvious that the s_i define an action of the symmetric group and that the relations $e_i e_j = e_j e_i$ are satisfied for $|i - j| > 1$. Moreover, $e_i^2 = \delta e_i$ by definition. To verify the relation $s_i e_i = e_i$ it suffices to assume $d = 2$. Let $v, w \in V$ be homogeneous then

$$\begin{aligned} (v \otimes w).s_i e_i &= (-1)^{|v| \cdot |w|} (w \otimes v).e_i = (-1)^{|v| \cdot |w|} \langle w, v \rangle \sum_{l \in I} (-1)^{|v_l|} v_l \otimes v_l^* \\ &= \langle v, w \rangle \sum_{l \in I} (-1)^{|v_l|} v_l \otimes v_l^* = (v \otimes w).e_i. \end{aligned}$$

For the equality $e_i = e_i s_i$ we note that $(v_j \otimes v_k^*).e_i = 0$ unless $j = k$, so we assume equality and obtain

$$\begin{aligned} (v_j \otimes v_j^*).e_i s_i &= \sum_{l \in I} (-1)^{|v_l|} (v_l \otimes v_l^*).s_i = \sum_{l \in I} (-1)^{|v_l|} (-1)^{|v_l|} v_l^* \otimes v_l \\ &= \sum_{l \in I} (-1)^{|v_l|} ((-1)^{|v_l|} v_l^*) \otimes v_l = \sum_{l \in I} (-1)^{|v_l|} v_l \otimes v_l^*, \end{aligned}$$

where the final equality holds because the $\{(-1)^{|v_l|} v_l^*\}$ also form a basis of V with right dual basis $\{v_l\}$.

For the final relations

$$e_i e_{i+1} e_i(x) = e_i(x) \quad \text{and} \quad e_{i+1} e_i e_{i+1}(x) = e_{i+1}(x), \quad (3.5)$$

$$e_i e_{i+1} s_i(x) = e_i s_{i+1}(x) \quad \text{and} \quad e_{i+1} e_i s_{i+1}(x) = e_{i+1} s_i(x) \quad (3.6)$$

for $x \in V^{\otimes d}$, it suffices to consider the case $d = 3$ and $x = u \otimes v \otimes w$ for homogeneous $u, v, w \in V$. The exact calculation is straight-forward and left to the reader. \square

4. THE ORIENTED BRAUER CATEGORY

We now recall the oriented Brauer category (or sometimes called walled Brauer category, see e.g. [SS]), which has as objects certain orientation sequences and morphism spaces given by oriented generalized Brauer diagrams. We then relate it to the space $V^{\otimes d}$.

We set $\text{OSeq}[d] = \{\wedge, \vee\}^{\times d}$ and $\widehat{\text{OSeq}}[d] = \{\wedge, \vee, \circ\}^{\times d}$ and call its elements *orientations* of length d respectively *generalized orientations* of length d . Let $\widehat{\mathcal{B}}[d]$ be the set of *generalized Brauer diagrams* on $2d$ vertices, that means diagrams which are Brauer diagrams, except that the partitioning is into subsets of cardinality 1 or 2. In other words, we allow vertices that are not connected to any other vertex.

Definition 4.13. An oriented generalized Brauer diagram is a triple $(\mathbf{t}, b, \mathbf{s})$ where $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}[d]$ and $b \in \widehat{\mathcal{B}}[d]$ such that the following conditions hold:


- (1) if $\{i, j\} \in b$ with $i \neq j$ then $\{\mathbf{t}_i, \mathbf{t}_j\} = \{\wedge, \vee\}$,
- (2) if $\{i^*, j^*\} \in b$ with $i \neq j$ then $\{\mathbf{s}_i, \mathbf{s}_j\} = \{\wedge, \vee\}$,
- (3) if $\{i, j^*\} \in b$ then $\{\mathbf{t}_i, \mathbf{s}_j\} \in \{\{\wedge\}, \{\vee\}\}$, and
- (4) if $\{i\} \in b$ or $\{i^*\} \in b$, then $\mathbf{t}_i = \circ$, respectively $\mathbf{s}_i = \circ$.

The pair (\mathbf{s}, \mathbf{t}) is then called the orientation.

We denote the set of diagrams oriented with orientation (\mathbf{s}, \mathbf{t}) by $\widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{t}}$ or by $\widehat{\mathcal{B}}[d]_{\mathbf{s}}^{\mathbf{t}}$ if we want to stress that the length is d , and draw them as oriented diagrams (in the obvious sense) with the orientation sequence \mathbf{s} at the bottom and the orientation sequence \mathbf{t} on the top.

It will be sometimes helpful to vary d and so we set $\widehat{\text{OSeq}} = \bigcup_{d' \geq 0} \widehat{\text{OSeq}}[d']$ and $\text{OSeq} = \bigcup_{d' \geq 0} \text{OSeq}[d']$. For $\mathbf{s} \in \widehat{\text{OSeq}}$ we denote by $\bar{\mathbf{s}} \in \text{OSeq} \subset \widehat{\text{OSeq}}$ the associated *reduced sequence* obtained by deleting all \circ 's. Similarly define \bar{b} for $b \in \widehat{\mathcal{B}}[d]$. Note that if $b \in \widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{t}}$ then $\bar{b} \in \widehat{\mathcal{B}}_{\bar{\mathbf{s}}}^{\bar{\mathbf{t}}}$. Moreover, we can allow pairs of sequences (\mathbf{s}, \mathbf{t}) not of the same length and define the sets $\widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{t}}$ and $\widehat{\mathcal{B}}_{\bar{\mathbf{s}}}^{\bar{\mathbf{t}}}$ of (reduced) oriented generalized Brauer diagrams with orientation (\mathbf{s}, \mathbf{t}) exactly as in Definition 4.13.

Example 4.14. The first diagram below is an element in $\widehat{\mathcal{B}}[4]_{(\vee, \wedge, \vee, \circ)}^{(\vee, \wedge, \vee, \circ)}$. The other two diagrams are not oriented, but removing the red part would give an element in $\widehat{\mathcal{B}}_{(\circ, \wedge, \vee)}^{(\circ)}$ and $\widehat{\mathcal{B}}_{(\vee, \wedge, \vee)}^{(\vee, \wedge, \vee, \circ)}$ respectively.



Definition 4.15. Let $d \in \mathbb{Z}_{\geq 0}$ and $\delta \in \mathbb{C}$. We define the oriented Brauer category $\mathcal{OB}_d(m-n)$ as the following \mathbb{C} -linear category: The set of objects is $\widehat{\text{OSeq}}[d]$, the morphism space $\text{Hom}(\mathbf{s}, \mathbf{t})$ for $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}[d]$ is the vector space with basis $\widehat{\mathcal{B}}[d]_{\mathbf{s}}^{\mathbf{t}}$, and the composition of morphisms

$$\text{Hom}(\mathbf{s}, \mathbf{t}) \times \text{Hom}(\mathbf{r}, \mathbf{s}) \longrightarrow \text{Hom}(\mathbf{r}, \mathbf{t}),$$

is done on basis vectors by putting the two diagrams on top of each other, glueing along the entries of \mathbf{s} and eliminating all internal \circ 's and also inner circles, each elimination of an internal circle resulting in multiplying the diagram with a factor of $m-n$. Similarly, define the category $\mathcal{OB}(m-n)$ by allowing as objects sequences of arbitrary finite length.

Lemma 4.16. Let $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}[d]$ and $\ell = |\{i \mid \mathbf{s}_i = \circ\}| - |\{i \mid \mathbf{t}_i = \circ\}|$. Then $\text{Hom}(\mathbf{s}, \mathbf{t}) = \{0\}$ if ℓ is odd or ℓ is even and additionally it holds

$$\frac{\ell}{2} \neq |\{i \mid \mathbf{s}_i = \wedge\}| - |\{i \mid \mathbf{t}_i = \wedge\}|.$$

Proof. This follows directly from the definitions. \square

Remark 4.17. We like to stress that our category $\mathcal{OB}(m-n)$ is slightly different from the category called oriented walled Brauer category in [BCNR] in the sense that we have more objects, since we also fix the places of the trivial tensor factors. Hence, the oriented Brauer category from [BCNR] is the full subcategory of ours, where we only allow objects without \circ 's.

To relate this category to the endomorphisms of the superspace $V^{\otimes d}$ let m be such that $\dim(V_0) = 2m$ or $\dim(V_0) = 2m+1$. Unfortunately we have to distinguish between these two cases. Therefore, we will refer to them as the *even case* respectively *odd case*. Fix the sets

$$I^\wedge = \{i \mid 1 \leq i \leq m+n\}, I^\vee = \{\bar{i} \mid 1 \leq i \leq m+n\}, I^\circ = \{0\},$$

and also $I = I^\wedge \cup I^\vee$ in the even case and $I = I^\wedge \cup I^\vee \cup I^\circ$ in the odd case. Denote by $\|i\|$ the absolute value for an element in $i \in I$, in particular $\|\bar{i}\| = i = \|i\|$ for $\bar{i} \in I^\vee$. To simplify calculations later on we use an explicit realization of $\mathfrak{osp}(V)$ in terms of matrices, see e.g. [Mu].

Definition 4.18. Let $V = \mathbb{C}^{2m+1|2n}$ be the superspace with basis v_i for $i \in I$ and $|v_i| = 0$ if $\|i\| \leq m$ and $|v_i| = 1$ otherwise. To write down matrices we order the basis elements as follows

$$0 < \bar{1} < \dots < \bar{m} < 1 < \dots < m < \overline{m+1} < \dots < \overline{m+n} < m+1 < \dots < m+n.$$

We denote by $\mathfrak{osp}(2m+1|2n) \subset \mathfrak{gl}(V)$ the Lie supersubalgebra given by

$$\left\{ \left(\begin{array}{ccc|cc} 0 & -u_1^T & -u_2^T & x_1 & x_2 \\ u_2 & -A^T & a_1 & C^T & z_1 \\ u_1 & a_2 & A & z_2 & B \\ \hline -x_2^T & -B^T & -z_1^T & -D^T & d_1 \\ x_1^T & z_2^T & C & d_2 & D \end{array} \right) \mid \begin{array}{l} A, a_1, a_2 \text{ are } m \times m \text{ matrices} \\ D, d_1, d_2 \text{ are } n \times n \text{ matrices} \\ u_1, u_2 \text{ are } m \times 1 \text{ matrices} \\ x_1, x_2 \text{ are } 1 \times n \text{ matrices} \\ B, z_1, z_2 \text{ are } m \times n \text{ matrices} \\ C \text{ is a } n \times m \text{ matrix} \\ a_1, a_2 \text{ are skew-symmetric} \\ d_1, d_2 \text{ are symmetric} \end{array} \right\}.$$

We denote by $\mathfrak{osp}(2m|2n)$ the Lie superalgebra produced in the same way but using the superspace $\mathbb{C}^{2m|2n}$ obtained by omitting the basis vector v_0 instead of V , and then using the matrices obtained by deleting the first row and column in the description of $\mathfrak{osp}(2m+1|2n)$.

Definition 4.18 is in the odd case consistent with the previous definition of $\mathfrak{osp}(V)$ if we take the bilinear form $\langle -, - \rangle$ given by the following block matrix where $\mathbb{1}_k$ denotes the identity matrix of size $k \times k$:

$$J = \left(\begin{array}{ccc|cc} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \mathbb{1}_m & 0 & 0 \\ 0 & \mathbb{1}_m & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & -\mathbb{1}_n \\ 0 & 0 & 0 & \mathbb{1}_n & 0 \end{array} \right),$$

In the even case one has to delete the first row and column.

To relate the action of $\mathfrak{osp}(2m+1|2n)$ on $V^{\otimes d}$ to the oriented Brauer category we will restrict the action to the general linear Lie supersubalgebra $\mathfrak{gl}(m|n)$. The embedding $\iota : \mathfrak{gl}(m|n) \hookrightarrow \mathfrak{osp}(2m+1|2n)$ is obtained by only allowing non-zero entries in the matrices A , B , C , and D in the notations above; similarly in the even case.

Then the restriction of V is given by the followings lemma.

Lemma 4.19. *As a $\mathfrak{gl}(m|n)$ module $V = W_\wedge \oplus W_\vee \oplus W_\circ$ in the odd case and $V = W_\wedge \oplus W_\vee$ in the even case, with*

$$W_\wedge = \text{span}\{v_i \mid i \in I^\wedge\}, W_\vee = \text{span}\{v_i \mid i \in I^\vee\}, \text{ and } W_\circ = \text{span}\{v_0\}.$$

Moreover, W_\wedge is isomorphic to the natural representation of $\mathfrak{gl}(m|n)$, W_\vee to its dual and W_\circ to the trivial representation.

Proof. Clearly, W_\wedge is isomorphic to the natural representation of $\mathfrak{gl}(m|n)$. The dual of the natural representation can be described in an appropriate basis w_i by $E_{i,j}w_k = -\delta_{i,k}(-1)^{(|i|+|j|)|i|}w_j$, and so it is isomorphic to W_\vee via $w_i \mapsto v_{\bar{i}}$. The statement for W_\circ is evident. \square

More generally we obtain the following as a direct consequence:

Corollary 4.20. *The space $V^{\otimes d}$ decomposes as a $\mathfrak{gl}(m|n)$ -module as*

$$V^{\otimes d} \cong \bigoplus_{\mathbf{s} \in \widehat{\text{OSeq}}[d]} W_{\mathbf{s}_1} \otimes \dots \otimes W_{\mathbf{s}_d}. \quad (4.7)$$

We abbreviate $W_{\mathbf{s}} = W_{\mathbf{s}_1} \otimes \dots \otimes W_{\mathbf{s}_d}$ and define for each $\mathbf{s} \in \widehat{\text{OSeq}}[d]$ in the odd case (respectively each $\mathbf{s} \in \text{OSeq}[d]$ in the even case) the set of \mathbf{s} -indices

$$\text{Vect}(\mathbf{s}) = \{\underline{i} = (i_1, \dots, i_d) \mid i_j \in I^{\mathbf{s}_j}\}.$$

Obviously this set labels the standard basis vectors in the summand corresponding to \mathbf{s} in the decomposition (4.7) with $v_{\underline{i}} = v_{i_1} \otimes \dots \otimes v_{i_d}$.

Lemma 4.21. *Let $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}[d]$ or $\mathbf{s}, \mathbf{t} \in \text{OSeq}[d]$ be two sequences, such that there exists $\sigma \in S_d$ with $\mathbf{s}_i = \mathbf{t}_{\sigma(i)}$. Then σ induces an isomorphism*

$$\psi_\sigma : W_{\mathbf{s}} \longrightarrow W_{\mathbf{t}}, \quad (4.8)$$

of $\mathfrak{gl}(m|n)$ -modules via the action of S_d given in Remark 3.12. Choosing σ to be of minimal length defines a distinguished isomorphism ψ_σ .

Proof. The action of the symmetric group on $V^{\otimes d}$ even commutes with $\mathfrak{gl}(V)$ and hence gives a morphism, and then clearly an isomorphism, of $\mathfrak{gl}(m|n)$ -modules when restricted to the subspaces. \square

As a consequence we have canonical isomorphisms of algebras

$$\text{End}_{\mathfrak{gl}(m|n)}(W_{\mathbf{s}}) \cong \text{End}_{\mathfrak{gl}(m|n)}(W_{\mathbf{t}})$$

for each pair \mathbf{s}, \mathbf{t} as above, namely given by conjugation with ψ_σ , where σ is the unique choice of minimal length. Note that these endomorphism rings

were described in detail in [BS2]. In case V has large enough dimension (in the sense that $d < (m+1)(n+1)$) and $\mathbf{t} = (\wedge, \dots, \wedge, \vee, \dots, \vee, \circ, \dots, \circ)$, where the symbols \wedge and \vee appear exactly r respectively s times, this endomorphism ring is exactly the *walled Brauer algebra* from [BS2, Theorem 7.8] originally introduced [Tu], [KM] and studied in [Ni].

Lemma 4.22. *Let $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}[d]$ and $\ell = |\{i \mid \mathbf{s}_i = \circ\}| - |\{i \mid \mathbf{t}_i = \circ\}|$. Then $\text{Hom}_{\mathfrak{gl}(m|n)}(W_{\mathbf{s}}, W_{\mathbf{t}}) = \{0\}$ if ℓ is odd or ℓ is even and then additionally*

$$\frac{\ell}{2} \neq |\{i \mid \mathbf{s}_i = \wedge\}| - |\{i \mid \mathbf{t}_i = \wedge\}|. \quad (4.9)$$

Proof. Let $\text{ht}(\lambda)$ denote the height of a $\mathfrak{gl}(m|n)$ weight, i.e. the sum of all coefficients written with respect to the standard $\{\varepsilon_i\}$ -basis. In this basis the weights occurring in W_{\wedge} are $\varepsilon_1, \dots, \varepsilon_{m+n}$, those in W_{\vee} are $-\varepsilon_1, \dots, -\varepsilon_{m+n}$, and in W_{\circ} only the weight 0 occurs. If ℓ is odd, then the height of the weights occurring in $W_{\mathbf{s}}$ will either all be even or all be odd, while in $W_{\mathbf{t}}$ it will be the other way around, thus there can't be a non-trivial $\mathfrak{gl}(m|n)$ -morphism.

If ℓ is even and (4.9) holds define $\mathbf{s}_{\wedge} = |\{i \mid \mathbf{s}_i = \wedge\}|$ and analogously \mathbf{s}_{\vee} , \mathbf{t}_{\wedge} , and \mathbf{t}_{\vee} . Assume now $\mathbf{s}_{\wedge} - \mathbf{t}_{\wedge} > \ell/2$, then it follows that $\mathbf{s}_{\vee} - \mathbf{t}_{\vee} < \ell/2$. By adjointness and Lemma 4.21 we know that

$$\text{Hom}_{\mathfrak{gl}(m|n)}(W_{\mathbf{s}}, W_{\mathbf{t}}) \cong \text{Hom}_{\mathfrak{gl}(m|n)}(W_{\wedge}^{\otimes(\mathbf{s}_{\wedge} + \mathbf{t}_{\vee})}, W_{\wedge}^{\otimes(\mathbf{t}_{\wedge} + \mathbf{s}_{\vee})}).$$

Then $\mathbf{s}_{\wedge} + \mathbf{t}_{\vee} < \mathbf{t}_{\wedge} + \mathbf{t}_{\vee} - \ell/2 < \mathbf{t}_{\wedge} + \mathbf{s}_{\vee}$, and so the height of any weight appearing in the domain is strictly less than the height of those appearing in the codomain and so all $\mathfrak{gl}(m|n)$ morphism are trivial. The remaining case $\mathbf{s}_{\wedge} - \mathbf{t}_{\wedge} < \ell/2$ is done in exactly the same way. \square

To extend the action from Theorem 3.11 to the oriented Brauer category we need the notion of a weight of an oriented Brauer diagram:

Definition 4.23. *Let $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}$. Assume $b \in \widehat{\mathcal{B}}[d]_{\mathbf{s}}^{\mathbf{t}}$ and $\underline{i} \in \text{Vect}(\mathbf{s})$ and $\underline{j} \in \text{Vect}(\mathbf{t})$. We denote by $b_{\underline{i}}^{\underline{j}}$ the diagram b with the vertices at the bottom labelled by the elements i_1, \dots, i_d and the vertices on the top by j_1, \dots, j_d (read always from left to right). We say that $b_{\underline{i}}^{\underline{j}}$ is consistently labelled if the labelling function $\| L(-) \|$ defined by $L(k) = j_k$ and $L(k^*) = i_k$, for $1 \leq k \leq d$, is constant on the subsets of the partition b .*

Example 4.24. *The first of the following diagrams is consistently labelled, the second and third are not:*

$$\begin{array}{ccc} \begin{array}{c} \overline{1} \quad 1 \quad \overline{2} \quad 0 \\ \swarrow \quad \searrow \quad \swarrow \quad \searrow \\ \overline{2} \quad 0 \quad 3 \quad \overline{3} \end{array} & \begin{array}{c} \overline{1} \quad 1 \quad \overline{3} \quad 0 \\ \swarrow \quad \searrow \quad \swarrow \quad \searrow \\ \overline{2} \quad 0 \quad 3 \quad \overline{3} \end{array} & \begin{array}{c} \overline{1} \quad 1 \quad \overline{2} \quad 0 \\ \swarrow \quad \searrow \quad \swarrow \quad \searrow \\ \overline{2} \quad \overline{2} \quad 3 \quad \overline{3} \end{array} \end{array} \quad (4.10)$$

Definition 4.25. The weight $\text{wt}\left(b_{\underline{i}}^j\right)$ of a labelled oriented diagram $b_{\underline{i}}^j$ is defined as follows:

$$\text{wt}\left(b_{\underline{i}}^j\right) := \begin{cases} \prod_c (-1)^{|c|} \prod_h (-1) \prod_{h'} (-1), & \text{if } b_{\underline{i}}^j \text{ is consistently labelled,} \\ 0, & \text{otherwise,} \end{cases},$$

where, viewed as diagrams,

- (1) the first product is over all crossings c of two different strands with the notation $|c| = |i| \cdot |j|$ if the two strands are labelled by i and j ;
- (2) the second product is over all labelled clockwise caps h with large labels, i.e. oriented horizontal strands at the bottom of the diagram with left endpoint oriented \wedge and with labelling set $\{a, \bar{a}\}$ such that $\|a\| > m$;
- (3) the third product is over all labelled anticlockwise cups h' with large labels, i.e. oriented horizontal strands at the top of the diagram with left endpoint oriented \vee and with labelling set $\{a, \bar{a}\}$ such that $\|a\| > m$.

Remark 4.26. Note that the weight of a diagram depends on V . For instance the first diagram in (4.10) has weight 0 if $m \geq 3$, has weight -1 if $m = 1, 2$ and has weight 1 if $m = 0$. Note that the weight of a crossing is by definition independent of the four possible orientations:

$$t_1 = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \quad t_2 = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \quad t_3 = \begin{array}{c} \diagdown \quad \diagup \\ \diagdown \quad \diagup \end{array} \quad t_4 = \begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array} \quad (4.11)$$

On the other hand, we can “rotate” each crossing by applying a cup and a cap on top of the diagram and compute then its weight. But in any case the weights of the added cup and cap multiply to 1, and so the weights don’t change. In fact, we will see in the proof of Theorem 4.28 that the weight is an invariant on equivalence classes of oriented labelled diagrams.

By a *representation* of $\mathcal{OB}_d(m-n)$ we mean a linear functor F from $\mathcal{OB}_d(m-n)$ to the category of complex vector spaces. Equivalently we say $\mathcal{OB}_d(m-n)$ acts on $X := \bigoplus_{\widehat{\text{OSeq}}[d]} F(\mathbf{s})$. If X is moreover an R -module for some ring R , then the action of $\mathcal{OB}_d(m-n)$ commutes with the action of R if $F(f)(r.x) = r.F(f)(x)$ for any $x \in X, r \in R$ and morphism f in $\mathcal{OB}_d(m-n)$.

Definition 4.27. For $\mathbf{s}, \mathbf{t} \in \widehat{\text{OSeq}}$ and $b \in \widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{t}}$ we define the linear map

$$F(b): W_{\mathbf{s}} \longrightarrow W_{\mathbf{t}}, \text{ via } F(b)(v_{\underline{i}}) := \sum_{\underline{j} \in \text{Vect}(\mathbf{t})} \text{wt}\left(b_{\underline{i}}^j\right) v_{\underline{j}}.$$

on basis vectors $v_{\underline{i}}$, where $\underline{i} \in \text{Vect}(\mathbf{s})$.

Theorem 4.28. Let δ be the supertrace of V . The assignment

$$\mathbf{s} \mapsto F(\mathbf{s}) := W_{\mathbf{s}} \quad \text{and} \quad f \mapsto F(f),$$

on objects respectively morphisms defines a functor from $\mathcal{OB}(m-n)$ to the category of finite dimensional $\mathfrak{gl}(m|n)$ -modules.

It restricts to an action of the oriented Brauer category $\mathcal{OB}_d(m-n)$ on $V^{\otimes d}$ which commutes with the action of $\mathfrak{gl}(m|n)$. The identity of the object $\mathbf{s} \in \widehat{\text{OSeq}}[d]$ acts by projecting onto the summand $W_{\mathbf{s}}$.

Proof. By definition, the identity of the object \mathbf{s} acts by projecting onto the summand $W_{\mathbf{s}}$. To see that the action is well-defined note that two diagrams $D, D' \in \widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{t}}$ are the same if and only if their reductions $\overline{D}, \overline{D}'$ are the same. Moreover, $F(D) = F(D')$ if and only if $F(\overline{D}) = F(\overline{D}')$. Finally assume $D_1 \in \widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{t}}, D_2 \in \widehat{\mathcal{B}}_{\mathbf{t}}^{\mathbf{u}}$, and $D' \in \widehat{\mathcal{B}}_{\mathbf{s}}^{\mathbf{u}}$ then $F(D_2 \cdot D_1) = F(D')$ if $F(\overline{D}_2 \cdot \overline{D}_1) = F(\overline{D}')$. Hence, to see that the action is well-defined and compatible with multiplication it is enough to restrict to reduced sequences. By Remark 4.17 it suffices then to show that the oriented Brauer category in the sense of [BCNR] acts. Luckily in this case we have explicit (monoidal) generators and relations for the morphisms, namely the generators

$$c = \text{cup} \quad d = \text{cap} \quad s = \text{crossing} \quad (4.12)$$

with all possible labellings and the relations (1.4)-(1.9) in [BCNR]. We only need to keep track of the weights: The first two of these relations amount to the fact that a consistently oriented and labelled kink built from a cup and cap has weight 1, the third just requires that $(-1)^{|i||j|}(-1)^{|i||j|} = 1$ for any labels i, j , the fourth just requires

$$(-1)^{|i||j|}(-1)^{|i||k|}(-1)^{|j||k|} = (-1)^{|j||k|}(-1)^{|i||k|}(-1)^{|i||j|}$$

and the relation [BCNR, (1.8)] amounts to the fact that the weight of a crossing does not depend on the orientation, but only on the labels, see Remark 4.26. Finally for [BCNR, (1.9)] it is enough to see that the sum of the weights of the diagram $d \cdot t_3 \cdot c$ over all labellings equals the supertrace δ . But the labellings contributing weight 1 are precisely those with absolute value at most m , hence there are precisely m . The other n possible labelling always create a cup of weight one and a cap of weight -1 or vice versa. Hence, the total weight is $\delta = m - n$. Hence we obtain an action when restricted to reduced sequences and therefore the claim follows. \square

5. THE ISOMORPHISM THEOREM

In this section we prove Theorem A from the introduction. The main step is to establish a commuting diagram of the form

$$\begin{array}{ccc} \text{End}_{\text{osp}(m'|2n)}(V^{\otimes d}) & \xrightarrow{\quad} & \text{End}_{\mathfrak{gl}(m|n)}(\oplus_{\mathbf{s}} W_{\mathbf{s}}) \\ \uparrow \Psi & \nearrow & \uparrow \Theta \\ \text{Br}_d(\delta) & \xrightarrow{\quad \Phi \quad} & \text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s}) \end{array} \quad (5.13)$$

where δ is the supertrace of V , $m' = 2m + 1$ or $m' = 2m$ and Φ a map constructed from the action of $\mathcal{OB}(m - n)$ from Theorem 4.28. For this we view the algebra $\text{Br}_d(\delta)$ as a category with one object, called \star , and endomorphism ring $\text{Br}_d(\delta)$, and construct a functor into $\text{Mat}(\mathcal{OB}(m - n))$, the additive closure of $\mathcal{OB}(m - n)$.

Definition 5.29. *The category $\text{Mat}(\mathcal{OB}(m - n))$ is defined as follows: The objects of $\text{Mat}(\mathcal{OB}(m - n))$ are formal finite direct sums of objects in $\mathcal{OB}(m - n)$, and the morphisms are matrices of morphisms between the summands with addition and composition given by the usual rules of matrix multiplication, see e.g. [B-N] for more details.*

The idea is now to construct a functor which sends the one object of $\text{Br}_d(\delta)$ to the direct sum of all $\mathbf{s} \in \widehat{\text{OSeq}}[d]$ and a Brauer diagram b to a matrix with rows and columns indexed by $\widehat{\text{OSeq}}[d]$. It will be convenient to write a matrix A as $\sum_{\mathbf{s}, \mathbf{t}} 1_{\mathbf{t}} A 1_{\mathbf{s}}$. Diagrammatically we will write the morphisms as formal sums of oriented diagrams, where the orientation sequences stand for the matrix idempotents, see e.g. (5.14).

Due to the fact that we also have the "place holder" symbols \circ in the sequences we need the following additional notion:

Definition 5.30. *Let $b, b' \in \widehat{\mathcal{B}}[d]$ be generalized Brauer diagrams. Then $b' \in \widehat{\mathcal{B}}[d]$ is called a subdiagram of $b \in \mathcal{B}[d]$, denoted $b' \triangleleft b$, if b' refines the partition of b . In other words, the diagram b' is obtained from b by removing some arcs with their orientations and replace them with the appropriate number of \circ 's.*

This allows us to define the functor from $\text{Br}_d(\delta)$ to $\text{Mat}(\mathcal{OB}(m - n))$.

Proposition 5.31. *The assignment*

$$\Psi: \quad \star \mapsto \bigoplus_{\mathbf{s} \in \widehat{\text{OSeq}}[d]} \mathbf{s} \text{ and } b \mapsto M(b),$$

with $M(b)_{(\mathbf{t}, \mathbf{s})} \in \text{Hom}_{\mathcal{OB}(m - n)}(\mathbf{s}, \mathbf{t})$ equal to the unique subdiagram of b in $\widehat{\mathcal{B}}[d]_{\mathbf{s}}^{\mathbf{t}}$ or zero if such a diagram does not exist, defines a faithful functor from $\text{Br}_d(\delta)$ to $\text{Mat}(\mathcal{OB}(m - n))$.

Proof. Clearly, the required subdiagram of b is unique in case it exists. That the functor is faithful is obvious by definition. To see that the functor is well-defined it remains to verify the relations (3.3). The element s_i is sent to the sum of all oriented crossings at place $i, i + 1$ plus its subdiagrams obtained by removing one or two strands in the crossing, for instance for $d = 2$:

$$M(s_i) = \begin{array}{c} \begin{array}{ccccccc} \nearrow & \nearrow & \searrow & \searrow & \circ & \circ & \circ \\ \nwarrow & \nwarrow & \nearrow & \nearrow & \nwarrow & \nwarrow & \nwarrow \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \end{array} \end{array} \quad (5.14)$$

If $d > 2$, then the assignment looks locally as above with the remaining strands oriented in all possible ways.

Note that at vertex i and $i + 1$ (resp. i^* and $(i + 1)^*$) every possible combination from $\{\wedge, \vee, \circ\}$ occurs precisely once and composing $\Psi(s_i)$ with itself gives the identity. The other two braid relations can also be checked easily. The image of e_i looks locally as follows:

$$M(e_i) = \begin{array}{cccccccccccc} \curvearrowright & \curvearrowright & \curvearrowright & \curvearrowright & \circ & \circ & \circ & \circ & \curvearrowright & \curvearrowright & \circ & \circ \\ + & + & + & + & + & + & + & + & + & + & + & + \\ \curvearrowleft & \curvearrowleft & \curvearrowleft & \curvearrowleft & \curvearrowleft & \curvearrowleft & \curvearrowleft & \curvearrowleft & \circ & \circ & \circ & \circ \end{array}$$

If we compose it with itself then we obtain the same sum, but each oriented diagram with an additional clockwise circle, an additional anticlockwise circle and in the odd case also the original diagram itself. Hence, we obtain δ times the original diagram as required. The fifth relation in (3.3) is clear. For the next relation note that $\Psi(e_i)\Psi(e_{i+1})\Psi(e_i)$ is a sum of oriented diagrams of the same underlying shape as the one for $e_i e_{i+1} e_i$ or subdiagrams of this form, but equipped with orientations. Note that each fixed pair of orientation at the bottom and the top appears in precisely one summand and so Ψ preserves the relation $e_i e_{i+1} e_i = e_i$ and similarly also $e_{i+1} e_i e_{i+1} = e_{i+1}$. One can easily check that it also preserves $e_i s_i = e_i = s_i e_i$ and the last two relations of (3.3). \square

Thus, we have now two actions of the Brauer algebra on $V^{\otimes d}$, one by Theorem 3.11 and another one given by Proposition 5.31 and Theorem 4.28.

Lemma 5.32. *The actions of the Brauer algebra given in Theorem 3.11 and the one given by Theorem 4.28 via the embedding of Proposition 5.31 agree.*

Proof. This follows directly from the definitions by a direct calculation on the generators. \square

By Proposition 5.31 we have the induced map Φ from the Brauer algebra to the algebra $\text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s})$. Moreover we have the action map Θ from $\text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s})$ to $\text{End}_{\mathfrak{gl}(m|n)}(V^{\otimes d})$ and know from Lemma 5.32 that the image of $\Theta \circ \Phi$ is contained in $\text{End}_{\mathfrak{osp}(2m+1|2n)}(V^{\otimes d})$. Hence we get the induced map Ψ as indicated in (5.13). We claim that Ψ is an isomorphism if $d \leq m+n$. First note that it is injective, since Φ is injective by definition, and Θ is injective if $d < (m+1)(n+1)$ by [BS2, Theorem 7.8], in particular it is injective if $d \leq m+n$.

Our strategy to prove surjectivity will be to use the surjectivity of Θ from [BS2, Theorem 7.8] and show that any element of $\text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s})$ that commutes with the action of the ortho-symplectic Lie superalgebra is already contained in the image of Φ .

We will show this by an inductive argument. For this we subdivide our set of generalized Brauer diagrams into smaller sets:

$$\widehat{\mathcal{B}}[d] = \coprod_{1 \leq k \leq d, 1 \leq r \leq d} \widehat{\mathcal{B}}[d]_{(k,r)},$$

where $\widehat{\mathcal{B}}[d]_{(k,r)}$ denotes the set of diagrams with exactly $2r$ singleton subsets and k vertical strands, i.e. subsets of the form $\{i, j^*\}$ for some i, j . Furthermore, we note that for the even case we can embed the set of Brauer diagrams into the set of generalized Brauer diagrams as

$$\mathcal{B}[d] = \coprod_{1 \leq k \leq d} \widehat{\mathcal{B}}[d]_{(k,0)}. \quad (5.15)$$

Theorem 5.33. *Assume that one of the following conditions holds:*

$$\text{sdim} V \neq 2m|0 \quad \text{and} \quad d \leq m + n \quad \text{or} \quad (5.16)$$

$$\text{sdim} V = 2m|0 \quad \text{with} \quad m > 0 \quad \text{and} \quad d < m. \quad (5.17)$$

Then the map Ψ is an isomorphism, i.e., $\text{Br}_d(\delta) \cong \text{End}_{\mathfrak{osp}(V)}(V^{\otimes d})$.

Proof. It only remains to prove the surjectivity. For this we proceed as follows: Given an element $f \in \text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s})$ which commutes with the action of $\mathfrak{osp}(V)$ we will show that there is a recursive procedure to write f as a linear combination of elements $M(b)$ for $b \in \mathcal{B}[d]$. This will be done by successively subtracting multiples of $M(b)$'s with decreasing numbers of vertical strands. Since the $M(b)$ for $b \in \mathcal{B}[d]$ span the image of the map Ψ , the claim follows. Write

$$f = \sum_{(\mathbf{t}, b, \mathbf{s}) \text{ oriented}} \gamma_{\mathbf{s}, b, \mathbf{t}} 1_{\mathbf{s}} b 1_{\mathbf{t}}.$$

Let k be maximal and then r minimal such that there exists an oriented generalized Brauer diagram $(\mathbf{t}, b, \mathbf{s})$ with $b \in \widehat{\mathcal{B}}[d]_{(k,r)}$ and $\gamma_{\mathbf{s}, b, \mathbf{t}} \neq 0$.

Claim 1. *The equality $r = 0$ holds. In particular we can find $b \in \widehat{\mathcal{B}}[d]_{(k,0)}$ such that there exists an oriented Brauer diagram $(\mathbf{t}, b, \mathbf{s})$ with $\gamma_{\mathbf{s}, b, \mathbf{t}} \neq 0$.*

For the even case this holds by definition and (5.15). For the odd case this is Corollary 5.35 below. The following is proved in the next paragraph:

Claim 2. *Let $(\mathbf{t}', b, \mathbf{s}')$ be an oriented Brauer diagram with the same underlying diagram b from Claim 1. Then $\gamma_{\mathbf{s}', b, \mathbf{t}'} = \gamma_{\mathbf{s}, b, \mathbf{t}}$.*

We assume for now the two claims hold and fix $b \in \widehat{\mathcal{B}}[d]_{(k,0)}$ as in Claim 1. We denote by $\gamma_b \neq 0$ its coefficient in f . By Claim 2 this is well-defined, i.e. independent of a chosen orientation. Then define

$$f' = f - \sum_{b \in \widehat{\mathcal{B}}[d]_{(k,0)}} \gamma_b M(b).$$

Since $M(b) \in \text{im}(\Phi)$ for all $b \in \widehat{\mathcal{B}}[d]_{(k,0)}$ the surjectivity of the theorem follows if we show that $f' \in \text{im}(\Phi)$. Thanks to Claim 1 we know that f' is contained in the span of the oriented generalized Brauer diagrams $(\mathbf{r}, c, \mathbf{p})$ with $c \in \widehat{\mathcal{B}}[d]_{(l,q)}$ where $l \leq k-1$ or $l = k$ and $q \neq 0$. But then by Claim 1 we have $l \leq k-1$ and some q . Hence, either $f' = 0$ or we can repeat our arguments for f' instead of f and our maximal choice of k strictly decreases in each step. Hence after finitely many steps we reduced the question whether $f \in \text{im}(\Phi)$

to the question whether $0 \in \text{im}(\Phi)$. This is certainly true, and thus the theorem follows. \square

Proof of Claim 2. Our proof treats the situations (5.16) and (5.17) separately distinguishing moreover in (5.16) the cases $n > 0$ respectively $n = 0$.

Let first X be the unique element in $\mathfrak{osp}(V)$ which maps v_{m+1} to $v_{\overline{m+1}}$ and annihilates all other basis elements, that is in terms of Definition 4.18 the matrix d_1 with exactly one non-zero entry 1 in the upper left corner. Its transpose X^T maps $v_{\overline{m+1}}$ to v_{m+1} and annihilates all other basis vectors.

Assume (5.16), ie. $n > 0$ and $d \leq n + m$: It is enough to consider the situation where $(\mathbf{t}, b, \mathbf{s})$ and $(\mathbf{t}', b, \mathbf{s}')$ differ only in the orientation of one strand S , since otherwise we can repeat the argument. Thanks to the assumption $d \leq n + m$ we can pick $\underline{i} \in \text{Vect}(\mathbf{s})$, $\underline{j} \in \text{Vect}(\mathbf{t})$ to get a consistent labelling of $(\mathbf{t}, b, \mathbf{s})$ with the following property

Different strands are labelled with different absolute values, and the strand S is the unique strand labelled with absolute value $m + 1$.

To prove Claim 2 we have to distinguish between three cases, namely the cases where S is a vertical strand, a cup or a cap respectively, ie. the cases where from the two labels $m + 1$ and $\overline{m + 1}$ exactly one, none or both occur in \underline{i} .

For an arbitrary labelling sequence \underline{i} we denote by \underline{i}^\uparrow , \underline{i}^\downarrow and $\underline{i}^\updownarrow$ the sequence obtained by changing all $m + 1$'s into $\overline{m + 1}$'s, by changing all $\overline{m + 1}$'s into $m + 1$'s, or by swapping the labels $\overline{m + 1}$ and $m + 1$ respectively.

Case I: Vertical Strand. Assume S is labelled $m + 1$ (the case of the label $\overline{m + 1}$ is done by replacing the role of X with X^T). Since the label $m + 1$ occurs only at strand S it follows that $Xv_{\underline{i}} = v_{\underline{i}^\uparrow}$. Moreover, $\underline{i}^\uparrow \in \text{Vect}(\mathbf{s}')$, $\underline{j}^\uparrow \in \text{Vect}(\mathbf{t}')$. The equivariance (2.2) of f implies

$$\left\langle f(Xv_{\underline{i}}), v_{\underline{j}^\uparrow}^* \right\rangle = - \left\langle f(v_{\underline{i}}), Xv_{\underline{j}^\uparrow}^* \right\rangle = \left\langle f(v_{\underline{i}^\uparrow}), v_{\underline{j}^\uparrow}^* \right\rangle = \gamma_{\mathbf{s}', b, \mathbf{t}'} \text{wt} \left(b_{\underline{i}^\uparrow}^{\underline{j}^\uparrow} \right),$$

Hence

$$\left\langle f(Xv_{\underline{i}}), v_{\underline{j}}^* \right\rangle = \left\langle f(v_{\underline{i}^\uparrow}), v_{\underline{j}^\uparrow}^* \right\rangle = \gamma_{\mathbf{s}', b, \mathbf{t}'} \text{wt} \left(b_{\underline{i}^\uparrow}^{\underline{j}^\uparrow} \right), \quad (5.18)$$

where the last equality is due to the fact that only $(\mathbf{t}', b, \mathbf{s}')$ can be consistently labelled by \underline{i}^\uparrow and \underline{j}^\uparrow . Similarly, $X^T v_{\underline{j}^\uparrow} = v_{\underline{j}}$

$$- \left\langle f(v_{\underline{i}}), Xv_{\underline{j}^\uparrow}^* \right\rangle = \left\langle f(v_{\underline{i}}), (X^T v_{\underline{j}^\uparrow})^* \right\rangle = \left\langle f(v_{\underline{i}}), v_{\underline{j}}^* \right\rangle = \gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt} \left(b_{\underline{i}}^{\underline{j}} \right). \quad (5.19)$$

The weights in equations (5.18) and (5.19) are equal, since we did not change the parity of the label. Therefore we obtain $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}', b, \mathbf{t}'}$.

Case II: Cup. In case S is a cup, \underline{i} contains no label $m + 1$ and so $Xv_{\underline{i}} = 0$, whereas \underline{j} contains $m + 1$ and $\overline{m + 1}$ and therefore $X^T v_{\underline{j}^\downarrow} = \pm(v_{\underline{j}} + v_{\underline{j}^\uparrow})$. Since

$$\left\langle f(Xv_{\underline{i}}), v_{\underline{j}^\downarrow}^* \right\rangle = - \left\langle f(v_{\underline{i}}), Xv_{\underline{j}^\downarrow}^* \right\rangle \text{ we obtain}$$

$$0 = \left\langle f(v_{\underline{i}}), Xv_{\underline{j}^\dagger}^* \right\rangle = \pm \left(\gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt} \left(b_{\underline{i}}^{\underline{j}} \right) + \gamma_{\mathbf{s}, b, \mathbf{t}'} \text{wt} \left(b_{\underline{i}}^{\underline{j}^\dagger} \right) \right).$$

As we changed only the orientation at a cup with large label, the two weights will exactly differ by a sign and since $\mathbf{s} = \mathbf{s}'$ in this case, we have $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}', b, \mathbf{t}'}$.

Case III: Cap. If S is a cap then $\overline{m+1}$ appears twice in \underline{i}^\dagger whereas $m+1$ does not appear, so $f(v_{\underline{i}^\dagger}) = 0$. To see this note that the label $m+1$ must correspond to vertical strands and therefore any consistently labeled diagram with labels \underline{i}^\dagger at the bottom must have more than k vertical strands and so the claim follows by the maximality of k . Therefore, $\left\langle X.f(v_{\underline{i}^\dagger}), v_{\underline{j}}^* \right\rangle = 0$. On the other hand $X.v_{\underline{i}^\dagger} = \pm (v_{\underline{i}} + v_{\underline{i}^\dagger})$, thus

$$0 = \pm \left\langle f(v_{\underline{i}} + v_{\underline{i}^\dagger}), v_{\underline{j}}^* \right\rangle = \pm \left(\gamma_{\mathbf{s}', b, \mathbf{t}} \text{wt} \left(b_{\underline{i}}^{\underline{j}} \right) + \gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt} \left(b_{\underline{i}^\dagger}^{\underline{j}} \right) \right).$$

We now switched the orientation of a cap with large label and so the two weights differ by a sign. Furthermore $\mathbf{t} = \mathbf{t}'$. Thus we obtain $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}', b, \mathbf{t}'}$.

Assume $\text{sdim } V = 2m+1|0$ and $d \leq m = m+n$: Let $(\mathbf{t}, b, \mathbf{s})$ be our oriented Brauer diagram. Thanks to the assumption on d we can pick $\underline{i} \in \text{Vect}(\mathbf{s})$, $\underline{j} \in \text{Vect}(\mathbf{t})$ to get a consistent labelling of $(\mathbf{t}, b, \mathbf{s})$ with the following property

Different strands are labelled with different absolute values and there is a unique strand S labelled with absolute value 1.

Let X be the unique element in $\mathfrak{osp}(V)$ that maps v_1 to v_0 , v_0 to $-v_{\bar{1}}$ and all other basis elements to zero. In the presentation from Definition 4.18 this means that only $-u_1^T$ contains a non-zero entry, namely a 1 as leftmost entry. Let $(\mathbf{t}^\circ, b^\circ, \mathbf{s}^\circ)$ be the unique oriented Brauer diagram obtained from $(\mathbf{t}, b, \mathbf{s})$ by deleting the strand S and replacing it by two singleton sets. Define \mathbf{t}° and \mathbf{t}° accordingly and let $\underline{i}^\circ \in \text{Vect}(\mathbf{s}^\circ)$ and $\underline{j}^\circ \in \text{Vect}(\mathbf{t}^\circ)$ be the consistent labelling of $(\mathbf{t}^\circ, b^\circ, \mathbf{s}^\circ)$ obtained from \underline{i} and \underline{j} by changing all 1's resp. $\bar{1}$'s into 0's.

We distinguish again three cases:

Case I: Vertical Strand. We assume that S is labelled 1, the case of label $\bar{1}$ is done analogously. Clearly, $Xv_{\underline{i}} = v_{\underline{i}^\circ}$, hence

$$\left\langle f(Xv_{\underline{i}}), v_{\underline{j}^\circ}^* \right\rangle = \left\langle f(v_{\underline{i}^\circ}), v_{\underline{j}^\circ}^* \right\rangle = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ} \text{wt} \left((b^\circ)_{\underline{i}^\circ}^{\underline{j}^\circ} \right), \quad (5.20)$$

where the last equality is due to the fact that only $(\mathbf{t}^\circ, b^\circ, \mathbf{s}^\circ)$ can be consistently labelled by \underline{i}° and \underline{j}° . On the other hand we can use equivariance of f and the fact that $X^T v_{\underline{j}^\circ} = v_{\underline{j}}$ to obtain

$$-\left\langle f(v_{\underline{i}}), Xv_{\underline{j}^\circ}^* \right\rangle = \left\langle f(v_{\underline{i}}), v_{\underline{j}}^* \right\rangle = \gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt} \left(b_{\underline{i}}^{\underline{j}} \right). \quad (5.21)$$

Since v_1 has even parity, the weights in equations (5.20) and (5.21) are equal. Therefore we obtain $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ}$.

Case II: Cup. Hence, \underline{i} contains no 1 and so $Xv_{\underline{i}} = 0$. Note that $\mathbf{s} = \mathbf{s}^\circ$. Now let \underline{j}'' be the sequence obtained from \underline{j} by switching the unique 1 to 0. It holds that $X^T v_{\underline{j}''} = v_{\underline{j}} - v_{\underline{j}^\circ}$ and thus

$$0 = \left\langle f(Xv_{\underline{i}}), v_{\underline{j}''}^* \right\rangle = - \left\langle f(v_{\underline{i}}), Xv_{\underline{j}''}^* \right\rangle = \gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt} \left(b_{\underline{i}}^{\underline{j}} \right) - \gamma_{\mathbf{s}, b^\circ, \mathbf{t}^\circ} \text{wt} \left((b^\circ)_{\underline{i}}^{\underline{j}^\circ} \right).$$

Since we changed a cup with small label, the two appearing weights agree and thus we obtain $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ}$, since $\mathbf{s} = \mathbf{s}^\circ$.

Case III: Cap Note that \underline{i}° is the labelling derived from \underline{i} by replacing 1 and $\bar{1}$ by 0. Let \underline{i}'' be obtained from \underline{i} by switching the unique $\bar{1}$ to 0. With the same argument as in Case III above, the maximality of k implies $f(v_{\underline{i}''}) = 0$. Hence also $\left\langle X.f(v_{\underline{i}''}), v_{\underline{j}}^* \right\rangle = 0$. On the other hand $X.v_{\underline{i}''} = v_{\underline{i}^\circ} - v_{\underline{i}}$, and therefore we obtain

$$0 = \left\langle f(Xv_{\underline{i}''}), v_{\underline{j}}^* \right\rangle = \left\langle f(v_{\underline{i}^\circ} - v_{\underline{i}}), v_{\underline{j}}^* \right\rangle = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ} \text{wt} \left((b^\circ)_{\underline{i}^\circ}^{\underline{j}} \right) - \gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt} \left(b_{\underline{i}}^{\underline{j}} \right).$$

using the fact that $\mathbf{t} = \mathbf{t}^\circ$. As above, we only changed a cap with small labels so the weights agree. Thus we obtain $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ}$.

In all three cases we proved $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ}$. The same arguments apply after we switched the orientation on the strand S , hence $\gamma_{\mathbf{s}', b, \mathbf{t}'} = \gamma_{\mathbf{s}^\circ, b^\circ, \mathbf{t}^\circ}$. This implies the claim $\gamma_{\mathbf{s}, b, \mathbf{t}} = \gamma_{\mathbf{s}', b, \mathbf{t}'}$.

Assume: $\text{sdim} V = 2m|0$ and $d < m = m + n$: This case is very similar to the previous one, but easier. We just replace the occurrences of the label 0 by 2 in the previous argument. For instance we choose X to be the unique element in $\mathfrak{osp}(V)$ that maps v_1 to $v_{\bar{2}}$, v_2 to $-v_{\bar{1}}$ and all other basis elements to zero (in the presentation from Definition 4.18 this means that only the matrix a_1 contains two non-zero entries). Furthermore we assume that no strand is labelled with absolute value 2, which is possible by the assumption $d < m + n$. Then the calculations are the same as in the previous case.

We are left with showing that claim 1 from the proof of Theorem 5.33 holds in the odd case.

Lemma 5.34. *Assume $d \leq m + n$. Let $f \in \text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s})$ be an element that commutes with the action of $\mathfrak{osp}(V)$ and write*

$$f = \sum_{(\mathbf{t}, b, \mathbf{s}) \text{ oriented}} \gamma_{\mathbf{s}, b, \mathbf{t}} 1_{\mathbf{s}} b 1_{\mathbf{t}}.$$

Assume that there exist $\gamma_{\mathbf{s}, b, \mathbf{t}} \neq 0$ such that $b \in \widehat{\mathcal{B}}[d]_{(k, r)}$ for some $r > 0$. Then there exists $\gamma_{\mathbf{s}', b', \mathbf{t}'} \neq 0$ such that $b' \in \widehat{\mathcal{B}}[d]_{(k, r-1)}$ and $b \triangleleft b'$.

Proof. Note that the assumptions of the lemma are never satisfied in the even case, so we can assume that we are in the odd case. Let $\underline{i} \in \text{Vect}(\mathbf{s})$ and $\underline{j} \in \text{Vect}(\mathbf{t})$ be such that $b_{\underline{i}}^{\underline{j}}$ is consistently labelled. In addition we assume that all strands in b are labelled by pairwise different absolute values and none of them is labelled with the absolute value 1. This is possible due

to the assumption $d \leq m + n$ and the assumption that there are less than d strands, since $r > 0$. By the definition of the action we know that the coefficient of $v_{\underline{j}}$ in $F(f)v_{\underline{i}}$, i.e., $\langle F(f)v_{\underline{i}}, v_{\underline{j}}^* \rangle$, is equal to $\gamma_{b, \mathbf{s}, \mathbf{t}} \text{wt}(b_{\underline{i}}^{\underline{j}})$, since this is the only diagram that can be consistently coloured by \underline{i} and \underline{j} .

We first assume that \mathbf{s} contains the symbol \circ at least once. Then let $\underline{i}' \in \text{Vect}(\mathbf{s}(\underline{i}'))$ be equal to \underline{i} except that the first occurring 0 is changed into a 1, furthermore let $X \in \mathfrak{osp}(V)$ be the unique element mapping v_0 to $v_{\bar{1}}$, v_1 to $-v_0$ and send all other basis elements to zero. Due to the assumption that f commutes with X we have the equalities

$$\langle F(f)X.v_{\underline{i}'}, v_{\underline{j}}^* \rangle = \langle F(f)v_{\underline{i}'}, X.v_{\underline{j}}^* \rangle = \langle F(f)v_{\underline{i}'}, (-X^T.v_{\underline{j}})^* \rangle.$$

Computing the very left side we obtain $X.v_{\underline{i}'} = -v_{\underline{i}} + \sum_{\underline{k}} v_{\underline{k}}$, where the sum runs over all those \underline{k} that differ from \underline{i}' by switching one 0 into a $\bar{1}$. Then

$$\langle F(f)X.v_{\underline{i}'}, v_{\underline{j}}^* \rangle = -\gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt}\left(b_{\underline{i}}^{\underline{j}}\right) + \sum_{\underline{k}} \gamma_{\mathbf{s}(\underline{k}), c(\underline{k}), \mathbf{t}} \text{wt}\left(c(\underline{k})_{\underline{k}}^{\underline{j}}\right),$$

where $c(\underline{k})$ is the unique diagram where the 1 and $\bar{1}$ in \underline{k} are connected by a horizontal arc and $\underline{k} \in \text{Vect}(\mathbf{s}(\underline{k}))$. For all $c(\underline{k})$ it holds that $b \triangleleft c(\underline{k})$ and they contain one more strand than b .

To compute the right side we see that $-X^T.v_{\underline{j}} = -\sum_{\underline{l}} v_{\underline{l}}$, where the sum runs over those \underline{l} that differ from \underline{j} by switching exactly one 0 into a 1. Hence

$$\langle F(f)v_{\underline{i}'}, X.v_{\underline{j}}^* \rangle = -\sum_{\underline{l}} \gamma_{\mathbf{s}(\underline{i}'), d(\underline{l}), \mathbf{t}(\underline{l})} \text{wt}\left(d(\underline{l})_{\underline{i}'}^{\underline{l}}\right),$$

where $d(\underline{l})$ is the unique diagram where the 1's in both \underline{i}' and \underline{l} are connected by a vertical strand and $\underline{l} \in \text{Vect}(\mathbf{t}(\underline{l}))$. Again it holds for all $d(\underline{l})$ that $b \triangleleft d(\underline{l})$ and that they contain one more strand than b .

Putting the two sides together we obtain

$$\gamma_{\mathbf{s}, b, \mathbf{t}} \text{wt}\left(b_{\underline{i}}^{\underline{j}}\right) = \sum_{\underline{k}} \gamma_{\mathbf{s}(\underline{k}), c(\underline{k}), \mathbf{t}} \text{wt}\left(c(\underline{k})_{\underline{k}}^{\underline{j}}\right) + \sum_{\underline{l}} \gamma_{\mathbf{s}(\underline{i}'), d(\underline{l}), \mathbf{t}(\underline{l})} \text{wt}\left(d(\underline{l})_{\underline{i}'}^{\underline{l}}\right).$$

Since all the occurring weights are ± 1 we see that if $\gamma_{\mathbf{s}, b, \mathbf{t}} \neq 0$ then there must be at least one coefficient on the right that is also non-zero.

The case that \mathbf{s} contains no \circ is dealt with in an analogous fashion by swapping the roles of \underline{i} and \underline{j} . \square

From this it follows that we can always find a chain of diagrams with non-zero coefficients that end in an oriented (non-generalized) Brauer diagram.

Corollary 5.35. *Assume $d \leq m + n$. Let $f \in \text{End}_{\text{Mat}(\mathcal{OB}_d(m-n))}(\oplus_{\mathbf{s}} \mathbf{s})$ be an element that commutes with the action of $\mathfrak{osp}(V)$ and write*

$$f = \sum_{(\mathbf{t}, b, \mathbf{s}) \text{ oriented}} \gamma_{\mathbf{s}, b, \mathbf{t}} 1_{\mathbf{s}} b 1_{\mathbf{t}}.$$

Assume that there exist $\gamma_{\mathbf{s}_r, b_r, \mathbf{t}_r} \neq 0$ such that $b_r \in \widehat{\mathcal{B}}[d]_{(k, r)}$ for some $r > 0$. Then there exists a sequence $((\mathbf{s}_i, b_i, \mathbf{t}_i))_{0 \leq i \leq r}$ of oriented generalized Brauer

diagrams with $b_i \in \widehat{\mathcal{B}}[d]_{(k_i, i)}$ for some $k_i \geq k$ such that $\gamma_{\mathbf{s}_i, b_i, \mathbf{t}_i} \neq 0$ for all i and $b_r \triangleleft b_{r-1} \triangleleft \dots \triangleleft b_1 \triangleleft b_0$.

Epecially note that $b_0 \in \widehat{\mathcal{B}}[d]_{(k_0, 0)}$.

Proof. This follows immediately by successively applying Lemma 5.34 to $(\mathbf{s}_i, b_i, \mathbf{t}_i)$ and setting $(\mathbf{s}_{i-1}, b_{i-1}, \mathbf{t}_{i-1}) := (\mathbf{s}', b', \mathbf{t}')$. \square

Remark 5.36. *In general, the map Ψ is not surjective. In [LZ1] it was shown that for $d \geq m(2n+1)$ in the even case the map Ψ is not surjective. This does not contradict Theorem 5.33 since in the case of $m, n \neq 0$ we have $m(2n+1) = 2mn + m$ which is strictly greater than the assumed bound in the theorem and in the case of $n = 0$ it holds $m(2n+1) = m$ which is also strictly greater.*

Remark 5.37. *Similar to the definition of the oriented Brauer category $\mathcal{OB}(m-n)$, one defines the Brauer category $\mathcal{Br}(\delta)$ using Brauer diagrams having different numbers of vertices at top and bottom. Both, the embedding into the additive closure of the oriented Brauer category as well as the proof of surjectivity generalise to this situation, as they only rely on the diagrammatic description. The number d has to be substituted for the number of arcs in each morphism space.*

6. THE ISOMORPHISM THEOREM FOR ARBITRARY FIELDS

Let \mathbb{k} be an arbitrary field. Consider $\mathfrak{osp}(V)$ in the special cases where $m = 0$ or $n = 0$. This is a classical complex reductive Lie algebra which we denote by \mathfrak{g} . Let $\mathcal{U}_{\mathbb{Z}}$ be Kostant's \mathbb{Z} -form, [H, Section 26], of the universal enveloping algebra of \mathfrak{g} and $\mathcal{U}_{\mathbb{k}} = \mathcal{U}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{k}$ its specialization to \mathbb{k} . Let $V_{\mathbb{Z}}$ be the integral version of the vector representation and $V_{\mathbb{k}}$ its specialisation to \mathbb{k} . On the other hand we denote by $\mathrm{Br}_d(\delta)_{\mathbb{k}}$ resp. $\mathcal{OB}_d(m-n)_{\mathbb{k}}$ the Brauer algebra, resp. oriented Brauer category, for the same parameters as above but defined now over \mathbb{k} , and by $\Phi_{\mathbb{k}}$ and $\Theta_{\mathbb{k}}$ the previous maps defined as maps of \mathbb{k} -vector spaces. (All of them as well as the action of the (oriented) Brauer category were obviously defined originally over the integers.)

Theorem 6.38. *Assume that one of the following conditions holds:*

- *the even orthogonal case: $\mathrm{sdim} V = 2m|0$ and $d < m$ or*
- *the odd orthogonal case: $\mathrm{sdim} V = 2m+1|0$ and $d \leq m$ or*
- *the symplectic case: $\mathrm{sdim} V = 0|2n$ with $d \leq n$.*

Then there is a natural isomorphism of \mathbb{k} -algebras $\mathrm{Br}_d(\delta)_{\mathbb{k}} \cong \mathrm{End}_{\mathcal{U}_{\mathbb{k}}}(V_{\mathbb{k}}^{\otimes d})$.

Proof. The map $\Phi_{\mathbb{k}}$ from $\mathrm{Br}_d(\delta)_{\mathbb{k}}$ to $\mathrm{End}_{\mathrm{Mat}(\mathcal{OB}_d(m-n)_{\mathbb{k}})}(\bigoplus_{\mathbf{s}} \mathbf{s})$ is defined exactly as before. All basis vectors were originally already mapped to a linear combination of basis vectors with all coefficients 0 or 1, and hence the map remains obviously injective. By [DDS, Corollary 6.2] and the fact that Lemma 4.21 remains true over an arbitrary field, the map $\Theta_{\mathbb{k}}$ is an isomorphism under our assumptions on d . Thus it only remains to show that $\Psi_{\mathbb{k}} := \Theta_{\mathbb{k}} \circ \Phi_{\mathbb{k}}$ remains surjective. This is true since the proof of Theorem 5.33

is characteristic free: a lift of the the elements X exist in the integral form $\mathcal{U}_{\mathbb{Z}}$ and all calculations are over the integers. \square

Remark 6.39. *Alternatively one can show the surjectivity of $\Psi_{\mathbb{k}}$ in case the characteristic of \mathbb{k} is not 2 by a dimension argument involving the theory of tilting modules: Since $V_{\mathbb{k}}$ is an irreducible $\mathcal{U}_{\mathbb{k}}$ -tilting module, see e.g. [AST], the dimension of $\text{End}_{\mathcal{U}_{\mathbb{k}}}(V_{\mathbb{k}}^{\otimes d})$ is independent of the field, [AST, (21)]. Thus our injective map (which is surjective over the complex numbers) remains surjective over any field, as the dimension of $\text{Br}_d(\delta)_{\mathbb{k}}$ is also independent of the field. (The assumption on the characteristic is necessary since otherwise $V_{\mathbb{k}}$ is not necessarily tilting in the even orthogonal case, see [J, p.20]).*

REFERENCES

- [AST] H.H. Andersen, C. Stroppel, and D. Tubbenhauer *Cellular Structures using U_q -tilting modules*, in preparation.
- [B-N] D. Bar-Natan, *Khovanov homology for tangles and cobordisms*, Geometry and Topology, **9** (2005) 1443–1499.
- [BCHLLS] G. Benkart, M. Chakrabarti, T. Halverson, R. Leduc, C. Lee and J. Stroomeer, *Tensor product representations of general linear groups and their connections with Brauer algebras*, J. Algebra **166** (1994), 529–567.
- [BR] A. Berele and A. Regev, *Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras*, Adv. Math. **64** (1987), 118–175.
- [Br] R. Brauer. *On algebras which are connected with the semisimple continuous groups*, Ann. of Math. (2), **38** (4):857–872, 1937.
- [BCNR] J. Brundan, J. Comes, D. Nash, and A. Reynolds *A basis theorem for the affine oriented Brauer category and its cyclotomic quotients* arXiv:1404.6574.
- [BS1] J. Brundan and C. Stroppel, *Highest weight categories arising from Khovanov’s diagram algebra III: category \mathcal{O}* , Represent. Theory **15** (2011), 170–243.
- [BS2] ———, *Gradings on walled Brauer algebras and Khovanov’s arc algebra*, Adv. Math. **231** (2012), no. 2, 709–773.
- [CW] J. Comes and B. Wilson, *Deligne’s category $\text{Rep}(\text{GL}(\delta))$, representations of general linear supergroups*. Represent. Theory **16**, 568–609, 2012.
- [Co] K. Coulembier, *The orthosymplectic superalgebra in harmonic analysis*. J. Lie Theory **23** (2013), no. 1, 5583.
- [De] P. Deligne: *La serie exceptionnelle de groupes de Lie*, C. R. Acad. Sci. Paris Ser. I Math. **322**, no. 4, 321–326, 1996.
- [DDS] R. Dipper, S. Doty and F. Stoll, *The quantized walled Brauer algebra and mixed tensor space*. Algebr. Represent. Theory **17** (2014), no. 2, 675–701.
- [ES1] M. Ehrig and C. Stroppel, *Nazarov-Wenzl algebras, coideal subalgebras and categorified skew Howe duality*, arXiv:1310.1972.
- [ES2] M. Ehrig and C. Stroppel, *Gradings on Brauer algebras*, in preparation.
- [GW] R. Goodman and N. R. Wallach, *Symmetry, representations, and invariants*, Graduate Texts in Mathematics **255**, Springer (2009).
- [GS] C. Gruson, and V. Serganova, *Bernstein-Gelfand-Gelfand reciprocity and indecomposable projective modules for classical algebraic supergroups*. Mosc. Math. J. **13**, no. 2, 281–313, 364, 2013.
- [H] J. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Math., Springer, 1972.
- [J] J.C. Jantzen, *Darstellungen halbeinfacher algebraischer Gruppen und zugeordnete kontravariante Formen*, Bonn. Math. Schr. No. 67, 1973.

- [KM] M. Kosuda and J. Murakami, *Centralizer algebras of the mixed tensor representations of quantum group $U_q(\mathfrak{gl}(n, \mathbb{C}))$* . Osaka J. Math. **30** (1993), no. 3, 475–507.
- [LZ1] G. Lehrer and R. Zhang. *The first fundamental theorem of invariant theory for the orthosymplectic supergroup*, arXiv:1401.7395.
- [LZ2] G. Lehrer and R. Zhang. *The second fundamental theorem of invariant theory for the orthosymplectic supergroup*, Ann. of Math. (2) **176** (2012), no. 3, 2031–2054.
- [Li] G. Li, *A KLR Grading of the Brauer Algebras*, arXiv:1409.1195.
- [L] G. Lusztig. *Quantum groups at roots of 1*, Geom. Dedicata **35** (1990), no. 1-3, 89–113.
- [Mu] I. M. Musson, *Lie superalgebras and enveloping algebras*, Graduate Studies in Mathematics **131**, American Mathematical Society (2012).
- [Ni] P. Nikitin, *The centralizer algebra of the diagonal action of the group $GL_n(\mathbb{C})$ in a mixed tensor space*, J. Math. Sci. **141** (2007), 1479–1493.
- [SS] A. Sartori and C. Stroppel: *Walled Brauer algebras as idempotent truncations of level 2-cyclotomic quotients*, arXiv:1411.2771 .
- [Sea] V. Serganova, *Finite dimensional representations of algebraic supergroups*, ICM Plenary talk Seoul 2014.
- [Sev] A. Sergeev, *Tensor algebra of the identity representation as a module over the Lie superalgebras $GL(n, m)$ and $Q(n)$* , Math. USSR Sbornik, **51** (1985), 419–427.
- [Tu] V. Turaev, *Operator invariants of matrices and R-matrices*, Izv. Akad. Nauk SSSR **53** (1989) 1073–1107.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BONN, 53115 BONN, GERMANY
E-mail address: mehrig@math.uni-bonn.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BONN, 53115 BONN, GERMANY
E-mail address: stroppel@math.uni-bonn.de