

# Week 7: Tests for Differences

## Session 1

Spring 2020

## iClicker Question 1

Which of the following should I use to read the file `mydata.csv` into a data frame called `dat` in R?

A

B

C

D

E

If you're feeling stuck, remember my office hours are Tuesday/Thursday 1:00 - 2:00.



# For Today

- ▶ Toward statistics
- ▶ Tests for differences

Follow-up questions from the Chapter 6 homework


- ▶ What questions do you have?

## Beyond graphs, Towards statistics

- ▶ Graphs are powerful tools that provide insight and understanding of the patterns and relationships in the data.
- ▶ Graphs alone don't give us the complete answer. We need to **quantify** the relationships we see in our plots.



# Beyond graphs, Towards statistics

- ▶ How can we **quantify** our evidence for relationships?
  - ▶ Are differences between groups *significant*?
  - ▶ Are differences between groups *meaningful*?



# Beyond graphs, Towards statistics

- How can we **quantify** our evidence for relationships?
  - Are associations between 2 variables *significant*?
  - Are associations between 2 variables *meaningful*?



# Beyond graphs, Towards statistics

- ▶ Statistics is the tool we use to formally answer these questions!
  - ▶ Are differences *are/are not* significant!
  - ▶ Are associations *are/are not* significant!

But... what do we mean when we say **significant**?



# Let's examine some plots to gain intuition:

- Scenario: We want to know whether the size of 3-year-old bluegill (*Lepomis macrochirus*) are larger in some Massachusetts lakes than others.
- We have collected data for bluegill from Wyola Lake and the Quabbin Reservoir in Western Mass.




---

<sup>1</sup>Image credit: New York Fish and Game Commission


# Bluegill Data I

- ▶ Are differences between lakes *significant*?
- ▶ Are differences between lakes *meaningful*?




# Bluegill Data II

- ▶ Are differences between lakes *significant*?
- ▶ Are differences between lakes *meaningful*?




# Bluegill Data III

- ▶ Are differences between lakes *significant*?
- ▶ Are differences between lakes *meaningful*?



# Bluegill Data IV

- ▶ Are differences between lakes *significant*?
- ▶ Are differences between lakes *meaningful*?






# Tests for differences

Often we want to know if two of more samples are different


- ▶ are the sample *means* different?
- ▶ are the sample *medians* different?
- ▶ are the differences *statistically significant*?



# Tests for differences

Often we want to know if two of more samples are different

- ▶ are the sample *means* different?
- ▶ are the sample *medians* different?
- ▶ are the differences *statistically significant*?






# Tests for differences

Often we want to know if two of more samples are different

- ▶ are the sample *means* different?
- ▶ are the sample *medians* different?
- ▶ are the differences *statistically significant*?



# Tests for differences

Often we want to know if two of more samples are different

- ▶ are the sample *means* different?
- ▶ are the sample *medians* different?
- ▶ are the differences *statistically significant*?

To determine the significance of differences between **two**, we need a statistical test

- ▶ *t-test*
- ▶ *U-test*

# Differences: t-test

Purpose:

- ▶ compare the means of two samples (say  $a$  and  $b$ )

Assumptions:

- ▶ both samples normally distributed
- ▶ both samples have equal variances

# Differences: t-test

Purpose:

- ▶ compare the means of two samples (say  $a$  and  $b$ )

Assumptions:

- ▶ both samples normally distributed
- ▶ both samples have equal variances

$$t = \frac{|\bar{x}_a - \bar{x}_b|}{\sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}}$$

- ▶  $t$ : the  $t$ -statistic
- ▶  $\bar{x}$ : sample mean
- ▶  $s$ : sample standard deviation
- ▶  $n$ : sample size

# Differences: t-test

Purpose:

- ▶ compare the means of two samples (say  $a$  and  $b$ )

Assumptions:

- ▶ both samples normally distributed
- ▶ both samples have equal variances

$$t = \frac{|\bar{x}_a - \bar{x}_b|}{\sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}}$$

- ▶ if  $|\bar{x}_a - \bar{x}_b|$  is large, then  $t$  is ????
- ▶ if  $\sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}$  is large, then  $t$  is ????

# Differences: t-test

Purpose:

- ▶ compare the means of two samples (say  $a$  and  $b$ )

Assumptions:


- ▶ both samples normally distributed
- ▶ both samples have equal variances

$$t = \frac{|\bar{x}_a - \bar{x}_b|}{\sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}}$$

- ▶ if  $|\bar{x}_a - \bar{x}_b|$  is large, then  $t$  is **large**
- ▶ if  $\sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}$  is large, then  $t$  is **small**


# Differences: t-test

Understanding the *t-distribution*:



# Differences: t-test


Understanding the *t*-distribution:






# Differences: t-test

Understanding the *t-distribution*:



# Differences: t-test


Understanding the *t-distribution*:



# Differences: t-test

Understanding the *t-distribution*:

- ▶ whether a difference is significant depends on:
  - ▶ the *t-statistic*
  - ▶ degrees-of-freedom ( $n_a - 1 + n_b - 1$ )
- ▶ larger *t-statistics* more likely to be significant



# Differences: t-test

Understanding the *p-value*:

- ▶ *p-value* is the probability of observing a *t-statistic* as high as we did by chance
- ▶ if *p-value* is lower than significance level (e.g. 5%):
  - ▶ difference is significant
  - ▶ reject the null hypothesis
  - ▶ accept the alternative hypothesis

# Differences: t-test

Which *t-test*?

- ▶ standard *t-test*
  - ▶ compare two independent samples
  - ▶ both normally distributed
  - ▶ equal (similar) variances
  - ▶ samples sizes can be the same or not

$$t = \frac{|\bar{x}_a - \bar{x}_b|}{\sqrt{\frac{s_a^2}{n_a} + \frac{s_b^2}{n_b}}}$$

- ▶  $t$ : the *t*-statistic
- ▶  $\bar{x}$ : sample mean
- ▶  $s$ : sample standard deviation
- ▶  $n$ : sample size

# Differences: paired t-test

Sometimes samples are not independent

- ▶ compare pairs of samples
  - ▶ e.g., before-after
  - ▶ e.g., north-south
  - ▶ e.g., left-right
- ▶ both normally distributed
- ▶ equal (similar) variances
- ▶ samples sizes *must* be the

# Differences: paired t-test

Which *t-test*?

- ▶ paired *t-test*
  - ▶ compare pairs of samples
  - ▶ both normally distributed
  - ▶ equal (similar) variances
  - ▶ samples sizes *must* be the

$$t = \frac{\bar{D}}{\sqrt{\frac{s_D^2}{n}}}$$


- ▶ *t*: the *t*-statistic
- ▶  $\bar{D}$ : mean of the *differences*
- ▶ *s*: standard deviation of the *differences*
- ▶ *n*: number of *paired* samples





# Differences: U-test

- ▶ compare two samples
- ▶ both *not* normally distributed
- ▶ based on *median*, *range*, and *ranks*
- ▶ rank all values as one sample, calculate group rank sums  $R$
- ▶ calculate a  $U$ -value, a measure of overlap



## Differences: U-test

- ▶ compare two samples
- ▶ both or differences *not* normally distributed
- ▶ based on *median*, *range*, and *ranks*
- ▶ rank all values as one sample, calculate group rank sums  $R$
- ▶ calculate a  $U$ -value, a measure of overlap

$$U_a = n_a \times n_b + \frac{n_a(n_a + 1)}{2} - R_a$$

$$U_b = n_b \times n_a + \frac{n_b(n_b + 1)}{2} - R_b$$

- ▶  $n_a$ : number of samples in sample  $a$
- ▶  $n_b$ : number of samples in sample  $b$
- ▶  $R_a$ : sum of the ranks of values in  $a$
- ▶  $R_b$ : sum of the ranks of values in  $b$

## Differences: U-test

- ▶ compare two samples
- ▶ both or differences *not* normally distributed
- ▶ based on *median*, *range*, and *ranks*
- ▶ rank all values as one sample, calculate group rank sums  $R$
- ▶ calculate a  $U$ -value, a measure of overlap

$$U_a = n_a \times n_b + \frac{n_a(n_a + 1)}{2} - R_a$$

$$U_b = n_b \times n_a + \frac{n_b(n_b + 1)}{2} - R_b$$

- ▶ smallest is used to find the  $p$ -value
- ▶ unlike the t-statistic, lower  $U$ -values are more likely to be significant

## Differences: Wilcoxon matched-pairs test

- ▶ both or differences *not* normally distributed
- ▶ based on ranked *differences*
  - ▶ first calculate the differences
  - ▶ second rank the differences
  - ▶ 0's not ranked
- ▶ sum and compare +ve and -ve ranks

$$W^+ = \sum R^+$$

$$W^- = \sum R^-$$

- ▶  $W^+$ : the Wilcoxon test statistic for positive differences
- ▶  $W^-$ : the Wilcoxon test statistic for negative differences
- ▶  $R^+$ : the sum of the ranks of positive differences
- ▶  $R^-$ : the sum of the ranks of negative differences

# Differences: Wilcoxon matched-pairs test

- ▶ pairs or differences *not* normally distributed
- ▶ based on ranked *differences*
  - ▶ first calculate the differences
  - ▶ second rank the differences
  - ▶ 0's not ranked
- ▶ sum and compare +ve and -ve ranks

$$W^+ = \sum R^+$$

$$W^- = \sum R^-$$

- ▶ smallest is used to find the  $p$ -value
- ▶ lower  $W$ -values are more likely to be significant

# Group Assignment

Using the whale count data, compare the differences between first and second abundance guesstimates using first `excel` and then `R`.

Submit a single written group report that outlines the following points:

1. state the null and alternative hypotheses being tested
2. the reason for choosing the statistical test you used
3. a summary of the results:
  - ▶ degrees-of-freedom, test statistic, p-values (at 5% level)?
  - ▶ did you accept or reject the null hypothesis?
  - ▶ is there a difference?
4. conduct the analysis in `R` and `excel` and submit:
  - ▶ a written report of points 1, 2 and 3 as **PDF**
  - ▶ an `excel` workbook showing your results
  - ▶ a saved `R` file showing results