

Linear Regression

Mario Parente

A First Supervised Learning Problem

How do you measure the biomass of a forest?

A First Supervised Learning Problem

How do you measure the biomass of a forest?

Hard to measure:

- ▶ Mass of tree

A First Supervised Learning Problem

How do you measure the biomass of a forest?

Hard to measure:

- ▶ Mass of tree
- ▶ Height of tree (but can be done)

A First Supervised Learning Problem

How do you measure the biomass of a forest?

Hard to measure:

- ▶ Mass of tree
- ▶ Height of tree (but can be done)

Easy to measure:

- ▶ Diameter at breast height (DBH)

A First Supervised Learning Problem

How do you measure the biomass of a forest?

Hard to measure:

- ▶ Mass of tree
- ▶ Height of tree (but can be done)

Easy to measure:

- ▶ Diameter at breast height (DBH)

Let's simplify the problem: devise method to easily estimate the height of a tree

A First Supervised Learning Problem

Idea?

A First Supervised Learning Problem

Idea?

- ▶ Collect data on DBH and height for some trees

A First Supervised Learning Problem

Idea?

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height


A First Supervised Learning Problem


Idea?

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

Some data


Development and Evaluation of Models for the Relationship between Tree Height and Diameter at Breast Height for Chinese-Fir Plantations in Subtropical China


Yan-qiong Li, Xiang-wen Deng , Zhi-hong Huang, Wen-hua Xiang, Wen-de Yan, Pi-feng Lei, Xiao-lu Zhou, Chang-hui Peng



Some data

Development and Evaluation of Models for the Relationship between Tree Height and Diameter at Breast Height for Chinese-Fir Plantations in Subtropical China

Yan-qiong Li, Xiang-wen Deng , Zhi-hong Huang, Wen-hua Xiang, Wen-de Yan, Pi-feng Lei, Xiao-lu Zhou, Chang-hui Peng



What do you predict for the height of a tree with DBH 15cm?
35cm? Why?

A First Supervised Learning Problem

Idea:

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

A First Supervised Learning Problem

Idea:

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

This is **supervised learning**:

A First Supervised Learning Problem

Idea:

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

This is **supervised learning**:

- ▶ Collect **training data**

A First Supervised Learning Problem

Idea:

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

This is **supervised learning**:

- ▶ Collect **training data**
- ▶ Use a **learning algorithm** to fit a **model**

A First Supervised Learning Problem

Idea:

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

This is **supervised learning**:

- ▶ Collect **training data**
- ▶ Use a **learning algorithm** to fit a **model**
- ▶ Use model to make a **prediction**

A First Supervised Learning Problem

Idea:

- ▶ Collect data on DBH and height for some trees
- ▶ Determine relationship between DBH and height
- ▶ Use DBH to predict height for a new tree

This is **supervised learning**:

- ▶ Collect **training data**
- ▶ Use a **learning algorithm** to fit a **model**
- ▶ Use model to make a **prediction**

What model? What algorithm? **Largely what this class is about.**


Supervised Learning

DBH (x)	Height (y)
17	63
19	65
20.5	66
...	...

Supervised Learning

DBH (x)	Height (y)
17	63
19	65
20.5	66
...	...

Find h such that $h(x) \approx y$



Supervised Learning: Notation and Terminology

- ▶ Observe m “training examples” of form $(x^{(i)}, y^{(i)})$
 - ▶ $x^{(i)}$: **features** / input / what we observe / DBH
 - ▶ $y^{(i)}$: **target** / output / what we want to predict / height

Supervised Learning: Notation and Terminology

- ▶ Observe m “training examples” of form $(x^{(i)}, y^{(i)})$
 - ▶ $x^{(i)}$: **features** / input / what we observe / DBH
 - ▶ $y^{(i)}$: **target** / output / what we want to predict / height
 - ▶ **Training set** $\{(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})\}$

Supervised Learning: Notation and Terminology

- ▶ Observe m “training examples” of form $(x^{(i)}, y^{(i)})$
 - ▶ $x^{(i)}$: **features** / input / what we observe / DBH
 - ▶ $y^{(i)}$: **target** / output / what we want to predict / height
 - ▶ **Training set** $\{(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})\}$
- ▶ Find function (“hypothesis”) h such that $h(x) \approx y$

Supervised Learning: Notation and Terminology

- ▶ Observe m “training examples” of form $(x^{(i)}, y^{(i)})$
 - ▶ $x^{(i)}$: **features** / input / what we observe / DBH
 - ▶ $y^{(i)}$: **target** / output / what we want to predict / height
 - ▶ **Training set** $\{(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})\}$
- ▶ Find function (“hypothesis”) h such that $h(x) \approx y$
 - ▶ $h(x^{(i)}) \approx y^{(i)}$ – good fit on training data

Supervised Learning: Notation and Terminology

- ▶ Observe m “training examples” of form $(x^{(i)}, y^{(i)})$
 - ▶ $x^{(i)}$: **features** / input / what we observe / DBH
 - ▶ $y^{(i)}$: **target** / output / what we want to predict / height
 - ▶ **Training set** $\{(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})\}$
- ▶ Find function (“hypothesis”) h such that $h(x) \approx y$
 - ▶ $h(x^{(i)}) \approx y^{(i)}$ – good fit on training data
 - ▶ **Generalize** well to new x values

Supervised Learning: Notation and Terminology

- ▶ Observe m “training examples” of form $(x^{(i)}, y^{(i)})$
 - ▶ $x^{(i)}$: **features** / input / what we observe / DBH
 - ▶ $y^{(i)}$: **target** / output / what we want to predict / height
 - ▶ **Training set** $\{(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})\}$
- ▶ Find function (“hypothesis”) h such that $h(x) \approx y$
 - ▶ $h(x^{(i)}) \approx y^{(i)}$ – good fit on training data
 - ▶ **Generalize** well to new x values

Variations: type of x , y , h

Linear Regression in One Variable

First example of supervised learning. Assume hypothesis is a linear function:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Linear Regression in One Variable

First example of supervised learning. Assume hypothesis is a linear function:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

► θ_0 : intercept, θ_1 : slope

Linear Regression in One Variable

First example of supervised learning. Assume hypothesis is a linear function:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

- ▶ θ_0 : intercept, θ_1 : slope
- ▶ “parameters” or “weights”


Linear Regression in One Variable

First example of supervised learning. Assume hypothesis is a linear function:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

- ▶ θ_0 : intercept, θ_1 : slope
- ▶ “parameters” or “weights”

How to find “best” θ_0, θ_1 ?



Finding the best hypothesis

Simplification: “slope-only” model $h_{\theta}(x) = \theta_1 x$

Finding the best hypothesis

Simplification: “slope-only” model $h_{\theta}(x) = \theta_1 x$

- We only need to find θ_1

Finding the best hypothesis

Simplification: “slope-only” model $h_{\theta}(x) = \theta_1 x$

- We only need to find θ_1

Idea: design **cost function** $J(\theta_1)$ to numerically measure the quality of hypothesis $h_{\theta}(x)$

Finding the best hypothesis

Simplification: “slope-only” model $h_{\theta}(x) = \theta_1 x$

► We only need to find θ_1

Idea: design **cost function** $J(\theta_1)$ to numerically measure the quality of hypothesis $h_{\theta}(x)$

Exercise: which cost functions below make sense?

A. $J(\theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$

B. $J(\theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

C. $J(\theta_1) = \sum_{i=1}^m |h_{\theta}(x^{(i)}) - y^{(i)}|$

1. A only

2. B only

3. C only

4. B and C

5. A, B, and C

Finding the best hypothesis

Simplification: “slope-only” model $h_{\theta}(x) = \theta_1 x$

► We only need to find θ_1

Idea: design **cost function** $J(\theta_1)$ to numerically measure the quality of hypothesis $h_{\theta}(x)$

Exercise: which cost functions below make sense?

A. $J(\theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$

B. $J(\theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

C. $J(\theta_1) = \sum_{i=1}^m |h_{\theta}(x^{(i)}) - y^{(i)}|$

1. A only

2. B only

3. C only

4. B and C

5. A, B, and C

Answer. 4

Squared Error Cost Function

The “squared error” cost function is:

$$J(\theta_1) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Squared Error Cost Function

The “squared error” cost function is:

$$J(\theta_1) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

► E.g., $\theta_1 = 3$:

x	y	$(3x - y)^2/2$
17	63	$(51 - 63)^2/2 = 144/2$

Squared Error Cost Function

The “squared error” cost function is:

$$J(\theta_1) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

► E.g., $\theta_1 = 3$:

x	y	$(3x - y)^2/2$
17	63	$(51 - 63)^2/2 = 144/2$
19	65	$(57 - 65)^2/2 = 64/2$

Squared Error Cost Function

The “squared error” cost function is:

$$J(\theta_1) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

► E.g., $\theta_1 = 3$:

x	y	$(3x - y)^2/2$
17	63	$(51 - 63)^2/2 = 144/2$
19	65	$(57 - 65)^2/2 = 64/2$
20.5	66	$(61.5 - 66)^2/2 = 20.25/2$

Squared Error Cost Function

The “squared error” cost function is:

$$J(\theta_1) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

► E.g., $\theta_1 = 3$:

x	y	$(3x - y)^2/2$
17	63	$(51 - 63)^2/2 = 144/2$
19	65	$(57 - 65)^2/2 = 64/2$
20.5	66	$(61.5 - 66)^2/2 = 20.25/2$

$$J(3) = (144 + 64 + 20.25)/2 = 220.25/2$$

Our First Algorithm

We can use calculus to find the hypothesis of minimum cost. Set the derivative of J to zero and solve for θ_1 . For this example:

Our First Algorithm

We can use calculus to find the hypothesis of minimum cost. Set the derivative of J to zero and solve for θ_1 . For this example:

$$J(\theta_1) = \frac{1}{2} \left((17 \cdot \theta_1 - 63)^2 + (19 \cdot \theta_1 - 65)^2 + (20.5 \cdot \theta_1 - 66)^2 \right)$$

Our First Algorithm

We can use calculus to find the hypothesis of minimum cost. Set the derivative of J to zero and solve for θ_1 . For this example:

$$\begin{aligned} J(\theta_1) &= \frac{1}{2} \left((17 \cdot \theta_1 - 63)^2 + (19 \cdot \theta_1 - 65)^2 + (20.5 \cdot \theta_1 - 66)^2 \right) \\ &= 535.125 \cdot \theta_1^2 - 3659 \cdot \theta_1 + 6275 \end{aligned}$$

Our First Algorithm

We can use calculus to find the hypothesis of minimum cost. Set the derivative of J to zero and solve for θ_1 . For this example:

$$\begin{aligned} J(\theta_1) &= \frac{1}{2} \left((17 \cdot \theta_1 - 63)^2 + (19 \cdot \theta_1 - 65)^2 + (20.5 \cdot \theta_1 - 66)^2 \right) \\ &= 535.125 \cdot \theta_1^2 - 3659 \cdot \theta_1 + 6275 \end{aligned}$$

$$0 = \frac{d}{d\theta_1} J(\theta_1) = 1070.25 \cdot \theta_1 - 3659$$

Our First Algorithm

We can use calculus to find the hypothesis of minimum cost. Set the derivative of J to zero and solve for θ_1 . For this example:


$$\begin{aligned} J(\theta_1) &= \frac{1}{2} \left((17 \cdot \theta_1 - 63)^2 + (19 \cdot \theta_1 - 65)^2 + (20.5 \cdot \theta_1 - 66)^2 \right) \\ &= 535.125 \cdot \theta_1^2 - 3659 \cdot \theta_1 + 6275 \end{aligned}$$

$$0 = \frac{d}{d\theta_1} J(\theta_1) = 1070.25 \cdot \theta_1 - 3659$$

$$\theta_1 = \frac{3659}{1070.25} = 3.4188$$

(See <http://www.wolframalpha.com>)

Our First Algorithm In Action



The General Algorithm

In general, we don't want to plug numbers into $J(\theta_1)$ and solve a calculus problem *every time*.

The General Algorithm

In general, we don't want to plug numbers into $J(\theta_1)$ and solve a calculus problem *every time*.

Instead, we can solve for θ_1 in terms of $x^{(i)}$ and $y^{(i)}$.

The General Algorithm

In general, we don't want to plug numbers into $J(\theta_1)$ and solve a calculus problem *every time*.

Instead, we can solve for θ_1 in terms of $x^{(i)}$ and $y^{(i)}$.

The general problem: find θ_1 to minimize

$$J(\theta_1) = \frac{1}{2} \sum_{i=1}^m (\theta_1 x^{(i)} - y^{(i)})^2$$

You will solve this in HW1.

Two Problems Remain

Problem one: we only fit the slope. What if $\theta_0 \neq 0$?

Two Problems Remain

Problem one: we only fit the slope. What if $\theta_0 \neq 0$?

Problem two: we will need a better optimization algorithm than “Set $\frac{d}{d\theta}J(\theta) = 0$ and solve for θ .”

- ▶ Wiggly functions
- ▶ Equation(s) may be non-linear, hard to solve

Two Problems Remain

Problem one: we only fit the slope. What if $\theta_0 \neq 0$?

Problem two: we will need a better optimization algorithm than “Set $\frac{d}{d\theta}J(\theta) = 0$ and solve for θ .”

- ▶ Wiggly functions
- ▶ Equation(s) may be non-linear, hard to solve

Exercise: ideas for problem one?

Solution to Problem One

Design a cost function that takes two parameters:

$$\begin{aligned} J(\theta_0, \theta_1) &= \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2 \\ &= \frac{1}{2} \sum_{i=1}^m (\theta_0 + \theta_1 x^{(i)} - y^{(i)})^2 \end{aligned}$$

Find θ_0, θ_1 to minimize $J(\theta_0, \theta_1)$

Functions of multiple variables!

Here is an example cost function:

$$J(\theta_0, \theta_1) = \frac{1}{2}(\theta_0 + 17 \cdot \theta_1 - 63)^2 + \frac{1}{2}(\theta_0 + 19 \cdot \theta_1 - 65)^2 \\ + \frac{1}{2}(\theta_0 + 20.5 \cdot \theta_1 - 66)^2 + \frac{1}{2}(\theta_0 + 18.9 \cdot \theta_1 - 62.9)^2 + \dots$$

Gain intuition on <http://www.wolframalpha.com>

- ▶ Surface plot
- ▶ Contour plot

Solution to Problem Two: Gradient Descent

- ▶ **Gradient descent** is a general purpose optimization algorithm.
A “workhorse” of ML.

Solution to Problem Two: Gradient Descent

- ▶ **Gradient descent** is a general purpose optimization algorithm. A “workhorse” of ML.
- ▶ Idea: repeatedly take steps in steepest downhill direction, with step length proportional to “slope”

Solution to Problem Two: Gradient Descent

- ▶ **Gradient descent** is a general purpose optimization algorithm. A “workhorse” of ML.
- ▶ Idea: repeatedly take steps in steepest downhill direction, with step length proportional to “slope”
- ▶ Illustration: contour plot and pictorial definition of gradient descent

Gradient Descent

To minimize a function $J(\theta_0, \theta_1)$ of two variables

- ▶ Initialize θ_0, θ_1 arbitrarily
- ▶ Repeat until convergence

$$\theta_0 := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

$$\theta_1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$

- ▶ α = step-size or **learning rate** (not too big)

Partial derivatives

- ▶ The **partial derivative with respect to** θ_j is denoted $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
- ▶ Treat all other variables as constants, then take derivative

Partial derivatives

- ▶ The **partial derivative with respect to** θ_j is denoted $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
- ▶ Treat all other variables as constants, then take derivative
- ▶ Example

$$\frac{\partial}{\partial u} 5u^2v^3 =$$

Partial derivatives

- ▶ The **partial derivative with respect to** θ_j is denoted $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
- ▶ Treat all other variables as constants, then take derivative
- ▶ Example

$$\frac{\partial}{\partial u} 5u^2 v^3 = 5v^3 \frac{\partial}{\partial u} u^2$$

Partial derivatives

- ▶ The **partial derivative with respect to** θ_j is denoted $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
- ▶ Treat all other variables as constants, then take derivative
- ▶ Example

$$\begin{aligned}\frac{\partial}{\partial u} 5u^2 v^3 &= 5v^3 \frac{\partial}{\partial u} u^2 \\ &= 5v^3 \cdot 2u\end{aligned}$$

Partial derivatives

- ▶ The **partial derivative with respect to** θ_j is denoted $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
- ▶ Treat all other variables as constants, then take derivative
- ▶ Example

$$\begin{aligned}\frac{\partial}{\partial u} 5u^2 v^3 &= 5v^3 \frac{\partial}{\partial u} u^2 \\ &= 5v^3 \cdot 2u \\ &= 10v^3 u\end{aligned}$$

Partial derivatives

- ▶ The **partial derivative with respect to** θ_j is denoted $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$
- ▶ Treat all other variables as constants, then take derivative
- ▶ Example

$$\begin{aligned}\frac{\partial}{\partial u} 5u^2 v^3 &= 5v^3 \frac{\partial}{\partial u} u^2 \\ &= 5v^3 \cdot 2u \\ &= 10v^3 u\end{aligned}$$

$$\frac{\partial}{\partial v} 5u^2 v^3 = ??$$

Partial derivative intuition

Interpretation of partial derivative: $\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$ is the rate of change along the θ_j axis

Example: illustrate function with elliptical contours

- ▶ Sign of $\frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$?
- ▶ Sign of $\frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$?
- ▶ Which has larger absolute value?

Gradient Descent


- ▶ Repeat until convergence

$$\theta_0 = \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

$$\theta_1 = \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$

- ▶ Issues (explore in HW1)
 - ▶ Pitfalls
 - ▶ How to set step-size α ?
 - ▶ How to diagnose convergence?

The Result in Our Problem



$$h_{\theta}(x) = 39.75 + 1.25x$$

Gradient descent intuition

$$\theta_0 := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$$

$$\theta_1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$$

- ▶ Why does this move in the direction of steepest descent?
- ▶ What would we do if we wanted to maximize $J(\theta_0, \theta_1)$ instead?

Gradient descent for linear regression

Algorithm

$$\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) \quad \text{for } j = 0, 1$$

Cost function

$$J(\theta_0, \theta_1) = \sum_{i=1}^m \frac{1}{2} (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

We need to calculate partial derivatives.

Linear regression partial derivatives

Let's first do this with a single training example (x, y) :

$$\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) = \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2$$

Linear regression partial derivatives

Let's first do this with a single training example (x, y) :

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\ &= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y)\end{aligned}$$

Linear regression partial derivatives

Let's first do this with a single training example (x, y) :

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\ &= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y) \\ &= (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (\theta_0 + \theta_1 x - y)\end{aligned}$$

Linear regression partial derivatives

Let's first do this with a single training example (x, y) :

$$\begin{aligned}\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) &= \frac{\partial}{\partial \theta_j} \frac{1}{2} (h_{\theta}(x) - y)^2 \\ &= 2 \cdot \frac{1}{2} (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (h_{\theta}(x) - y) \\ &= (h_{\theta}(x) - y) \cdot \frac{\partial}{\partial \theta_j} (\theta_0 + \theta_1 x - y)\end{aligned}$$

So we get

$$\begin{aligned}\frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) &= (h_{\theta}(x) - y) \\ \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) &= (h_{\theta}(x) - y)x\end{aligned}$$

Linear regression partial derivatives

More generally, with many training examples (work this out):

$$\frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

$$\frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})x^{(i)}$$

Linear regression partial derivatives

More generally, with many training examples (work this out):

$$\frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

$$\frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) = \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})x^{(i)}$$

So the algorithm is:

$$\theta_0 := \theta_0 - \alpha \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

$$\theta_1 := \theta_1 - \alpha \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})x^{(i)}$$

Demo: parameter space vs. hypotheses

Show gradient descent demo

Summary

- ▶ What to know
 - ▶ Supervised learning setup
 - ▶ Cost function
 - ▶ Convert a learning problem to an optimization problem
 - ▶ Squared error
 - ▶ Gradient descent
- ▶ Next time
 - ▶ More on gradient descent
 - ▶ Linear algebra review