Assignment 2

Michael Lee MATH1853 University Number 3035569110

October 22, 2018

Answer Q. 1.

For eigenvalues of A, we have

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \tag{1}$$

with

$$\begin{bmatrix} a & b \\ b & -a \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}$$
$$= \begin{bmatrix} a - \lambda & b \\ b & -a - \lambda \end{bmatrix}$$

hence

$$\begin{vmatrix} a - \lambda & b \\ b & -a - \lambda \end{vmatrix} = 0$$

$$-(\alpha + \lambda)(\alpha - \lambda) - b^2 = 0 - a^2 + \lambda^2 - b^2$$

$$\lambda^2 = a^2 + b^2$$

$$\lambda = \pm \sqrt{a^2 + b^2}$$

For eigenvalue $\sqrt{a^2+b^2}$ and eigenvector v_1 of \mathbf{A} , we have

$$\begin{pmatrix} \mathbf{A} - \sqrt{a^2 + b^2} \mathbf{I} \end{pmatrix} \mathbf{v}_1 = 0$$

$$\begin{bmatrix} a + \sqrt{a^2 + b^2} & b \\ b & -a + \sqrt{a^2 + b^2} \end{bmatrix} \mathbf{v}_1 = 0$$

Hence we have

$$\begin{bmatrix} a + \sqrt{a^2 + b^2} & b & 0 \\ b & -a + \sqrt{a^2 + b^2} & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & \frac{b}{a + \sqrt{a^2 + b^2}} & 0 \\ 1 & \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & \frac{b}{a + \sqrt{a^2 + b^2}} - \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \\ 2 & \frac{b}{a + \sqrt{a^2 + b^2}} + \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & \frac{b(\sqrt{a^2 + b^2 - a})}{(a + \sqrt{a^2 + b^2})(\sqrt{a^2 + b^2 - a})} - \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \\ 2 & \frac{b(\sqrt{a^2 + b^2 - a})}{(a + \sqrt{a^2 + b^2})(\sqrt{a^2 + b^2 - a})} + \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & \frac{\sqrt{a^2 + b^2 - a}}{b} - \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \\ 2 & \frac{\sqrt{a^2 + b^2 - a}}{b} + \frac{-a + \sqrt{a^2 + b^2}}{b} & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & \frac{a + \sqrt{a^2 + b^2}}{b} & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

which can be reduced to