

MATH475: Combinatorics and Graph Theory

Michael Li

Contents

1	Chapter 1	2
1.1	The Basics - Permutations, Combinations, and General Counting	2
1.1.1	Permutation	2
1.1.2	Combinations	2
1.2	Counting in 2 Ways	5
1.3	Pigeon Hole Principle	6
1.4	Compositions and Set Partitions	6
1.5	Integer Partitions	9
1.6	Twelffold Way	10
1.7	Inclusion-Exclusion Principle	11
2	Chapter 2	13
2.1	Review of Power Series	13
2.1.1	OGF's and Recurrences	14
2.2	OGFs and Counting	15
2.3	Exponential Generating Functions	18
2.3.1	Compositions of EGF	23
2.4	Catalan Numbers	23
2.4.1	Combinatoric Proof of Catalan Numbers	24
3	Graph Theory	25
3.1	Introduction to Graph Theory	25
3.2	Vertex Degree	27
3.3	Properties of Trees	31
3.4	Spanning Trees	33
3.4.1	Rooted Plane Trees	35
3.4.2	Rooted Forest	36
3.4.3	Car Parking Problem	37

1 Chapter 1

1.1 The Basics - Permutations, Combinations, and General Counting

1.1.1 Permutation

Definition - Permutation: A **permutation** of an n -element set is an arrangement of the elements in a specific order

- k -permutation: arrangement of k elements from the set
- The total number of k -permutation of an n -element set is

$$P(n, k) = \frac{n!}{(n-k)!} = n(n-1) \cdots (n-k+1)$$

Example: 10 people can run for office for a committee with a President, Vice President, and Treasurer. The total number of possible committees is $P(10, 3)$

Definition - k th Falling Factorial of n : Let $n \geq k$, both positive integers. Then the **k th Falling Factorial of n** is $(n)_k = n(n-1) \cdots (n-k+1)$

Example - Permutations with Repetitions: How many rearrangements of $AAAABBBCCD$ are there?

- If we were only looking at distinct elements, there would be $10!$ factorial. However, the elements aren't distinct

As an example, look at how we can reorder the A 's

$$A_1 A_2 A_3 A_4 B B B C C D = A_2 A_1 A_3 A_4 B B B C C D = \cdots$$

Thus there are $4!$ of these representations that are equivalent and we divide $10!$ by $4!$ to account for this. Similar idea for the other letters

Thus the number of rearrangements is $\frac{10!}{4!3!2!1!}$

Theorem: Suppose object 1 occurs a_1 times, object 2 occurs a_2 times, ..., object k occurs a_k times. Furthermore, suppose $a_1 + \cdots + a_k = n$

Then the total number of arrangements is $\frac{n!}{a_1! \cdots a_k!} = \binom{n}{a_1, a_2, \dots, a_k}$

- **Definition - Multinomial Coefficients:** The notation above is called a multinomial coefficient

1.1.2 Combinations

Definition - Combinations: The total number of ways to create a k -element subset from $[n] = \{1, 2, \dots, n\}$ is denoted

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{P(n, k)}{k!}$$

- **Note:** the definition comes from $n!$ total possible distinct permutations of which we can permute a k -element subset $k!$ ways

Example: There are 5 cats, 5 dogs, and 4 mice. 3 are chosen at once

- How many total number of ways to get 2 cats, 1 dog?

$\binom{5}{2} \binom{5}{1}$. This comes from choosing 2 cats from 5 cats, and then choosing 1 dog from 5 dogs

- How many total ways to get at least 1 cat?

$\binom{14}{3} - \binom{9}{3}$. This comes from subtracting the possible groupings with no cats from the total number of possible groupings

$= \binom{5}{1}\binom{9}{2} + \binom{5}{2}\binom{9}{1} + \binom{5}{3}$. This comes from enumerating over possible groupings with 1 cat and 2 other animals, 2 cats and 1 other animal, and 3 cats and no other animals

Binomial Theorem: If n is a positive integer then

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

Proof: $(x + y)^n = (x + y) \cdots (x + y)$

The coefficients of $x^k y^{n-k}$ is the product of n terms distributing from k x -terms and $n - k$ y -terms

So out of the n terms that are multiplied together, $\binom{n}{k}$ contribute the x -term, leaving $n - k$ unused terms for y

Thus we have

$$\binom{n}{k} x^k \binom{n-k}{n-k} y^{n-k} = \binom{n}{k} x^k y^{n-k}$$

Multinomial Theorem: For $n \in \mathbb{Z}^+$, we have

$$(x_1 + \cdots + x_k)^n = \sum_{\substack{a_1 + \cdots + a_k = n \\ a_1, \dots, a_k \geq 0}} \binom{n}{a_1, a_2, \dots, a_k} x_1^{a_1} \cdots x_k^{a_k}$$

Proof: $(x_1 + \cdots + x_k)^n = (x_1 + \cdots + x_k) \cdots (x_1 + \cdots + x_k)$

The coefficients of $x_1^{a_1} \cdots x_k^{a_k}$ is the product of n terms resulting from a permutation with repetition of a_1, \dots, a_k

Thus there are $\binom{n}{a_1, \dots, a_k}$ distinct ways of representing the exponents of x_1, \dots, x_k

Worksheet Examples:

1. • How many strictly increasing 6-digit numbers are there (first digit can be 0)

Since the numbers are strictly increasing, they numbers must be distinct

Furthermore, there is only one possible way ordering of these selected digits (in increasing order), so order “doesn’t matter”

Thus we can just count the ways to select 6 digits: $\boxed{\binom{10}{6}}$

- What about when the first digit cannot be zero?

Since 0 can only appear as the first digit, this is identical to the first question, but restricted to $\{1, \dots, 9\}$

$\boxed{\binom{9}{6}}$ ways

2. A quiz has 5 problems with 3 different answers each. How many ways can the quiz be completed?

$\boxed{3^5}$ ways

3. • How many rearrangements of MISSISSIPPI are there?

$$\boxed{\binom{11}{4, 4, 2, 1}}$$

- How many rearrangements do not have all S's together?

Treat all S's as one unit $\Rightarrow \binom{8}{4, 2, 1, 1}$

Thus there are $\boxed{\binom{11}{4, 4, 2, 1} - \binom{8}{4, 2, 1, 1}}$ rearrangements

4. Suppose each digit in a 5-digit code can have any number zero to nine

- How many 5-digit codes are there with no restrictions?

$$\boxed{10^5}$$

- How many 5-digit codes have distinct digits?

$$\boxed{P(10, 5)}$$

- How many codes begin and end with an even number?

$$\boxed{5 * 10^4 * 5}$$

- How many 5-digit codes use exactly 2 different numbers?

$$\binom{5}{2} \text{ ways to choose two digits}$$

2^5 ways to write a five digit number using these two digits. However, we need to subtract the 2 cases where they use the same digit five times

Thus answer is $\boxed{\binom{5}{2}(2^5 - 2)}$

5. Consider a standard 52 card deck with 13 ranks and 4 suits

- Probability of getting a full house (triple of the same rank, pair of another rank)

$$\boxed{\frac{\binom{13}{1}\binom{4}{3}\binom{12}{1}\binom{4}{2}}{\binom{52}{5}}}$$

- Probability of getting 2 pairs of different ranks and a 5th card of a different rank

$$\boxed{\frac{\binom{13}{2}\binom{4}{2}\binom{4}{2} * 44}{\binom{52}{5}}}$$

Logic is to choose 2 types of ranks AT THE SAME TIME and then choosing possible pairs from those ranks. Finally multiply by 44 (the remaining number of possible cards for the 5th card)

Note: the answer below is NOT correct but is added to show an easy mistake to make

$$\frac{\binom{13}{1}\binom{4}{2}\binom{12}{1}\binom{4}{2} * 44}{\binom{52}{5}} \text{ is WRONG because we need to consider duplicate pairs (e.g. QQ, KK and KK, QQ)}$$

This solution ends up overcounting since we can choose Queens then Kings versus Kings then Queens, which yields the same hand

6. 15 people want to buy pre-made sandwiches. There are 7 H sandwiches, 4 C sandwiches, and 4 P sandwiches. If each person gets 1 sandwich, how many ways can the sandwiches be distributed?

$$\boxed{\binom{15}{7,4,4}}$$

7. 10 people are sitting at a circular table. How many seating arrangements are there?

$$\frac{10!}{10} = \boxed{9!}$$

1.2 Counting in 2 Ways

Given an equation, we count combinatorially in 2 ways

Example: $\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$

LHS: chooses a k -size subset from $[n]$

RHS: Partitions subsets into 2 cases, containing element 1 or not containing 1

- Containing 1: Pick $k-1$ elements from the remaining $n-1$ elements $\binom{n-1}{k-1}$
- Not containing 1: Pick k from $n-1$ elements $\binom{n-1}{k}$

Thus we have $\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$

Example: $\sum_{k=0}^n k \binom{n}{k} = n2^{n-1}$

RHS: Picks 1 element to be the leader (n choices)

Then from the remaining $n-1$ people, pick a subset (2^{n-1} ways)

LHS: Picks a group of size k ($\binom{n}{k}$ ways) and then select a leader amongst them (k possible choices)

Thus $\sum_{k=0}^n k \binom{n}{k} = n2^{n-1}$

Example: $\sum_{i=1}^n i(n-i+1) = \binom{n+2}{3}$

LHS: Let a, b, c be numbers with $a < b < c$ from $[n+2]$. Fix $b = i+1$ (the i th $+1$ element). Then there are i choices for the element a and $(n+2) - (i+1) = n-i+1$ choices for the element c

Thus the total ways to create a 3-size subset $\{a, b, c\}$ from $[n+2]$ is $\sum_{i=1}^n i(n-i+1) = \binom{n+2}{3}$

RHS: Counts the number of subsets of size 3 from $[n+2]$

Example: $\binom{2n}{n} = \sum_{k=0}^n \binom{n}{k}^2$

LHS: Count the number of size n subsets of $[2n]$

RHS: Can be written as $\binom{n}{k} \binom{n}{n-k}$

So choose k people from a group of size n and $n - k$ people from another group of size n

Thus $\binom{n}{k} \binom{n}{n-k} = \binom{2n}{n}$

1.3 Pigeon Hole Principle

Theorem - Pigeon Hole Principle: If $n + 1$ pigeons are placed into n pigeonholes, then there is at least one pigeonhole with at least 2 pigeons

Theorem - Generalized Pigeon Hole Principle: If n pigeons are placed into k holes, then at least one hole has

- At most $\lfloor \frac{n-1}{k} \rfloor$ pigeons (round down)
- At least $\lceil \frac{n}{k} \rceil$ pigeons (round up)
- **Note:** the statements above are equivalent

Proof: Suppose all holes have $\leq \lfloor \frac{n-1}{k} \rfloor$ pigeons

Then there are at most $k \lfloor \frac{n-1}{k} \rfloor \leq n-1 < n$ pigeons total. Which is a contradiction

Example: Prove that in a group of 40 people, at least 4 people have a same birthday in the same month

40 pigeons and 12 holes $\implies \lceil \frac{40}{12} \rceil = 4$. Thus by PHP, the statement holds

Example: Prove that if 6 distinct numbers are chosen from $[9]$, then 2 numbers will sum to 10

6 pigeons and 5 holes $(\{(1,9), \dots, (5,5)\} \implies \lceil \frac{6}{5} \rceil = 2$. Thus by PHP, 2 of the numbers belong to the same pair and sum to 10

Example; An athlete wants to work out 45 hours over a 30-day month and will work out at least 1 hour each day. Prove that there is a period of consecutive days such that the cumulative hours he has worked out is exactly 14

Let a_i be the total hours accumulated up to day i

This creates a strictly increasing sequence $a_1, a_2, \dots, a_{30} = 45$ where each a_i is distinct

Define a new sequence $b_i = a_i + 14$

This creates a strictly increasing sequence $b_1, b_2, \dots, b_{30} = 59$ where each b_i is distinct

Since $\{a_i\}_{i=1}^{30}$ is distinct and $\{b_i\}_{i=1}^{30}$ is distinct (only within their own respective sequence), we can have at most 60 possible values, which we treat as pigeons

Furthermore, b_i can take on any of 59 distinct values, which we treat as the holes

Thus by PHP, some $a_j = b_k = a_k + 14$

Thus from day $k + 1$ to day j , the total number of hours worked out is 14

1.4 Compositions and Set Partitions

Definition - Weak Composition: Take $a_1, \dots, a_k \geq 0$ such that $\sum_{i=1}^k a_i = n$. Then the ordered k -tuple (a_1, \dots, a_k) is a **weak composition** of n into k partitions

- **Note:** This can be seen as distributing n unlabelled balls into k labelled boxes

Theorem - Stars and Bars: The number of weak compositions of n into k parts is $\binom{n+k-1}{k-1}$

Proof:

Denote the n objects as stars. We model a distribution as follows:

Place k bars where stars to the right of each bar is how many stars the corresponding part contains

Note: we must begin with a bar. Thus the rest involves n stars and $k - 1$ bars

Thus we can choose $k - 1$ from these $n + k - 1$ objects, resulting in $\binom{n + k - 1}{k - 1}$

Each one of these combination generates a different weak composition

Example: How many ways can we distribute \$100 to 5 people?

- Can also be rephrased as how many non-negative integer solutions are there to $x_1 + \dots + x_5 = 100$

Using the Stars and Bars Theorem, we have $\binom{100 + 5 - 1}{5 - 1} = \binom{104}{4}$ ways

Example: Suppose we want to choose 3 lollipop flavors from 5 orange, 4 strawberry, and 3 grapes. How many ways can we choose 3 lollipops?

Here we want $x_O + x_S + x_G = 3$ where $x_O, x_S, x_G \geq 0$

Thus $\binom{3 + 3 - 1}{3 - 1}$

Definition - Compositions: Same as weak compositions but with the restriction that each $x_i \geq 1$

We can derive this with the formula above but ensuring that each of the k boxes starts with 1 star, and then we distribute the remaining $n - k$ stars in a weak composition fashion. Thus the number of compositions of n into k parts is

$$\binom{n - k + k - 1}{k - 1} = \binom{n - 1}{k - 1}$$

Note: This problem is placing unlabelled balls into labelled bins

Definition - Partition: Let X be a finite set and A_1, \dots, A_k be non-empty subsets of X . Then a **partition** of X , $\{A_1, \dots, A_k\}$ satisfies

- $\bigcup_{i=1}^k A_i = X$
- Each A_i, A_j are pairwise disjoint

Note: This problem can be stated as: for a given k , X is partitioned into k **blocks**

Example: $\{\{1, 2, 4\}, \{3, 5\}, \{6\}\}$ is a partition of $[6]$

The first inner set here can be considered placing balls 1, 2, 4 into its own (unlabelled) bin

Definition - Stirling Number of the 2nd Kind: The number of partitions of $[n]$ into k **blocks**, denoted $S(n, k)$

- $S(0, 0) = 1$ by convention
- $S(n, 0) = 0$ for $n \geq 1$ is a partition of $[6]$

Example: Use counting arguments to compute $S(n, 2)$

For each ball, we decide if it either it goes into block 1 or block 2, resulting in 2^n

However, we require that each block has ≥ 1 elements. Thus we need to subtract by 2 for the cases where block 1 has all balls or block 2 has all balls by 2 for the cases where block 1 has all balls or block 2 has all balls

Finally, we need to divide by 2 since the order of the blocks doesn't matter

$$\frac{2^n - 2}{2}$$

Theorem: For $n \geq k$, we have

$$S(n, k) = S(n-1, k-1) + kS(n-1, k)$$

Proof:

LHS: Counts the number of ways to partition $[n]$ into k blocks

RHS looks at partitions where

- $\{1\}$ is in its own bin. Then we need to partition $[n-1]$ into $k-1$ bins, resulting in $S(n-1, k-1)$
- $\{1\}$ is in a bin with other elements. Then we need to partition $[n-1]$ into k bins, and then choose a bin to place $\{1\}$ into, resulting in $kS(n-1, k)$

Theorem:
$$S(n, k) = \frac{1}{k!} \sum_{i=0}^k (-1)^i \binom{k}{i} (k-i)^n$$

Proof: will show later

Definition - Bell Number: Total number of partitions of $[n]$, denoted $B(n) = \sum_{k=1}^n S(n, k)$

- **Note:** $B(0) = 1$ by convention

Example: Take $n = 3$. Then we have

$$\{\{1, 2, 3\}\}, \{\{1\}, \{2\}, \{3\}\}, \{\{1\}, \{2, 3\}\}, \{\{2, 1, 3\}\}, \{\{3\}\{1, 2\}\}$$

Thus $B(3) = 5$

Theorem:
$$B(n) = \sum_{i=1}^n \binom{n-1}{i-1} B(n-i)$$

Proof:

LHS: counts the total number of partitions of $[n]$

RHS:

- Fix the n th element and determine the number of subsets of size i that contain this element: $\binom{n-1}{i-1}$ ways
- Partition the remaining $n-i$ elements: $B(n-i)$ ways

Thus we see that
$$B(n) = \sum_{i=1}^n \binom{n-1}{i-1} B(n-i)$$

Example: Consider $f: A \rightarrow B$ where $|A| = n$ and $|B| = k$ for $n \geq k$. What is the total number of surjective functions?

- Recall that f is surjective if for $\forall b \in B$, there exists at least one $a \in A$ such that $f(a) = b$

This problem can be viewed as partitioning A into $\{A_1, \dots, A_k\}$ and determining how they map to B , where $A_i = \{a \in A \mid f(a) = i\}$

Clearly the number of ways to partition $[n]$ into k parts is $S(n, k)$

However, we also need to consider the order of the boxes (since they determine which subsets map to which values of B). Thus we need to multiply by $k!$

Thus the final answer is $k!S(n, k)$

1.5 Integer Partitions

Definition - Partition of Integer n : Positive sequence of integers a_1, \dots, a_k , with $a_1 \geq a_2 \geq \dots \geq a_k$ such that

$$a_1 + a_2 + \dots + a_k = n$$

Total number of such sequences is denoted $p(n)$

- **Note:** Observe that since the values are non-increasing, the order “doesn’t matter” so we don’t need to distinguish $1 + 3 = 4$ from $3 + 1 = 4$
- **Note:** By convention, we have $p(0) = 1$


Example: Take $n = 4$. Then we have

$$4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 = 1 + 1 + 1 + 1$$

Thus $p(4) = 5$

Definition - Ferrers Diagram: Represents a partition (a_1, \dots, a_k) by taking a partial rectangular grid with k rows whose rows contain a_i dots, such that $a_1 \geq \dots \geq a_k$

Example:




This corresponds to $4 + 3 + 1 + 1 + 1 = 10$

Definition - Conjugate of a Partition: Diagram created by reflecting a Ferrers Diagram along the diagonal

- **Note:** Can be seen as replacing columns as rows and rows as columns

Example: Conjugate of the previous example




This corresponds to $5 + 2 + 2 + 1 = 10$

Theorem: Let P_1 denote the number of partitions where $a_1 = a_2 \geq a_3 \geq \dots \geq a_k$ (for unfixed k). Let P_2 be the number of partitions whose smallest part is at least 2. Then

$$P_1 = P_2$$

Proof: \implies Take a partition where $a_1 = a_2$. Then take its conjugate.



Clearly every part of the conjugate has length ≥ 2

Adding more dots to the original diagram doesn't affect the minimal lengths of the conjugate

Thus the conjugate assures each row has at least 2 dots \implies each part has at least size 2

\Leftarrow Given a partition with $a_k \geq 2$, the conjugate creates a partition with 2 rows that are the largest and have the same size. Thus $a_1 = a_2$ in the conjugate

Example: Show that the total number of partitions of n containing a part of size 1 is $p(n-1)$

\implies Take partition P of n containing a part of size 1

This part will be the bottom row. Deleting that row yields $p(n-1)$

\Leftarrow Given a partition P_2 of $n-1$, add a dot to the bottom. This yields a partition of n with a part of size 1

Definition - $p_k(n)$: Number of partitions of n into exactly k parts

Theorem: For $1 < k < n$

$$p_k(n) = p_{k-1}(n-1) + p_k(n-k)$$

Proof:

LHS: counts the number of partitions of n into k parts

RHS:

- Count the number of partitions such that there is a partition of size 1: $p_{k-1}(n-1)$
- Count the number of partitions such that every part has size ≥ 1 : $p_k(n-k)$

Note: Each part must have size ≥ 1 by definition

1.6 Twelfold Way

Goal is to distribute n (labelled or unlabelled) balls into k (labelled or unlabelled) bins, depending on if the bins

- Have no restriction
- Have at least 1 ball
- Have at most 1 ball

This results in 12 possible scenarios

1. n Labelled Balls into k Labelled Bins

- No restrictions: For each ball, pick a bin to put it into $\implies k^n$
- At least 1: Can be viewed as the number of surjective functions $\implies k!S(n, k)$
- At most 1:
 - If $n > k \implies 0$
 - Otherwise $n \leq k$ and we pick n bins in a specific order $\implies P(k, n)$

2. n Unlabelled Balls into k Labelled Bins

- No restrictions: Weak composition $\implies \binom{n+k-1}{k-1}$
- At least 1: Composition $\implies \binom{n-1}{k-1}$
- At most 1:

- If $n > k \implies 0$
- Otherwise $n \leq k$ and we just need to choose n bins to use $\implies \binom{k}{n}$

3. n Labelled Balls into k Unlabelled Bins

- No restrictions: Need to count all possible partitions with 0 to k parts $\implies \sum_{i=1}^k S(n, i)$
 - **Note:** This is NOT Bell's Number since we might have $k < n$
- At least 1: Partition $[n]$ into k parts $\implies S(n, k)$
- At most 1:
 - If $n > k \implies 0$
 - Otherwise $n \leq k$ and the bins are indistinguishable $\implies 1$ way

4. n Unlabelled Balls into k Unlabelled Bins

- No restrictions: need to count all possible partitions with 0 to k parts $\implies \sum_{i=1}^k p_i(n)$
- At least 1: Partition n into k parts $\implies p_k(n)$
- At most 1:
 - If $n > k \implies 0$
 - Otherwise $n \leq k$ and the bins are indistinguishable $\implies 1$ way

Definition - Multiset: Set of elements such that elements may repeat

Example: How many multisets of size 5 can be created using [7]?

Weak compositions for $1, 2, \dots, 7$ such that the number of balls in each bin totals up to 5 $\implies \binom{5+7-1}{7-1}$

Example: How many injective functions $f : [n] \rightarrow [k]$ are there?

n labelled balls into k labelled bins where each bin gets at most 1 $\implies P(k, n)$

Example: 20 kids split into 4 study groups. 2 brothers must be in the same group and 2 sisters must be in the same group. What is the total number of ways to split the kids

Treat the brothers and sisters each as one unit then partition $[18]$ into 4 bins $\implies S(18, 4)$

1.7 Inclusion-Exclusion Principle

Theorem: Let $A_1, \dots, A_n \subseteq X$ where X is finite and each A_i is non-empty. Also let

- $I \subseteq [n]$
- $A_I = \bigcap_{i \in I} A_i$ with $A_\emptyset = X$

Then

$$\left| \bigcup_{i=1}^n A_i \right| = |X| - \sum_{I \subseteq [n]} (-1)^{|I|} |A_I|$$

Note: By De Morgan's Law, this is equivalent to

$$\left| \bigcap_{i=1}^n \bar{A}_i \right| = |X| - \sum_{I \subseteq [n]} (-1)^{|I|} |A_I|$$

Example: For $n = 2 \implies I = \emptyset, \{1\}, \{2\}, \{1, 2\}$

$$\begin{aligned}
\left| \bigcup_{i=1}^2 A_i \right| &= |X| - ((-1)^0 |X| + (-1)^1 |A_1| + (-1)^1 |A_2| + (-1)^2 |A_{\{1,2\}}|) \\
&= |A_1| + |A_2| - |A_1 \cap A_2|
\end{aligned}$$

Example: How many integer solutions satisfy

$$x_1 + x_2 + x_3 = 35 \quad \text{for } 0 \leq x_1, x_2, x_3 \leq 15$$

Consider the problem where $X_1 \geq 16$. We can place 16 balls into the first bin then perform weak composition. We now use the negation form of the theorem above

Let X be the set of all weak compositions. Now define

- $A_1 = \{ \text{weak compositions} \mid x_1 \geq 16 \}$
- $A_2 = \{ \text{weak compositions} \mid x_2 \geq 16 \}$
- $A_3 = \{ \text{weak compositions} \mid x_3 \geq 16 \}$

Now we see that

$$\begin{aligned}
|\bar{A}_1 \cap \bar{A}_2 \cap \bar{A}_3| &= \sum_{I \subseteq [3]} (-1)^{|I|} |A_I| \\
&= \underbrace{\binom{35+3-1}{3-1}}_{I=\emptyset} - \underbrace{\binom{3}{1} \binom{35-16+3-1}{3-1}}_{I=\{1\}, \{2\}, \{3\}} + \underbrace{\binom{3}{2} \binom{35-32+3-1}{3-1}}_{I=\{1,2\}, \{1,3\}, \{2,3\}} - \underbrace{\binom{3}{3} (0)}_{\text{impossible since } 48 > 35}
\end{aligned}$$

Example: Count the total number of surjective functions $f : [n] \rightarrow [k]$, where $n \geq k$

Let X be all functions from $[n] \rightarrow [k]$ and let A_i be the set of functions that DO NOT map to $i \in [k]$. Then the total number of surjective functions is

$$\begin{aligned}
\left| \bigcap_{i=1}^k \bar{A}_i \right| &= \sum_{I \subseteq [k]} (-1)^{|I|} |A_I| \\
&= \sum_{i=0}^k \binom{k}{i} (-1)^i (k-i)^n \quad \text{out of } k, \text{ choose a set of image points not mapped to} \\
&= k! S(n, k)
\end{aligned}$$

Example: Consider the Euler Totient Function

$$\phi(n) = |\{i \in \mathbb{Z}^+ \mid 1 \leq i \leq n, \gcd(i, n) = 1\}|$$

Let $p_1^{e_1} p_2^{e_2} \cdots p_t^{e_t}$, for $e_i \geq 1$ be the prime factorization

Let $A_i = \{a \in [n] \mid p_i \mid a\}$. Then we see value of $\phi(n)$ is

- Note $|A_1| = \frac{n}{p_1}$ how many numbers in $[n]$ have 2 as a factor
- Note $|A_{\{1,2\}}| = \frac{n}{p_1 p_2}$ how many numbers in $[n]$ have 2 AND 3 as factors

$$\begin{aligned}
\left| \bigcap_{i=1}^t \bar{A}_i \right| &= \sum_{I \subseteq [t]} (-1)^{|I|} |A_I| \\
&= \underbrace{n}_{I=\emptyset} - \left(\frac{n}{p_1} + \frac{n}{p_2} + \cdots + \frac{n}{p_t} \right) + \left(\frac{n}{p_1 p_2} + \frac{n}{p_2 p_3} + \cdots \right) - \left(\frac{n}{p_1 p_2 p_3} + \cdots \right) + \cdots \\
&= n \left(1 - \sum_{i=1}^t \frac{1}{p_i} + \sum_{1 \leq i \leq j \leq t} \frac{1}{p_i p_j} - \sum_{1 \leq i \leq j \leq k} \frac{1}{p_i p_j p_k} \right) \\
&= n \left(1 - \frac{1}{p_1} \right) \left(1 - \frac{1}{p_2} \right) \cdots \left(1 - \frac{1}{p_t} \right) \\
&= n \prod_{i=1}^t \left(1 - \frac{1}{p_i} \right)
\end{aligned}$$

Example: How many ways to seat 20 couples in a circular, unlabelled table with 40 seats such that each significant other doesn't sit with each other

Let A_i be the set of seatings where couple 1 is sitting next to each other. Then we see

$$\begin{aligned}
\left| \bigcap_{i=1}^{20} \bar{A}_i \right| &= \sum_{I \subseteq [20]} (-1)^{|I|} |A_I| \\
&= \sum_{i=0}^{20} \underbrace{\binom{20}{i}}_{\text{select couples}} (-1)^i \underbrace{\frac{(40-i)!}{40-i}}_{\text{glue couples}} \overbrace{2^i}^{\text{order of couples}}
\end{aligned}$$

2 Chapter 2

2.1 Review of Power Series

Definition - Formal Power Series: For a sequence $\{a_n\}_{n \geq 0}$, the **Formal Power Series**

$$F(x) = \sum_{n=0}^{\infty} a_n x^n$$

Is the **Ordinary Generating Function** of $\{a_n\}_{n \geq 0}$

Example: Let $\{a_n\}$ be given by

$$\frac{1}{2!}, \frac{1}{3!}, \dots$$

Then the generating function is

$$\begin{aligned}
F(x) &= \frac{1}{2!} + \frac{1}{3!}x + \frac{1}{4!}x^2 + \cdots \\
&= \frac{1}{x^2} \left(\frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \cdots \right) \\
&= \frac{1}{x^2} \sum_{n=2}^{\infty} \frac{x^n}{n!} \\
&= \frac{1}{x^2} (e^x - x - 1)
\end{aligned}$$

Same principle can be applied to find a sequence $\{a_n\}_{n \geq 0}$ given an OGF $F(x)$

Example: Find the sequence $\{a_n\}$ corresponding to

$$F(x) = \sum_{n=0}^{\infty} a_n x^n = \frac{-4x + 3}{(1-x)(1-2x)}$$

Using partial fractions we see

$$\frac{A}{1-x} + \frac{B}{1-2x} = -4x + 3 = A(1-2x) + B(1-x) \implies A = 1, B = 2$$

Thus we see that

$$\begin{aligned} F(x) &= \frac{-4x + 3}{(1-x)(1-2x)} \\ &= \frac{1}{1-x} + \frac{2}{1-2x} \\ &= \sum_{n=0}^{\infty} x^n + 2 \underbrace{\sum_{n=0}^{\infty} (2x)^n}_{\sum 2^{n+1} x^n} \\ &= \sum_{n=0}^{\infty} (1 + 2^{n+1}) x^n \end{aligned}$$

Thus $\{a_n\}_{n \geq 0} = \{1 + 2^{n+1}\}$

2.1.1 OGF's and Recurrences

We can replace a_n with recursions for OGFs $\sum a_n x^n$

Example: Consider codes of length n using A, B, C, D

Let a_n be the total codes containing an even number of A's

1. Find a recurrence for a_n in terms of a_{n-1}

$$a_n = 3a_{n-1} + 4^{n-1} - a_{n-1}$$

This comes from

- Code doesn't end in A $\implies a_n = 3a_{n-1}$ (there are 3 possible letters for the last letter)
- Code ends in A $\implies 4n^{n-1} - a_{n-1}$ (All codes of length $n-1$ minus codes of length $n-1$ with an even number of A)

2. Find the OGF for $\{a_n\}$

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} a_n x^n = \underbrace{1}_{a_0} + \sum_{n=1}^{\infty} a_n x^n \\ &= 1 + \sum_{n=1}^{\infty} (2a_{n-1} + 4^{n-1}) x^n \\ &= 1 + \sum_{n=1}^{\infty} 2a_{n-1} x^n + \sum_{n=1}^{\infty} 4^{n-1} x^n \\ &= 1 + 2x \underbrace{\sum_{n=1}^{\infty} a_{n-1} x^{n-1}}_{f(x)} + x \underbrace{\sum_{n=1}^{\infty} (4x)^{n-1}}_{\text{geometric sum}} \\ &= 1 + 2x(f(x) + x(\frac{1}{1-4x})) \end{aligned}$$

Thus $f(x) = \frac{1}{1-2x} + \frac{x}{(1-2x)(1-4x)}$

3. Find the closed form of a_n

By partial fractions, we have

$$\begin{aligned} f(x) &= \frac{1}{1-2x} + \frac{-1}{2(1-2x)} + \frac{1}{2(1-4x)} \\ &= \frac{1}{2} \left(\sum_{n=0}^{\infty} (2x)^n \right) + \frac{1}{2} \sum_{n=0}^{\infty} (4x)^n \\ &= \sum_{n=0}^{\infty} \underbrace{\left(\frac{2^n}{2} + \frac{4^n}{2} \right)}_{a_n} x^n \end{aligned}$$

Thus we have $\{a_n\} = \{2^{n-1} + \frac{4^n}{2}\}$

2.2 OGFs and Counting

Using $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$, we can differentiate to get

$$\begin{aligned} \frac{1}{(1-x)^2} &= \sum_{n=1}^{\infty} n x^{n-1} = 1 + 2x + \cdots = \sum_{n=0}^{\infty} (n+1) x^n \\ \frac{2}{(1-x)^3} &= \sum_{n=2}^{\infty} n(n-1) x^{n-2} = 1 + 2x + \cdots = \sum_{n=0}^{\infty} (n+2)(n+1) x^n \\ &\vdots \\ \frac{m!}{(1-x)^{m+1}} &= \sum_{n=0}^{\infty} \underbrace{(n+m)(n+m-1) \cdots (n+1)}_{(n+m)_m} x^n = \sum_{n=0}^{\infty} \frac{(n+m)!}{n!} x^n \end{aligned}$$

Thus we have

$$\frac{1}{(1-x)^{m+1}} = \sum_{n=0}^{\infty} \frac{(n+m)!}{n!m!} x^n = \sum_{n=0}^{\infty} \binom{n+m}{m} x^n$$

For a fixed m , we have

$$\frac{1}{(1-x)^{m+1}} \iff \{a_n\} \quad a_n = \binom{n+m}{m}$$

Which is the OGF for weak compositions into $m+1$ bins. However, we can now interpret

$$\frac{1}{(1-x)^k} = \underbrace{\left(\frac{1}{1-x} \right) \left(\frac{1}{1-x} \right) \cdots}_{\text{apply geometric series k times}} = (1+x+x^2+\cdots)(1+x+x^2+\cdots)\cdots$$

Consider the question of getting coefficients of x^7 . This corresponds to the weak compositions of 7 unlabelled balls into k bins. However, now the chosen exponents correspond to the bin sizes. Thus we see that the OGF for weak compositions is

$$F(x) = \frac{1}{(1-x)^k} = \sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} \binom{n+k-1}{k-1} x^n$$

To obtain coefficients of a_n , contributions come from x^{n_1} from group 1, x^{n_2} from group 2, etc. such that

$$x^{n_1} x^{n_2} \cdots x^{n_k} = x^n \implies n_1 + n_2 + \cdots + n_k = n \quad n_i \geq 0$$

Example: 5 apple, 4 pear, and 6 orange lollipops. Find the OGF that would yield the total ways to get 5 lollipops. State the term that we want

$$F(x) = \underbrace{(1 + x + \cdots + x^5)}_{\text{apples}} \underbrace{(1 + x + x^4)}_{\text{pears}} \underbrace{(1 + x + \cdots + x^6)}_{\text{oranges}}$$

We want coefficients of x^5 in $F(x)$. Here the OGF is

$$\frac{1}{(1-x)^k} \longleftrightarrow \left\{ \binom{n+k-1}{k-1} \right\}$$

Theorem: Let k be a fixed positive integer. Then the OGF for Stirling Numbers $S(n, k)$ is

$$F_k(x) = \sum_{n=0}^{\infty} a_n x^n = \frac{x^k}{(1-x)(1-2x) \cdots (1-kx)}$$

Proof: Note that

$$S(0, 0) = 1 \quad S(n, 0) = 0 \quad S(0, k) = 0$$

Also recall the recurrence

$$S(n, k) = S(n-1, k-1) + kS(n-1, k)$$

Then we see that

$$\begin{aligned} F_k(x) &= \sum_{n=0}^{\infty} S(n, k) x^n = 0 + \sum_{n=1}^{\infty} (S(n-1, k-1) + kS(n-1, k)) x^n \\ &= \left(x \sum_{n=1}^{\infty} S(n-1, k-1) x^{n-1} \right) + \left(kx \sum_{n=1}^{\infty} S(n-1, k) x^{n-1} \right) \\ &= xF_{k-1}(x) + kxF_k(x) \end{aligned}$$

Thus $F_k(x) = \frac{x F_{k-1}(x)}{(1-kx)}$ Finally, applying the above equation recursively, we see that

$$\begin{aligned} F_k(x) &= \frac{x}{1-kx} \frac{x}{1-(k-1)x} F_{k-2} = \cdots \\ &= \frac{x^k}{(1-x)(1-2x) \cdots (1-kx)} \end{aligned}$$

Thus the OGF is

$$\prod_{t=1}^k \frac{x}{(1-tx)} \longleftrightarrow \{S(n, k)\}$$

Theorem: Let $p(n)$ denote the total partitions of an integer n . Then the OGF for $p(n)$ is

$$F(x) = \sum_{n=0}^{\infty} p(n) x^n = \frac{1}{(1-x)(1-x^2) \cdots} = \prod_{k=1}^{\infty} \frac{1}{1-x^k}$$

Proof:

$$\begin{aligned} F(x) &= \frac{1}{1-x} \frac{1}{1-x^2} \frac{1}{1-x^3} \cdots \\ &= \underbrace{(1 + x + x^2 + x^3 + \cdots)}_{\text{how many 1's}} \underbrace{(1 + x^2 + x^4 + x^6 + \cdots)}_{\text{how many 2's}} \underbrace{(1 + x^3 + x^6 + x^9 + \cdots)}_{\text{how many 3's}} \cdots \\ &= \prod_{k=1}^{\infty} (1 + x^k + x^{2k} + \cdots) \end{aligned}$$

Consider the question: how many combinations of x^n ?

For $x^{m(k)*k}$, $m(k)$ is the total number of k 's in the partition. For example, $m(1)$ is the total 1's in the partition

All possible contributions of x^n is exactly the total partitions of integer n

Example: Given an unlimited number of pennies, nickels, dimes, quarters, and half dollars, find the OGF that would count the total ways to make \$1. What is the coefficient?

We want coefficients of x^{100} where

$$F(x) = \underbrace{(1 + x + \cdots + x^{100})}_{\text{number of pennies}} \underbrace{(1 + x^5 + \cdots + x^{100})}_{\text{number of nickels}} \underbrace{(1 + x^{10} + \cdots + x^{100})}_{\text{number of dimes}} \underbrace{(1 + x^{25} + \cdots + x^{100})}_{\text{number of quarters}} \underbrace{(1 + x^{50} + x^{100})}_{\text{number of half dollars}}$$

Expanding (done on computer) yields $292x^{100} \implies 292$ ways

Theorem: Total ways to partition n into distinct parts equals the total ways to partition n into odd sized parts

Proof: Distinct parts OGF is

$$\begin{aligned} F(x) &= (1+x)(1+x^2) + \cdots && \text{this comes from each number only appears once} \\ &= \left(\frac{1-x^2}{1-x}\right)\left(\frac{1-x^4}{1-x^2}\right) \cdots \\ &= \left(\frac{1}{1-x}\right)\left(\frac{1}{1-x^3}\right)\left(\frac{1}{1-x^5}\right) \cdots \\ &= (1+x+x^2+\cdots)(1+x^3+x^6+\cdots) \end{aligned}$$

From above, we see that the even terms get cancelled and we are only left with the odd terms

Thus number of ways to partition n into distinct parts equals the total number of ways to partition n into odd sized parts

Theorem: OGF for $p_k(n)$, the total number of partitions into exactly k parts is

$$F(x) = \sum_{n=0}^{\infty} p_k(n)x^n = \frac{x^k}{(1-x)(1-x^2)\cdots(1-x^k)} = (1+x+x^2+\cdots)(1+x^2+x^4+\cdots)\cdots(x^k+x^{2k}+\cdots)$$

Proof: Consider the OGF

$$F(x) = \frac{1}{(1-x)(1-x^2)\cdots(1-x^k)} = (1+x+x^2+\cdots)(1+x^2+x^4+\cdots)\cdots(1+x^k+x^{2k}+\cdots)$$

This counts the number of partitions of n where each part has size at most k

Looking at the Ferrer Diagram of this, we see that each row can have at most k dots

Thus the conjugate will have columns of size at most k , which establishes a bijection between counting the total partitions of n with at most k part

To manipulate this into an OGF for $p_k(n)$, we need a exactly k rows in the conjugate, and thus need a part of exactly size k in the original diagram

Thus we can create a new OGF that enforces an x^k term

$$F(x) = \frac{x^k}{(1-x)(1-x^2)\cdots(1-x^k)} = (1+x+x^2+\cdots)(1+x^2+x^4+\cdots)\cdots(x^k+x^{2k}+\cdots)$$

2.3 Exponential Generating Functions

Definition - EGF: Function for a sequence $\{a_n\}$

$$G(x) = \sum_{n=0}^{\infty} a_n \frac{x^n}{n!}$$

Recall that for OGFs $A(x) = \sum a_n x^n$ and $B(x) = \sum b_n x^n$, we have that

$$A(x)B(x) = \sum_{n=0}^{\infty} c_n x^n \quad c_n = \sum_{i=0}^n a_i b_{n-i} \implies \sum_{n=0}^{\infty} c_n x^n = (a_0 + b_0) + (a_0 b_1 + a_1 b_0)x + (a_0 b_2 + a_1 b_1 + a_2 b_0)x^2 + \dots$$

Lemma: For OGFs $F(x) = \sum \bar{a}_n \frac{x^n}{n!}$ and $G(x) = \sum \bar{b}_n \frac{x^n}{n!}$, we have that

$$F(x)G(x) = \sum_{n=0}^{\infty} \bar{c}_n \frac{x^n}{n!} \quad \bar{c}_n = \sum_{i=0}^n \binom{n}{i} \bar{a}_i \bar{b}_{n-i}$$

Proof: let $a_n = \frac{\bar{a}_n}{n!}$ and $b_n = \frac{\bar{b}_n}{n!}$. Then we have

$$F(x)G(x) = \sum_{n=0}^{\infty} c_n x^n \quad c_n = \sum_{i=0}^n a_i b_{n-i} = \frac{1}{n!} \sum_{i=0}^n \binom{n}{i} \bar{a}_i \bar{b}_{n-i}$$

This last part is apparent from

$$\begin{aligned} c_n &= \sum_{i=0}^n \frac{\bar{a}_i}{i!} \frac{\bar{b}_{n-i}}{(n-i)! n} \\ &= \frac{1}{n!} \sum_{i=0}^n \frac{n!}{i!(n-i)!} \bar{a}_i \bar{b}_{n-i} \\ &= \frac{1}{n!} \sum_{i=0}^n \binom{n}{i} \bar{a}_i \bar{b}_{n-i} \end{aligned}$$

Example: For a fixed m , let $\{a_n\} = \left\{ \binom{m}{n} \right\}$. Find the EGF for this

We know that the OGF for $\{a_n\}$ is

$$(1+x)^m = \sum_{n=0}^{\infty} \binom{m}{n} x^n = \sum_{n=0}^{\infty} \frac{m!}{n!(m-n)!} x^n$$

Thus this gives

$$\sum_{n=0}^{\infty} \frac{m!}{(m-n)!} \frac{x^n}{n!} = \sum_{n=0}^{\infty} P(m, n) \frac{x^n}{n!}$$

Thus the EGF is $\{P(m, n)\}$

Example: For a fixed k the EGF for $\{k^n\}$ is

$$\sum_{n=0}^{\infty} k^n \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{(kx)^n}{n!} = e^{kx}$$

Theorem: For a fixed k , the EGF for Stirling Numbers $S(n, k)$ is

$$G(x) = \sum_{n=0}^{\infty} S(n, k) \frac{x^n}{n!} = \frac{(e^x - 1)^k}{k!}$$

Corollary: $B(n) = \sum_{k=0}^n S(n, k)$. Then the EGF is

$$G(x) = \sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} = e(e^x - 1)$$

Proof: We know that

$$G(x) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{\infty} S(n, k) \right) \frac{x^n}{n!}$$

- **Note:** If $n < k$, then $S(n, k) = 0$

Applying the previous theorem, we see that

$$\begin{aligned} G(x) &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^{\infty} S(n, k) \right) \frac{x^n}{n!} \\ &= \sum_{k=0}^{\infty} \frac{1}{k!} (e^x - 1)^k \\ &= e^{(e^x - 1)} \quad \text{Note : } e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \end{aligned}$$

Consider the EGF of permutations with repetitions of S_1, S_2, S_3

$$\left(\sum_{\substack{S_1 + S_2 + S_3 = n \\ S_1, S_2, S_3 \geq 0}} \frac{n!}{S_1! S_2! S_3!} \right) \frac{x^n}{n!}$$

What is the coefficient of all possible permutations of length n using A, B, C?

Combinatorically, we know that this is 3^n . Thus we see that the EGF is

$$G(x) = \sum_{n=0}^{\infty} 3^n \frac{x^n}{n!} = e^{3x}$$

Example: Find the EGF that counts the total number of permutations of length n containing an odd number of As, even number of Bs, and any number of Cs. Then find a closed form for the value

$$\begin{aligned}
& \underbrace{\left(x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots\right)}_A \underbrace{\left(1 + \frac{x^2}{2!} + \dots\right)}_B \underbrace{\left(1 + x + \frac{x^2}{2!} + \dots\right)}_C \\
&= \left(\left(1 + x + \frac{x^2}{2!} + \dots\right) - \left(1 - x + \frac{(-x)^2}{2!} + \frac{(-x)^3}{3!} + \dots\right) \right) / 2 \\
&\quad * \left(\left(1 + x + \frac{x^2}{2!} + \dots\right) - \left(1 - x + \frac{(-x)^2}{2!} + \frac{(-x)^3}{3!} + \dots\right) \right) / 2 \\
&\quad * \left(1 + x + \frac{x^2}{2!} + \dots\right) / 2 \\
&= \left(\frac{e^x - e^{-x}}{2}\right) \left(\frac{e^x + e^{-x}}{2}\right) (e^x) \\
&= \frac{e^{3x} - e^{-x}}{4} \\
&= \frac{1}{4} \left(\sum_{n=0}^{\infty} \frac{(3x)^n}{n!} - \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} \right) \\
&= \sum_{n=0}^{\infty} \frac{(3^n - (-1)^n) x^n}{4 n!}
\end{aligned}$$

Consider: What about multiplying other functions besides e^x

Recall that if $F(x) = \sum_{n=0}^{\infty} \bar{a}_n \frac{x^n}{n!}$, $G(x) = \sum_{n=0}^{\infty} \bar{b}_n \frac{x^n}{n!}$, the product had coefficients

$$\bar{c}_n = \sum_{i=0}^n \binom{n}{i} \bar{a}_i \bar{b}_{n-i}$$

This can be seen as separating $[n]$ into 2 blocks and assign group 1 to do something in \bar{a}_i ways and $n - i$ to do something in \bar{b}_{n-i} ways

Example: Find EGF and closed form a_n that counts total permutations of length n using A, B, C, D where the total number of A, B used is odd

EGF using odd numbers A, B in total is

$$G(x) = 2 \frac{x}{1!} + 2^3 \frac{x^3}{3!} + \dots = \sum_{k=0}^{\infty} \frac{(2x)^{2k+1}}{(2k+1)!} = \frac{e^{2x} - e^{-2x}}{2}$$

EGF for permutations of C, D (with no restriction)

$$H(x) = \left(1 + 1 \frac{x}{1} + 1 \frac{x^2}{2!}\right) \left(1 + 1 \frac{x}{1} + 1 \frac{x^2}{2!}\right) = e^x e^x$$

Thus closed form EGF for total permutations is $\frac{e^{2x} - e^{-2x}}{2} e^x e^x = \frac{e^{4x} - 1}{2}$

To find a closed for total ways a_n , we have

$$\begin{aligned}
\frac{1}{2}(e^{4x} - 1) &= \frac{1}{2} \sum_{n=1}^{\infty} \frac{(4x)^n}{n!} \\
&= \sum_{n=1}^{\infty} \frac{1}{2} 4^n \frac{x^n}{n!}
\end{aligned}$$

Thus $a_n = \frac{1}{2}4^n$

For a counting argument, for each good permutation, we can swap A, C to get a bad permutation or swap B, D to get a bad permutation. Thus exactly half of the permutations are good

Example: Place n distinct books into 3 bookshelves. What is the total number of ways to arrange the books such that each shelf has ≥ 1 book?

Counting argument: Look at all $n!$ permutations of books then split them into 3 bins $\implies n! \binom{n-1}{3-1}$

EGF argument: Let a_i be the total number of ways to place i books into shelf 1. Here $a_i = i!$

Thus for a single shelf, the EGF is $\sum_{i=1}^{\infty} a_i \frac{x^i}{i!} = \sum_{i=1}^{\infty} x^i$

To answer the original problem, we want coefficients of $\frac{x^n}{n!}$ in

$$\begin{aligned} \left(\sum_{i=1}^{\infty} x^i\right)^3 &= \left(\frac{1}{1-x} - 1\right)^3 \\ &= \left(\frac{x}{1-x}\right)^3 \\ &= x^3 \sum_{j=0}^{\infty} \underbrace{\binom{j+3-1}{3-1}}_{\text{weak comp into 3 parts OGF}} x^j \\ &= \sum_{j=0}^{\infty} \binom{j+3-1}{3-1} x^{j+3} \\ &= \sum_{n=3}^{\infty} \binom{n-1}{3-1} x^n \\ &= \sum_{n=3}^{\infty} n! \binom{n-1}{2} \frac{x^n}{n!} \end{aligned}$$

Thus the coefficients for the EGF is $n! \binom{n-1}{2}$

Example: EGF of D_n Derangements of $f: [n] \rightarrow [n]$

Let $D(x)$ be the EGF for D_n

$$D(x) = \sum_{n=0}^{\infty} D_n \frac{x^n}{n!}$$

Consider the EGF for all bijections where $f(i) = i$: $\sum_{n=0}^{\infty} 1 \frac{x^n}{n!} = e^x$ (only one way)

Now consider $D(x)e^x = (D_0 + D_1x + D_2\frac{x^2}{2!} + \dots)(1 + x + \frac{x^2}{2!} + \dots)$

- The coefficients of $\frac{x^n}{n!}$ is $\sum_{i=0}^n \binom{n}{i} D_i b_{n-i}$ (choose i to derange and $n-i$ to fix)
- This counts all possible bijections

Thus EGF of all bijections is

$$F(x) = \sum_{n=0}^{\infty} n! \frac{x^n}{n!} = \sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

Thus $\frac{1}{1-x} = D(x)e^x \implies D(x) = e^{-x} \frac{1}{1-x}$

To get the sequence a_n we have

$$\begin{aligned}
 D(x) &= e^{-x} \frac{1}{1-x} \\
 &= \left(\sum_{j=0}^{\infty} (-1)^j \frac{x^j}{j!} \right) \left(\sum_{i=0}^{\infty} x^i \right) \\
 &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \right) x^n \\
 &= \sum_{n=0}^{\infty} n! \sum_{k=0}^n \frac{(-1)^k}{k!} = D_n
 \end{aligned}$$

TODO LOOK AT THIS AGAIN

Example: How many ways can ≥ 1 people be split into any number of non-empty groups such that each group sits at an unlabelled circular table

EGF for 1 table with n people is

$$\begin{aligned}
 G(x) &= \sum_{n=1}^{\infty} a_n \frac{x^n}{n!} \\
 &= \sum_{n=1}^{\infty} (n-1)! \frac{x^n}{n!} \\
 &= \sum_{n=1}^{\infty} \frac{x^n}{n} \\
 &= \sum_{n=0}^{\infty} \frac{x^{n+1}}{n+1} \\
 &= \int \sum_{x=0}^{\infty} x^n \\
 &= \int \frac{1}{1-x} dx \\
 &= \ln\left(\frac{1}{1-x}\right)
 \end{aligned}$$

Thus EGF for k tables is

$$\frac{\ln\left(\frac{1}{1-x}\right) \ln\left(\frac{1}{1-x}\right) \cdots \ln\left(\frac{1}{1-x}\right)}{k!} \quad (\text{need to consider overcounting since tables are unlabelled})$$

For any number of tables, we need to sum over k and want the coefficient of $\frac{x^n}{n!}$

$$\sum_{k=0}^{\infty} \frac{\left(\ln\left(\frac{1}{1-x}\right)\right)^k}{k!} e^{\ln\left(\frac{1}{1-x}\right)} = \frac{1}{1-x}$$

Thus we see that

$$\sum_{n=0}^{\infty} x^n = \sum_{n=0}^{\infty} n! \frac{x^n}{n!}$$

Thus there are $n!$ ways to split n people into any number of tables

TODO LOOK AT THIS AGAIN

2.3.1 Compositions of EGF

We want to partition $[n]$ into nonempty blocks and do task 1 to each block, then task 2 to the set of blocks

If $F(x)$ and $G(x)$ are the EGFs for each respective task, provided $F(x) = \sum_{n=1}^{\infty} a_n \frac{x^n}{n!}$, the total number of ways to partition $[n]$ into any number of blocks with task 1 on each block and task 2 on the set of blocks is

$$H(x) = G(F(x)) = \sum_{n=0}^{\infty} h_n \frac{x^n}{n!}$$

Example What if $F(x), G(x)$ are trivial, i.e. total ways to partition $[n]$ such that each block is represented in 1 way and the set of blocks is seen in 1 way

$$F(x) = \sum_{n=1}^{\infty} \frac{x^n}{n!} = e^x - 1$$

$$G(x) = \sum_{k=0}^{\infty} \frac{x^k}{k!} = e^x$$

$$H(x) = e^{e^x - 1}$$

Example: Given n people, count the ways to seat them at any number of tables where each table is served red or white wine
To seat i people at a table, the EGF was

$$F(x) = \sum_{n=1}^{\infty} (i-1) \frac{x^i}{i!} = \ln\left(\frac{1}{1-x}\right)$$

Now given k tables, EGF for ways to serve red or white wine is

$$G(x) = \sum_{k=0}^{\infty} 2^k \frac{x^k}{k!} = e^{2x}$$

Thus we see that

$$H(x) = G(F(x)) = e^{2 \ln(\frac{1}{1-x})} = \frac{1}{(1-x)^2} = \frac{d}{dx} \left(\frac{1}{1-x} \right) = \frac{d}{dx} \left(\sum_{n=0}^{\infty} x^n \right) = \sum_{n=0}^{\infty} (n+1) n! \frac{x^n}{n!}$$

Thus there are $(n+1)!$ total ways

2.4 Catalan Numbers

Given a $m \times n$ lattice, how many UP, RIGHT paths from $(0,0)$ to (m,n) are there?

- This requires $m+n$ total steps $\implies \binom{n+m}{m} = \binom{n+m}{n}$

Now consider $n \times n$ case that don't cross the diagonal

Let C_n be the total number of paths that don't cross the diagonal

Partition these into classes where the last time the diagonal is touched is (i,i) for $0 \leq i \leq n-1$

Once we hit (i,i) , we have C_{n-i-1} paths left. Thus the total paths is

$$c_n = \sum_{i=0}^{n-1} C_i C_{n-i-1} \quad c_0 = 1$$

Theorem: If C_n denotes the total UP/RIGHT paths from $(0,0)$ to (n,n) that don't go over the diagonal is

$$C_n = \sum_{n=0}^{\infty} C_i C_{n-i-1}$$

Moreover, the OGF is

$$F(x) = \frac{1 - \sqrt{1 - 4x}}{2x} = \sum_{n=0}^{\infty} C_n x^n \quad C_n = \frac{\binom{2n}{n}}{n+1}$$

Proof:

$$\begin{aligned} F(x) &= 1 + \sum_{n=1}^{\infty} C_n x^n = 1 + \sum_{n=1}^{\infty} \left(\sum_{i=0}^{n-1} C_i C_{n-i-1} \right) x^n \\ &= 1 + x \sum_{n=1}^{\infty} \left(\sum_{i=0}^{n-1} C_i C_{n-i-1} \right) x^{n-1} \\ &= 1 + x(F(x))^2 \\ &= \frac{1 \pm \sqrt{1 - 4x}}{2x} \end{aligned}$$

For the algebra above, recall that for $f = \sum a_n x^n$ and $g = \sum b_n x^n$, we have

$$fg = \sum d_n x^n \quad d_n = \sum a_i b_{n-i}$$

Thus we see that

$$fg = \sum_{n=1}^{\infty} d_{n-1} x^{n-1} \implies \sum_{i=0}^{n-1} C_i C_{n-i-1} = F(x)^2$$

Additional algebra shows that

$$F(x) = \sum_{n=0}^{\infty} \frac{\binom{2n}{n}}{n+1} x^n$$

2.4.1 Combinatoric Proof of Catalan Numbers

We know that the total number of UP/RIGHT paths to (n, n) is $\binom{2n}{n}$. We now remove the bad paths

Observe that the first time we touch a bad diagonal (diagonal above main diagonal) is after i RIGHTS and $i+1$ UPS

- This results in $n-i$ RIGHTS and $n-(i+1)$ UPS to reach (n, n)

We alter the remaining path after touching the bad diagonal by reflecting it over the bad diagonal (so all RIGHTS are now UPS)

Now beyond the diagonal, we go $n-i-1$ RIGHTS and $n-i$ UPS

Thus total RIGHTS is $i + (n-i-1) = n-1$ and total UPS is $i+1 + (n-i) = n+1$

Consider which paths to $(n-1, n+1)$ will touch the bad diagonal? All of them

Thus any bad path to (n, n) that goes over the main diagonal corresponds to a path reaching $(n-1, n+1)$, establishing a bijection

$$\binom{(n-1) + n + 1}{n-1} = \binom{2n}{n-1}$$


Thus the total paths (n, n) that don't go over the main diagonal is

$$\binom{2n}{n} - \binom{2n}{n-1} = \frac{\binom{2n}{n}}{n+1}$$

Example: Total ways to correctly group n pairs of parentheses

Treat (as RIGHT and) as UP. Then we have the Catalan number C_n

Example: Total number of binary trees with n internal nodes and $n + 1$ leaves is



The root is ignored and we recurse over the subcases. Thus there are

$$C_i = \sum_{i=0}^{n-1} C_i C_{n-i-1}$$

total trees

Example: Total ways to triangulate a regular (labelled) polygon with $n + 2$ sides i.e. total ways to draw $n - 1$ lines connecting 2 vertices such that no lines cross

To triangulate a polygon with $n + 2$ sides, we create a triangle using the line joining vertex $n + 1$ and $n + 2$ as the base

Assume that these vertices join with a vertex k for $1 \leq k \leq n$. This creates 2 sub-polygons

- Polygon formed by $n + 2, 1, 2, \dots, k \implies k + 1$ vertices $\implies C_{k-1}$
- Polygon formed by $k, k + 1, \dots, n + 1 \implies n - k + 2$ vertices $\implies C_{n-k}$

Thus total number of triangulations is

$$\sum_{k=1}^n C_{k-1} C_{n-k} = \sum_{n=0}^n C_k C_{n-k-1} = C_n$$

3 Graph Theory

3.1 Introduction to Graph Theory

Definition - (Simple) Graph G: ordered pair (V, E) where $V = V(G)$ is a set of **vertices** and $E = E(G)$ is a set of **edges** (2-size subsets of V)

Definition - Adjacent: Two vertices $v_1, v_2 \in V$ are **adjacent** if $\{v_1, v_2\} \in E$, denoted as $v_1 \sim v_2$

Definition - Incident: An edge $e = \{v_1, v_2\}$ is **incident** to a vertex v if v is one of the endpoints of e

Definition - Subgraph: Graph H with $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$

- **Spanning Subgraph** if $V(H) = V(G)$
- **Vertex Induced Subgraph** if whenever $u, v \in V(H)$ and $u \sim v \in E(G)$, then $u \sim v \in E(H)$
- **Subgraph Induced by S** if $S = V(H) \subseteq V(G)$, denoted $G[S]$

Definition - Walk: Sequence of vertices beginning at u and ending at v such that consecutive vertices are adjacent

- **Closed** if $u = v$
- **Length:** total edges traversed

Definition - Trail: Walk with no edge traversed more than once

Definition - Path: Walk with no vertex traversed more than once

- **Note:** path \implies trail \implies walk

Definition - Circuit: Closed trail of length 3 or more

- **Cycle:** Circuit with only the first and last vertex repeated
 - **Even Cycle** if total edges traversed is even
 - **Odd Cycle** if total edges traversed is odd

Definition - Connected Graph: Graph G such that for any distinct vertices u, v there is a path uv

Definition - Distance: Minimum length of a path uv , denoted $d(u, v)$

Definition - Diameter: Distance of the farthest apart vertices, denoted $\text{diam}(G) = \max_{u, v \in V(G)} \{d(u, v)\}$

Definition - Path Graph: Graph on n vertices, denoted P_n , with edge set $E(P_n) = \{\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_n\}\}$

- **Note:** P_n will have $n - 1$ edges

Definition - Cycle Graph: Graph on n vertices, denoted C_n , with edge set $E(C_n) = \{\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_1\}, \{v_n, v_1\}\}$

- **Note:** C_n will have n edges


Definition - Complete Graph: Graph on n vertices, denoted K_n with edge set $E(K_n) = \{\{v_i, v_j\} \mid 1 \leq i \neq j \leq n\}$

- **Note:** $E(K_n)$ has size $\binom{n}{2}$

Definition - Complement: The **complement** of graph G , denoted \overline{G} , has vertex set $V(\overline{G}) = V(G)$ and edge set $E(\overline{G}) = \{v_i, v_j \mid v_i, v_j \notin E(G)\}$

Definition - Bipartite: G is **bipartite** if its vertex set can be partitioned into 2 subsets A, B , called **partite sets**: such that every edge of G has one endpoint in A and the other in B

Example



Above graph cannot be bipartite since the neighbors of ?? lie in A and B

Theorem: G is bipartite if and only if G contains no odd cycles

Proof: \Rightarrow By contraposition, assume G has an odd cycle $v_1, v_2, \dots, v_{2k-1}, v_1$

Then we have $v_1 \in A, v_2 \in B, \dots, v_{2k-1} \in A, v_1 \in A$. Thus G is not bipartite

\Leftarrow WLOG, let G be connected (if it isn't take any connected component in G)


Let $u \in V(G)$ and assume G has no odd cycles (all cycles are even)

Define $A = \{w \mid d(u, w) \text{ is even}\}$ and $B = \{w \mid d(u, w) \text{ is odd}\}$ (these are our partites)

BWOC, let $w_1, w_2 \in B$ such that $w_1 \sim w_2$ (similar idea can be shown for A)

Consider the paths $u = v_0, v_1, \dots, v_{2s+1} = w_1$ and $u = u_0, u_1, \dots, u_{2t+1} = w_2$

Let x be the last vertex shared by the 2 paths



Necessarily, we must have that $x = v_i = u_i$. Otherwise a shorter path exists

But then the cycle containing $x, \dots, w_1, w_2, \dots, x$ is of odd length (since the paths xw_1 and xw_2 have the same parity)

Thus it is impossible for an edge to connect two nodes in B

Similarly, an edge cannot connect two nodes in A

Thus any edge must connect between a node in A and a node in B , which are our partites, and thus G is bipartite

Definition - Complete Bipartite Graph with partites A, B of size a, b is denoted $K_{a,b}$ where every vertex in A is adjacent to every vertex in B

3.2 Vertex Degree

Definition - Vertex Degree: Total edges incident to v , denoted $\deg(v)$

Definition - Neighborhood: Set of vertices adjacent to v , denoted $N(v)$

Definition - Min [Max] Degree: Lowestest [highest] degree of G , denoted $\delta(G)$ [$\Delta(G)$]

- **Note:** An isolated vertex has degree 0

Theorem: A Graph G with size m (m edges) has $\sum_{v \in V(G)} \deg(v) = 2m$

Proof: Summing over all vertices, every edge gets counted twice by the 2 endpoints

Corollary: Graph G must have an even number of vertices with odd degree

Definition - d-regular: Graph G where every vertex has degree d

Example: K_4 is 3-regular with 4 vertices




Non-Example: A 3-regular graph with 5 vertices doesn't exist by the Corollary above (there must be an even number of vertices with odd degree)


Theorem: There exists a d -regular graph on n vertices if and only if at least one of d, n is even

Proof: \implies By Contraposition. If d, n are both odd, then a d -regular graph doesn't exist by the Corollary above

\Leftarrow If $d = 2k$, arrange the vertices in a cycle. Then v_i joins to k preceding and k succeeding vertices

Otherwise assume d is odd and n is even. We join edges as before, along with v_i joining to $v_{i+t \pmod n}$ where $n = 2t$ (the opposite vertex)





Theorem: For a given graph G , there exists a d -regular graph H such that G is an induced subgraph of H

Proof: If G is d -regular, then $H = G$ and we are done

Otherwise, create a copy of G , call it G'

Now construct G_1 by adding $v_i \in G \sim v'_i \in G'$ for vertices of G with $\deg(v) < d$

Thus $\delta(G_1) = \delta(G) + 1$. If G_1 is d -regular, then we are done

Otherwise, apply the procedure recursively on G_1

Example:




Figure 1: Non-regular Graph Induced in a Regular Graph

Definition - Degree Sequence: Given a graph G with n vertices, its **degree sequence** is a non-increasing sequence of length n whose i th term is the degree of vertex i

Theorem: Let $d = d_1, \dots, d_n$ be a non-increasing sequence. There exists a graphical graph G with these degrees if and only if the sequence

$$s_1 = d_2 - 1, d_3 - 1, \dots, d_{d_1+1} - 1, \underbrace{d_{d_1+2}, \dots, d_n}_{\text{no minus term}}$$

is graphical. That is, delete d_1 then delete 1 edge from the next d_1 terms (may require reordering the sequence)

Proof: \Leftarrow Given $d_2 - 1, \dots, d_n$ is graphical, we can add a new vertex v_1 and join v_1 to the first d_1 vertices in the sequence \Rightarrow Suppose that there is a graph H with the degree sequence s and a vertex u of degree d_1 that is adjacent with vertices of degree $d_2, d_3, \dots, d_{d_1+1}$. Then clearly s_1 is the sequence of $H \setminus u$ and is graphical

Now we show that such a graph H has to exist

BWOC, suppose no graph H exists such that v_1 is adjacent to vertices of degree $d_2, d_3, \dots, d_{d_1+1}$

Among all graphs with degree sequence s , let G be the graph with the largest sum of degree of neighbors of v_1

Since v_1 is not adjacent to vertices with degree d_2, \dots, d_{d_1+1} , v_1 is NOT adjacent to the next d_1 vertices with the highest degrees

Thus $v_1 \sim v_s$, where $\deg(v_s) < \deg(v_r)$ for some $v_r \in V(G)$ that is NOT adjacent to v_1

Since $\deg(v_r) > \deg(v_s)$ there is some $v_t \in V(G)$ such that $v_t \sim v_r$ but NOT adjacent to v_s

Now consider the graph G' where we remove $v_1 \sim v_s$ and $v_r \sim v_t$ and replace with $v_1 \sim v_r$ and $v_s \sim v_t$




Figure 2: Degree Sequence Proof Diagram

Clearly the G, G' have the same vertex set and satisfy the degree sequence s

However, now the sum of the degrees of vertices adjacent to v_1 is larger than our proposed maximal. Contradiction

Thus such a graph H must exist and the sequence s_1 is graphical

Example: $s = (5, 4, 3, 3, 2, 2, 2, 1, 1, 1)$

$s_1 = (3, 2, 2, 1, 1, 2, 1, 1, 1) \leftrightarrow (3, 2, 2, 2, 1, 1, 1, 1, 1)$

$s_2 = (1, 1, 1, 1, 1, 1, 1, 1, 1)$

- **Note:** When going from s_1 to s , we need to add edges to the ORIGINAL sequence




Figure 3: Degree Sequence Example 1

Non-Example: $s = 7, 7, 4, 3, 3, 3, 2, 1$

$s_1 = (6, 3, 2, 2, 2, 1, 0)$

$s_2 = (2, 1, 1, 1, 0, -1)$ impossible to have a vertex of negative degree

Thus s is NOT graphical

- **Note:** As another argument, G has order 8 and s has 2 vertices with degree 7, thus each vertex must have at least degree 2. Thus s is NOT graphical

3.3 Properties of Trees

Definition - Acyclic: A graph is **acyclic** if it contains no cycles

Definition - Tree: A **tree** is an acyclic, connected graph

Definition - Leaf: A **leaf** is a vertex with degree 1

Definition - Forest: A **forest** is an acyclic graph (union of trees)

Notation $G \setminus v$: Denotes the graph created by deleting vertex v and all adjacent edges to it

Example: Consider K_4 and removing the bottom-right corner vertex



Theorem: Every tree on 2 or more vertices has at least 2 leaves

Proof: Since a tree is acyclic, take a path of longest length with endpoints u, v

Neither u nor v can have $\deg \geq 2$ since otherwise a longer path would exist

Thus $\deg(u) = \deg(v) = 1$

Theorem: Let G be on ≥ 1 vertices. The following are equivalent

1. G is a tree (connected and acyclic)
2. G is connected with $n - 1$ edges
3. G is acyclic with $n - 1$ edges
4. There exists a unique path for any 2 distinct vertices $u, v \in V(G)$

Proof $1 \rightarrow 2, 3$: By Induction on n

Base case: $n = 1$. Here G is an isolated vertex with $1 - 1 = 0$ edges

IH: Suppose that for a G on $i \leq n$ vertices, G is connected and acyclic with $i - 1$ edges

IS: Let G be on $n + 1$ vertices. Necessarily, G must have a leaf v , so let $G' = G \setminus v$

Here G' is on n vertices, and by IH must be connected with $n - 1$ edges

Adding back v and its incident edge, we get that G has $n + 1 - 1 = n$ edges

Furthermore, since a leaf has degree 1, G must be connected and acyclic

Proof $2 \rightarrow 3$: Let G be a connected graph with $n - 1$ edges.

BWOC, assume that G has cycles

We can delete edges so that there are no cycles but G remains connected

This is possible since this only removes 1 of 2 paths between vertices

Thus this new graph call it G' , is acyclic and connected \implies it's a tree

However, G' has $< n - 1$ edges which contradicts the previous proof.

Thus G must be acyclic with $n - 1$ edges

Proof $3 \rightarrow 1$: Let G be an acyclic graph with $n - 1$ edges. We show that G is connected

Let C_1, \dots, C_k be the components of G and assume that C_i has n_i vertices

Then by the previous proof, each C_i is connected and acyclic and is thus a tree

From the first proof, each C_i has $n_i - 1$ edges. By assumption, we also have that G has $n - 1$ total edges. Thus we have

$$n - 1 = \sum_{i=1}^k n_i - 1 = n - k \implies k = 1$$

Thus G is connected

Proof $1 \rightarrow 4$: Let G be a tree. BWOC, suppose that there exists 2 uv -paths

Let w be the first vertex shared such that the next one is not shared

Let z be the vertex after w that is shared by both paths

This creates a cycle, contradicting that G is a tree

Thus there exists a unique path for any 2 distinct vertices u, v

Proof $4 \rightarrow 1$: Consider a unique uv -path between any 2 arbitrary vertices u, v

By assumption, G is connected since any path can be formed

BWOC, suppose that G has a cycle. Then there are some vertices that have 2 possible paths between each other

Contradiction. Thus G must be acyclic and thus a tree

Theorem: The minimum number of edges in a connected graph on n vertices is $n - 1$

Proof: The statement clearly holds for $n = 1, 2, 3$. Thus we look at $n \geq 4$

BWOC, suppose that G be a graph of smallest order that is connected on $m \leq n - 2$ edges

First we show that G has a leaf. BWOC, suppose G has no leaves.

Since $\sum_{v \in V(G)} \deg(v) = 2m$, if there were no leaves, then $2m \geq \underbrace{2n}_{\deg(v) \geq 2} \implies m \geq n$, contradicting that $m \leq n - 2$

Thus G has a leaf, call it v

Looking at $G' = G \setminus v$, we see that G' is connected on $n - 1$ vertices with $n - 3$ or fewer edges

This contradicts the minimality of n since we claimed that G is the graph of smallest order that is connected on $m \leq n - 2$ edges

Thus the minimum number of edges in a connected graph on n vertices is $n - 1$

Example: Suppose that G is a tree on 13 vertices with degrees 1, 2, 5. If G has 3 vertices of degree 2, how many leaves does it have?

Let x be the total number of leaves. Then we have $13 - 3 - x = 10 - x$ vertices of degree 5

Moreover, we see that

$$\sum_{v \in V} \deg(v) = (3)2 + (x)1 + (10 - x)5 = 2(13 - 1) = 24 \implies x = 8$$

Definition - Bridge: An edge e is a **bridge** of a connected graph if removing it ($G \setminus e$) disconnects the graph

Theorem: An edge e is a bridge if and only if e isn't in any cycles

3.4 Spanning Trees

Definition - Spanning Tree: Tree T such that $V(T) = V(G)$ and $E(T) \subseteq E(G)$

Consider the total number of trees on n vertices


- If the vertices are unlabelled, there is no easy formula
- If the vertices are labelled, we can look at the problem: How many labelled spanning trees are in K_n

Definition - Prüfer Code: A sequence of length $n - 2$ where each entry can take on 1 to n , which corresponds to a labelled tree

Converting a Spanning Tree to a Prüfer Code:

1. Delete the leaf with the lowest index and write down the vertex adjacent to the leaf
2. Repeat until only an edge remains

Example: Consider the tree



The Prüfer code is 2, 3, 2, 6. This comes from

- Removing 1 \implies write down 2
- Removing 4 \implies write down 3
- Removing 3 \implies write down 2
- Removing 2 \implies write down 6
- Stop on the edge $6 \sim 5$

Prüfer Code to Spanning Tree:


1. Given $(a_1, a_2, \dots, a_{n-2})$, let b_1 be the smallest index not used
2. Create $a_1 \sim b_1$
3. Delete a_1 and append b_1 to the end of the sequence
4. Let b_2 be the lowest index showing up in the new sequence

5. Create the edge $b_2 \sim a_2$
6. Delete a_2 and append b_2 to the end of the sequence
7. Repeat until the sequence is b_1, \dots, b_{n-2}
8. There will be 2 indices i, j NOT in the sequence. Create an edge $i \sim j$ to form a tree

Example: Consider the sequence (2, 2, 6, 4)

- $b_1 = 1 \implies (2, 6, 4, 1)$
- $b_2 = 3 \implies (6, 4, 1, 3)$
- $b_3 = 2 \implies (4, 1, 3, 2)$
- $b_4 = 5 \implies (1, 3, 2, 5)$
- Stop. 4, 6 aren't in the graph, so connect them

The corresponding tree is



Where the order of edge insertion is red, orange, green, blue, brown

Theorem: Each Prufer code corresponds to a unique tree. Hence the total number of labelled spanning trees of K_n is n^{n-2}

- This comes from having ending the algorithm with a sequence b_1, \dots, b_{n-2} where each b_i can take on any n value

Proof: Given a tree, using our algorithm, we have a sequence

Now we show that every sequence corresponds to a unique by induction on the number of vertices n

Base Case: Easily verifiable for $n = 3$

IH: Assume that the claim is true up to some n

IS: Given (a_1, \dots, a_{n-1}) (so the tree has $n + 1$ vertices), show that the sequence comes from exactly 1 tree

Let x be the smallest index not in the sequence. Then in our construction of the tree, x will be a leaf and $x \sim a_1$


Now consider the subtree excluding x and the edge incident on it. By IH, this tree must be unique

But then the sequence (a_1, \dots, a_{n-1}) must correspond to a unique tree with $x \sim a_1$. Otherwise a_1 would not be the first value of the sequence

Consider: Given a vertex i with degree d_i , how many times will the value show up in the code?

Every time we delete a leaf adjacent to i , it will appear in the sequence until i is a leaf

Thus i shows up $d_i - 1$ times




Corollary: The total trees (labelled) such that vertex i has degree d_i is

$$\binom{n-2}{d_1-1, d_2-1, \dots} \quad \text{Permutation with repetition}$$


Example: Let $n = 6$ and consider $d_1 = d_2 = 3$ and $d_3 = d_4 = d_5 = d_6 = 1$

By the formula, we have $\binom{6-2}{2,2} = 6$ trees



Note that the tree is determined by picking 2 of the leaf indices $\implies \binom{4}{2} = 6$ ways to create labelled trees with thus Prufer code

This follows from swapping labels on either ends of the tree correspond to the same tree. That is




Correspond to the same tree (by rotation argument)

3.4.1 Rooted Plane Trees

Definition - Rooted Plane Tree: Trees that have a starting root vertex and the children of the tree have a left, right ordering

- **Note:** Vertices are NOT labelled

Example: Consider $n = 4$. There are 5 rooted plane trees



- **Note:** The 3rd and 4th trees differ because of left, right ordering

On rooted plane trees, we can do a clockwise walk on the border of the tree starting at the root

In this traversal, each edge is covered twice

In particular, as we move down, we can never go up more than we have gone down (similar to Catalan Numbers)

Thus the number of rooted plane trees on $n + 1$ vertices (because of the root) equals the Catalan Numbers


$$C_n = \frac{\binom{2n}{n}}{n+1}$$

3.4.2 Rooted Forest


Definition - Rooted Forest: A forest where each tree (component) has a distinguishable vertex as the root

Consider: What is the total number of labelled rooted forests?

Example: $n = 3$



In particular, given a rooted forest, we can add a vertex $(n + 1)$ that joins the roots to create a labelled tree on $n + 1$ vertices



Here the red vertices were the roots of the rooted forest and the blue vertex is the added vertex $n + 1$

Conversely, we can also reverse the process (take a labelled tree and convert it to a rooted forest) by deleting vertex $n + 1$

The vertices adjacent to the vertex $n + 1$ will now become the roots of the rooted forest

This creates a bijection between rooted labelled forests and labelled trees on $n + 1$ vertices

Theorem: The number of labelled rooted forests on n vertices is

$$(n + 1)^{n-1}$$

Proof: The formula comes from labelled trees on $n + 1$ vertices

3.4.3 Car Parking Problem

Suppose n cars go down a 1-way street. There are n parking spots and each car has a favorite spot

Each car drives to its favorite spot

- If it available, the car will park at the spot
- Otherwise the car will drive to the next open spot

Example: The following set up works

Car	Favorite
1	1
2	1
3	2
4	2

Non-Example: The following set up does NOT work since car 4 can't park

Car	Favorite
1	2
2	2
3	2
4	2

Consider: How many preferences are there such that all cars can park?