

EE4211: Data Science for the Internet of Things

Logistic Regression

Biplab Sikdar

Agenda

2

- Examples
- Logit function
- Fitting a logistic regression
- Understanding the parameters

Example

3

- **Regression analysis:** describes the relationship between two (or more) variables
- Examples:
 - Income and educational level
 - Demand for electricity and the weather
 - Tire tread depth and mileage
- Linear regression:
 - The data is represented as $(\epsilon_i \sim N(0, \sigma^2))$:
$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i$$
 - The fitted model:

$$\hat{y} = \beta_0 + \beta_1 x$$

Example

4

- Typical retail/restaurant chain: hundreds of pieces of equipment in each store, spread across thousands of store locations across geographic locations.
- Preventive maintenance in such scenarios: difficult problem.
- Sensors mounted on IoT devices, equipment, and process monitoring controllers continuously collect and transmit data.
- Rule based systems raise false alarms that turn out to be less serious alerts upon actual inspection, such as an open door on a refrigeration unit.
- Use analytics to learn, enhance, and implement a new method for predicting equipment failure using IoT sensor data.

Example

5

iot_sensor_dataset (version 1) [Autosaved] - Excel

File Home Insert Page Layout Formulas Data Review View Acrobat Tell me what you want to do... Biplab Sikdar Share

PivotTable Recommended PivotTables Tables

Pictures Online Pictures Illustrations

Shapes SmartArt Screenshot

Store My Add-ins Add-ins

Recommended Charts Charts

PivotChart

3D Map Tours

Line Column Win/Loss Sparklines

Slicer Timeline Filters

Hyperlink Links

Text Box Header & Footer Text

Equation Symbol

Symbols

W23


Name Box

| | A | B | C | D | E | F | G | H | I |
|----|----------|-------|-----|-------------|------------|------|------|---|---|
| 1 | football | atemp | PID | outpressure | inpressure | temp | fail | | |
| 2 | 0 | 7 | 6 | 36 | 3 | 1 | 1 | | |
| 3 | 190 | 1 | 1 | 20 | 4 | 1 | 0 | | |
| 4 | 31 | 7 | 1 | 24 | 6 | 1 | 0 | | |
| 5 | 83 | 4 | 1 | 28 | 6 | 1 | 0 | | |
| 6 | 640 | 7 | 0 | 68 | 6 | 1 | 0 | | |
| 7 | 110 | 3 | 1 | 21 | 4 | 1 | 0 | | |
| 8 | 100 | 7 | 1 | 77 | 4 | 1 | 0 | | |
| 9 | 31 | 1 | 4 | 21 | 4 | 1 | 0 | | |
| 10 | 180 | 7 | 3 | 31 | 4 | 1 | 0 | | |
| 11 | 2800 | 0 | 0 | 39 | 3 | 1 | 0 | | |

iot_sensor_dataset


Example

6




Example

7



Example


8




Best Fit Curve

9

- Data that has a sharp cut off point between categories:



- Data with a lengthy transition between categories:



Logistic Regression

10

- Response/outcome/dependent variable: y
- Predictor/explanatory/independent variable: x
- Logistic regression extends the ideas of linear regression to the situation where the dependent variable, y , is categorical.
- **Categorical variable:** divides the observations into classes.
 - y : state of a machine (normal, alarm, failure)
 - Each of the measurements in the dataset (the observations) may be classified as belonging to one of three classes.

Logistic Regression

11

- Uses a logarithmic transformation to allow a non-linear relationship to be expressed in a linear way.
- Logistic regression can be used for classifying a new observation into one of the classes, based on the values of its predictor variables (called “**classification**”).
- It can also be used in data (where the class is known) to find similarities between observations within each class in terms of the predictor variables (called “**profiling**”).

Logistic Regression Model

12

- Example:
 - We are interested evaluating the success (y) of a new fire-retardant spray is at preventing “fire” and how it depends on the amount (x) that is applied.
 - The values of y are 1 (Success) or 0 (Failure).
 - The values of x range over a continuum

Logistic Regression Model

13



Logistic Regression Model

14

- $p \triangleq P[y = 1] = P[\text{success}]$
- p in general will increase with x
- **Odds ratio:** $\frac{p}{1-p}$
- The odds ratio also increases with the value of x , ranging from zero to infinity
- **Log odds ratio (logit):** $\ln\left(\frac{p}{1-p}\right)$

Example: Odds Ratio

15

- We roll a regular, six-faced die
- Success: “roll a six”
- $P[\text{success}] = p = \frac{1}{6}$
- **Odds ratio:** $\frac{p}{1-p} = \frac{\frac{1}{6}}{1-\frac{1}{6}} = \frac{1}{5}$
- **Log odds ratio (logit):** $\ln\left(\frac{p}{1-p}\right) = \ln\left(\frac{1}{5}\right) = -1.69$

Better Example: Odds Ratio

16

| | | |
|---|------------------------|-------------|
|  | MANCHESTER CITY | 4/6 |
|  | LIVERPOOL | 11/4 |
|  | CHELSEA | 10/1 |
|  | TOTTENHAM | 16/1 |
|  | MANCHESTER UTD | 25/1 |

Logistic Regression Model

17

- The logistic regression model assumes that the log odds ratio is linearly related to x :

$$z \rightarrow \ln \left(\frac{p}{1-p} \right) = \beta_0 + \beta_1 x$$

- This can be solved using the technique used solving an ordinary (linear) regression!
- Fit observed data to obtain β_0 and β_1

Logistic Regression Model

18

- In terms of the odd ratio:

$$\frac{p}{1-p} = e^{\beta_0 + \beta_1 x}$$

- Solving for p in terms of x :

$$\frac{p}{1-p} = e^{\beta_0 + \beta_1 x}$$

$$\Rightarrow p = (1-p)e^{\beta_0 + \beta_1 x}$$

$$\Rightarrow p + pe^{\beta_0 + \beta_1 x} = e^{\beta_0 + \beta_1 x}$$

$$\Rightarrow p = \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}}$$


$$\Rightarrow p = \frac{1}{1 + e^{-(\beta_0 + \beta_1 x)}}$$

Sigmoid
function

Interpretation: β_0

19


- β_0 : determines the intercept



Interpretation: β_1

20

- β_1 : determines when $p = 0.5$ (along with β_0)



Interpretation: β_1

21

□ Also:


$$\begin{aligned}\frac{dp}{dx} &= \frac{d}{dx} \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}} \\ &= \frac{e^{\beta_0 + \beta_1 x} \beta_1 (1 + e^{\beta_0 + \beta_1 x}) - e^{\beta_0 + \beta_1 x} \beta_1 e^{\beta_0 + \beta_1 x}}{(1 + e^{\beta_0 + \beta_1 x})^2} \\ &= \frac{e^{\beta_0 + \beta_1 x} \beta_1}{(1 + e^{\beta_0 + \beta_1 x})^2} \\ &= \frac{\beta_1}{4} \quad \text{when } x = -\frac{\beta_0}{\beta_1}\end{aligned}$$

□ $\frac{\beta_1}{4}$: rate of increase in p with respect to x when $p = 0.5$

Interpretation: β_1

22

- β_1 : determines slope when $p = 0.5$




Example 1

23

- In a study of the effectiveness of coupons offering a price reduction, 1,000 homes were selected and coupons mailed
- Coupon price reductions: 5, 10, 15, 20, and 30 dollars
- 200 homes assigned at random to each coupon value
- x : amount of price reduction
- y : binary variable indicating whether or not coupon was redeemed


Example 1

24



Example 1

25



□ Fitted response function: $\beta_0 = -2.04$ and $\beta_1 = 0.097$

Example 2

26

- For each value of x , we may not have probability but rather a number of $\{x, y\}$ pairs
- **Step 1:** Extract frequencies from the pairwise data and hence probabilities
 - Raw data: $\{12, 0\}, \{12, 1\}, \{14, 0\}, \{12, 1\}, \{14, 1\}, \{14, 1\}, \{12, 0\}, \{12, 0\}$
 - Probability data: ($p = 1$ for each x , third entry is number of occurrences in raw data): $\{12, 0.4, 5\}, \{14, 0.67, 2\}$
- **Step 2:** Obtain the odds ratio data
- **Step 3:** Obtain β_0 and β_1 using linear regression model

Example 2

27

| Number of Trips | Require Maintenance? | | Total |
|-----------------|----------------------|-----|-------|
| | No | Yes | |
| 1 | 9 | 1 | 10 |
| 2 | 13 | 2 | 15 |
| 3 | 9 | 3 | 12 |
| 4 | 10 | 5 | 15 |
| 5 | 7 | 6 | 13 |
| 6 | 3 | 5 | 8 |
| 7 | 4 | 13 | 17 |
| 8 | 2 | 8 | 10 |
| Total | 57 | 43 | 100 |

Example 2

28

| Number of Trips | P(maintenance) | odds | Log odds | Number of occurrences |
|-----------------|----------------|--------|----------|-----------------------|
| 1 | 0.1000 | 0.1111 | -2.1972 | 10 |
| 2 | 0.1333 | 0.1538 | -1.8718 | 15 |
| 3 | 0.2500 | 0.3333 | -1.0986 | 12 |
| 4 | 0.3333 | 0.5000 | -0.6931 | 15 |
| 5 | 0.4615 | 0.8571 | -0.1542 | 13 |
| 6 | 0.6250 | 1.6667 | 0.5108 | 8 |
| 7 | 0.7647 | 3.2500 | 1.1787 | 17 |
| 8 | 0.8000 | 4.000 | 1.3863 | 10 |

Example 2

29

| x (trips) | y (log odds) | x^2 | xy | Number of occurrences |
|-------------|-----------------|-------------|-----------------|-----------------------|
| 1 | -2.1972 | 1 | -2.1972 | 10 |
| 2 | -1.8718 | 4 | -3.7436 | 15 |
| 3 | -1.0986 | 9 | -3.2958 | 12 |
| 4 | -0.6931 | 16 | -2.7726 | 15 |
| 5 | -0.1542 | 25 | -0.7708 | 13 |
| 6 | 0.5108 | 36 | 3.0650 | 8 |
| 7 | 1.1787 | 49 | 8.2506 | 17 |
| 8 | 1.3863 | 64 | 11.0904 | 10 |
| 448 | -37.6471 | 2504 | 106.3981 | 100 |

Bottom row: weighted sum ($\sum_{i=1}^8 value_i \times occurrences_i$)

Example 2

30

□ Results from regression

□ $\beta_0 = -2.856$ and $\beta_1 = 0.5535$

| Number of Trips | P(maintenance) | \hat{p} |
|-----------------|----------------|-----------|
| 1 | 0.1000 | 0.0909 |
| 2 | 0.1333 | 0.1482 |
| 3 | 0.2500 | 0.2323 |
| 4 | 0.3333 | 0.3448 |
| 5 | 0.4615 | 0.4788 |
| 6 | 0.6250 | 0.6142 |
| 7 | 0.7647 | 0.7346 |
| 8 | 0.8000 | 0.8280 |


SSE 0.0028

SST 0.5265

R^2 0.9946

Logistic vs Linear

31



Poisson Regression

32

- Y is a non-negative integer: $0, 1, 2, \dots$ (usually small values)

- Model:

$$\ln(E[Y]) = \beta_0 + \beta_1 x$$

- This implies:

$$E[Y] = e^{\beta_0 + \beta_1 x}$$

Acknowledgements

33

- A number of the slides in this lecture are based on material from various sources:
 - Luke Sloan
 - Luiz Pessoa