


EE4704 Image Processing and Analysis

Tutorial Set E – Solution

Question 1

Part (a)



$$(x_i, y_i) \rightarrow b = -x_i a + y_i$$

$$P1: (-4, 0) \rightarrow b = 4a$$


$$P2: (-2, 1) \rightarrow b = 2a + 1$$

$$P3: (0, 1) \rightarrow b = 1$$

$$P4: (2, 1.5) \rightarrow b = -2a + 1.5$$

$$P5: (4, 2) \rightarrow b = -4a + 2$$

$$P6: (6, 2) \rightarrow b = -6a + 2$$



Part (b)

$$(x_i, y_i) \rightarrow \rho = \sqrt{x_i^2 + y_i^2} \cos(\theta - \tan^{-1}(y_i/x_i))$$

$$P1: (-4, 0) \rightarrow \rho = 4 \cos(\theta - \tan^{-1}(\frac{0}{-4})) = 4 \cos(\theta - 3.14)$$


$$P2: (-2, 1) \rightarrow \rho = \sqrt{5} \cos(\theta - \tan^{-1}(\frac{1}{-2})) = \sqrt{5} \cos(\theta - 2.68)$$

$$P3: (0, 1) \rightarrow \rho = 1 \cos(\theta - \tan^{-1}(\frac{1}{0})) = 1 \cos(\theta - 1.57)$$

$$P4: (2, 1.5) \rightarrow \rho = \sqrt{6.25} \cos(\theta - \tan^{-1}(\frac{1.5}{2})) = \sqrt{6.25} \cos(\theta - 0.64)$$

$$P5: (4, 2) \rightarrow \rho = \sqrt{20} \cos(\theta - \tan^{-1}(\frac{2}{4})) = \sqrt{20} \cos(\theta - 0.46)$$

$$P6: (6, 2) \rightarrow \rho = \sqrt{40} \cos(\theta - \tan^{-1}(\frac{2}{6})) = \sqrt{40} \cos(\theta - 0.32)$$



Detected line parameters:

$$\rho = -1, \quad \theta = -1.4 \text{ rad}$$

Line equation:

$$x \cos(-1.4) + y \sin(-1.4) = -1$$

$$y = 0.17x + 1.0$$

Question 2

$$(x_i, y_i) \rightarrow b = -x_i a + y_i$$

The accumulator array has limits

$$-1 \leq a \leq +1, \quad -2 \leq b \leq +14$$

with $\Delta a = 0.2$ and $\Delta b = 1$.

The center coordinates of the cells are

$$\begin{aligned} a_i &= -1, -0.8, -0.6, \dots, 0, \dots, 0.8, 1 & (\Delta a = 0.2) \\ b_j &= -2, -1, 0, \dots, 13, 14 & (\Delta b = 1) \end{aligned}$$



The cell values can be obtained using a table:

$a_i :$		-1	-0.8	-0.6	-0.4	-0.2	0	0.2	0.4	0.6	0.8	1
(4, 9)	$b = -4a + 9$	13.0	12.2	11.4	10.6	9.8	9.0	8.2	7.4	6.6	5.8	5.0
(5, 3)	$b = -5a + 3$	8	7	6	5	4	3	2	1	0	-1	-2
(6, 9)	$b = -6a + 9$...										
(8, 5)	$b = -8a + 5$											
(8, 8)	$b = -8a + 8$											
(10, 10)	$b = -10a + 10$											
(12, 6)	$b = -12a + 6$											
(14, 10)	$b = -14a + 10$											
(16, 5)	$b = -16a + 5$											
(16, 12)	$b = -16a + 12$											

From the completed accumulator array, the largest cell values are 5 obtained at cell $(-0.4, 11)$, and 4 obtained at $(0.8, -1)$. Hence, the corresponding line equations are

$$L1 : y = -0.4x + 11$$

$$L2 : y = 0.8x - 1$$



Question 3

Gray level, z_k :	0	1	2	3	4	5	6	7	8	9
No. of pixels, n_k :	400	800	800	1200	400	800	2000	1600	1200	800

Initial estimate

$$T_0 = \text{mean} = \frac{1}{\sum_k n_k} \sum_k n_k z_k = 5.16$$

The histogram is partitioned into two groups:

$$0 \rightarrow 5 \quad \mu_1 = 2.63 \quad ; \quad 6 \rightarrow 9 \quad \mu_2 = 7.14$$

$$\begin{aligned} T_1 &= \frac{1}{2}(2.63 + 7.14) \\ &= 4.89 \quad (\text{Threshold} = 4) \end{aligned}$$

Next, we have


$$\begin{aligned} 0 \rightarrow 4 \quad \mu_1 = 2.11 \quad ; \quad 5 \rightarrow 9 \quad \mu_2 = 6.88 \\ T_2 &= \frac{1}{2}(2.11 + 6.88) \\ &= 4.49 \quad (\text{Threshold} = 4) \end{aligned}$$

The threshold value is unchanged; hence the selected threshold is

$$T = 4$$



i.e.

$$g(x, y) = \begin{cases} 1 & z > 4 \\ 0 & z \leq 4 \end{cases}$$



Question 4

Part (a)



Starting from the leftmost pixel of the top row, we have

00000654664443212

This code is normalised for starting point.

Part (b)


Denote the 17 boundary pixels by $b_i(x_i, y_i)$, $i = 1, 2, \dots, 17$. The centroid is estimated by

$$\begin{aligned}\bar{x} &= \frac{1}{17} \sum_{i=1}^{17} x_i && \text{[icon]} \\ &= 4.00 \\ \bar{y} &= \frac{1}{17} \sum_{i=1}^{17} y_i && \text{[icon]} \\ &= 3.235\end{aligned}$$

Radial distance: $r_i = \sqrt{(x_i - \bar{x})^2 + (y_i - \bar{y})^2}$


Polar angle: $\theta_i = \tan^{-1} \frac{y_i - \bar{y}}{x_i - \bar{x}}$

where $i = 1, 2, \dots, 17$.




Starting
at
pixel P

Point	theta	r
(7,4)	14	3.1
(7,5)	30	3.48
(6,5)	41	2.67
(5,5)	60	2.03
(4,5)	90	1.77
(3,5)	120	2.03
(2,5)	139	2.67
(2,4)	159	2.14
(1,3)	184	3.01
(1,2)	202	3.24
(2,1)	228	3
(3,1)	246	2.45
(4,1)	270	2.24
(5,1)	294	2.45
(5,2)	309	1.59
(5,3)	347	1.03
(6,3)	353	2.01



centroid =
(4, 3.235)




Minimum radial distance, r_{min} = 1.03

Maximum radial distance, r_{max} = 3.48


Average radial distance, r_{ave} = 2.41

Eccentricity = r_{max}/r_{min} = 3.38

Part (c)



Part (c)




With A as the origin,
equation of AB is $x - 2y = 0$,
 C is the point $(1, 3)$ and F is the point $(4, -1)$

$$\begin{aligned} \text{Length of major axis } a &= \overline{AB} \\ &= \sqrt{3^2 + 6^2} \\ &= 6.71 \end{aligned}$$


$$\begin{aligned} \text{Orientation, } \theta &= \tan^{-1}(3/6) \\ &= 26.6^\circ \end{aligned}$$

$$\begin{aligned} \text{Length of minor axis } b &= \overline{CD} + \overline{EF} \\ &= \frac{|1 - 6|}{\sqrt{1^2 + 2^2}} + \frac{|4 + 2|}{\sqrt{1^2 + 2^2}} \\ &= \frac{11}{\sqrt{5}} \\ &= 4.92 \end{aligned}$$

Question 5



Part (a)



Point	s	ψ
$P+$	0	0°
$Q-$	40	0°
$Q+$	40	90°
R	50	90°
$S-$	81.4	270°
$S+$	81.4	207°
$P-$	103.7	207°

Part (b)

