MiraVista DIAGNOSTICS
Rapid Fungal Testing. Accurate Results.

Medical Test Menu

Antigen Detection

Test Code	Test Name	CPT Code	Specimen Requirements	Specimen Rejection	Shipping	Interfering Substances & Cross-Reactions	TAT
310	**MVista®** *Histoplasma* Ag Quantitative EIA	87385	**Acceptable Specimen:** Urine, serum, CSF, BAL and plasma (EDTA, heparin, or sodium citrate). Serum/plasma separated from the clot, in a leak-proof container. Urine, CSF & BAL in sterile leak-proof container. **Specimen Stability:** **Histo:** 2 weeks RT/RF, indefinitely if FZ **Blasto:** 2 weeks RT/RF, 60 days if FZ **Cocci:** 28 days RT/RF, 60 days if FZ **Minimum Volume**: Urine, Serum, Plasma, CSF, BAL, other body fluids* = 0.8mL	Tissue, biopsy, sputum, bronchial brush, tracheal aspirate, FNA, bone marrow aspirate, or stool. Samples in transport media, fixative or Isolator tubes.	Samples may be shipped on dry ice, frozen ice packs, or ambient.	Sodium hydroxide and sputolysin. Cross-reactions occur with Blastomyces spp., Coccidioides spp., and Paracoccidioides brasiliensis.	Same Day
316	**MVista®** *Blastomyces* Ag Quantitative EIA	87449				When tested in cultures of 10^5 – 10^6 organisms/mL, cross-reactions occurred with Histoplasma spp., Coccidioides spp., Paracoccidioides brasiliensis, Talaromyces marneffei, Aspergillus nidulans, and Candida tropicalis.	Same Day
315	**MVista®** *Coccidioides* Ag Quantitative EIA	87449				Low positive cross-reactions can occur in patient populations with Rheumatoid Arthritis.	Same Day**
309	*Aspergillus* Galactomannan EIA	87305	**Acceptable Specimen**: Serum/EDTA Plasma separated from the clot in a leak-proof container. CSF & BAL in sterile leak-proof container. **Specimen Stability Serum/BAL:** 48hrs RT, 5 days RF or up to 5 months FZ. **Specimen Stability CSF:** 30 days RT/RF, 60 days FZ **Minimum volume**: Serum, EDTA Plasma*, CSF, BAL, other body fluids* = 0.8mL.	Nasal washings, plasma from any tube other than EDTA. Samples in fixative, transport media or PLASMA-LYTE.	Ship on dry ice for next day service. Frozen ice packs may be substituted if sample is shipped the day of collection.	Piperacillin-tazobactam, PLASMA-LYTE or other fluids containing sodium gluconate. Patients must be off Plasma-Lyte for at least 3 days before collection. Infection with *Penicillum, Alternaria, Fusarium Geotrichum, Paecilomyces, Histoplasma, Blastomyces;* or airway colonization with above organisms. In CSF, cross reaction can be seen with *Histoplasma spp, Blastomyces spp, Cryptococcus neoformans,* and *Pseudomonas aeruginosa.*	Same Day
319	Cryptococcal Antigen, Latex Agglutination	86403	**Acceptable Specimen**: Serum separated from the clot in a leak-proof container. CSF in sterile leak-proof container. **Specimen Stability:** 2 months RF/FZ **Minimum volume:** Serum, CSF = 0.25mL. **Positive samples will automatically reflex to a titer.*	Specimen types other than serum or CSF. Samples shipped at ambient temperature.		False-positive results may occur in patients with *Trichosporonosis*.	Same Day
317	(1→3)-β-D-Glucan Colorimetric Assay	87449	**Acceptable Specimen**: Collect serum in red-top or serum separator tube and separate from clot. Pipette into a suitable container that is free of interfering levels of (1→3)-β-D-Glucan. Submit CSF in sterile leak-proof container. **Specimen Stability:** 15 days RF, 27 days FZ **Minimum Volume:** Serum, CSF = 0.25mL.	Lipemic, hemolyzed or icteric samples or any specimen type other than serum or CSF. Samples shipped at ambient temperature.		False-positive results may occur in hemodialysis patients, those receiving therapy with certain fractionated blood products such as serum albumin and immunoglobulins and in specimens of patients exposed to glucan-containing gauzes.	Same Day**

Additional Information:

Two unique patient identifiers are required on specimen containers.

Specimens with particulate matter or viscosity that would not allow the specimen to be pipetted will be rejected.

**Reported with rare specimen comment applied*

**Test Set up: Samples for Antigen testing received by 10:30AM will have same day test set up. Positive samples may require confirmation which could extend TAT. Serum & CSF specimens are pretreated to improve sensitivity which can add one day to TAT.

MiraVista DIAGNOSTICS
Rapid Fungal Testing. Accurate Results.

Medical Test Menu

Antibody Detection

Test Code	Test Name	CPT Code	Specimen Requirements	Specimen Rejection	Shipping	Interfering Substances & Cross-Reactions	TAT
326	**MVista®** *Histoplasma* Antibody IgG, IgM EIA	86698 x2	**Acceptable Specimen**: Serum/Plasma separated from the clot in a leak-proof container. CSF in a sterile leak-proof container. **Specimen Stability:** 28 days RT, 6 months RF, indefinite if FZ. **Minimum volume:** Serum, Plasma*, CSF* = 0.25mL	Specimen types other than serum, plasma, or CSF.	Samples may be shipped on dry ice, frozen ice packs, or ambient.	12% cross reactivity with Blastomycosis. 24% cross reactivity with Coccidioidomycosis.	Same Day
331	**MVista®** *Blastomyces* Antibody IgG EIA	86612	**Acceptable Specimen**: Serum separated from the clot in a leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** Serum & Plasma* = 0.25mL	Specimen types other than serum or plasma.		Specimens positive for Histoplasmosis may cross react and test positive.	Same Day
325	**MVista®** *Coccidioides* Antibody IgG, IgM EIA	86635 x2	**Acceptable Specimen**: Serum/Plasma separated from the clot in a leak-proof container. CSF in a sterile leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** Serum, Plasma*, CSF* = 0.25mL	Specimen types other than serum, plasma, or CSF.		Cross-reactions occur with Aspergillus spp., Histoplasma capsulatum, Paracoccidioides brasiliensis, and Talaromyces marneffei.	Same Day
321	*Histoplasma* Antibody, by Immunodiffusion	86698	**Acceptable Specimen**: Serum/Plasma separated from the clot in a leak-proof container. CSF in a sterile leak-proof container. **Specimen Stability:** 28 days RT, up to 6 months RF, indefinite if FZ. **Minimum volume:** Serum, Plasma*, CSF* = 0.25mL per antibody	Specimen types other than serum, plasma, or CSF.		C reactive protein present in the patient's serum may interfere with interpretation of results for *Aspergillus* Antibody.	3 days from set up date
322	*Blastomyces* Antibody, by Immunodiffusion	86612					
320	*Coccidioides* Antibody, by Immunodiffusion	86635 x2					
324	*Aspergillus* Antibody, by Immunodiffusion	86606					

Additional Information:

Two unique patient identifiers are required on specimen containers.

Specimens with particulate matter or viscosity that would not allow the specimen to be pipetted will be rejected.

**Reported with rare specimen comment applied*

**Test Set up: Samples for Antigen testing received by 10:30AM will have same day test set up. Positive samples may require confirmation which could extend TAT. Serum & CSF specimens are pretreated to improve sensitivity which can add one day to TAT.

MiraVista DIAGNOSTICS
Rapid Fungal Testing. Accurate Results.

Medical Test Menu

Molecular Detection

Test Code	Test Name	CPT Code	Specimen Requirements	Specimen Rejection	Shipping	Interfering Substances & Cross-Reactions	TAT
403	**MVista®** *Histoplasma* DNA, PCR	87798	**Acceptable Specimen**: BAL in sterile leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** BAL, Tracheal Aspirate*, Bronchial Wash* = 0.5mL	Sputum and all other non-respiratory samples	Samples may be shipped on dry ice, frozen ice packs, or ambient.	Specimens positive for *Paracoccidioides brasiliensis*, *Aspergillus niger*, *Aspergillus flavus*, and *Aspergillus nidulans* may cross react and test positive.	Same Day**
404	**MVista®** *Blastomyces* DNA, PCR	87798	**Acceptable Specimen**: BAL in sterile leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** BAL, Tracheal Aspirate*, Bronchial Wash* = 0.5mL			Specimens positive for *Cryptococcus neoformans* may cross react and test positive.	Same Day**
405	**MVista®** *Coccidioides* DNA, PCR	87798	**Acceptable Specimen**: BAL in sterile leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** BAL, Tracheal Aspirate*, Bronchial Wash* = 0.5mL			No cross-reactions or interfering substances were identified during validation.	Same Day**
402	**MVista®** *Pneumocystis* DNA, PCR	87798	**Acceptable Specimen**: BAL in sterile leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** BAL, Tracheal Aspirate*, Bronchial Wash* = 0.5mL			No cross-reactions or interfering substances were identified during validation.	Same Day**

Additional Information:

Two unique patient identifiers are required on specimen containers.

Specimens with particulate matter or viscosity that would not allow the specimen to be pipetted will be rejected.

**Reported with rare specimen comment applied*

**Test Set up: Samples for Antigen testing received by 10:30AM will have same day test set up. Positive samples may require confirmation which could extend TAT. Serum & CSF specimens are pretreated to improve sensitivity which can add one day to TAT.

MiraVista DIAGNOSTICS
Rapid Fungal Testing. Accurate Results.

Medical Test Menu

Panels

Test Code	Test Name	CPT Code	Specimen Requirements	Specimen Rejection	Shipping	Included Tests and Samples	TAT
1000	**MVista®** Pulmonary Fungal PCR Panel	87798 x4	**Acceptable Specimen**: BAL in sterile leak-proof container. **Specimen Stability:** 30 days RT, RF, or FZ. **Minimum volume:** BAL, Tracheal Aspirate*, Bronchial Wash* = 0.5mL	Sputum and all other non-respiratory samples	Samples may be shipped on dry ice, frozen ice packs, or ambient.	Histo 403 Blasto 404 Cocci 405 Pneumo 402	Same Day**

Additional Information:

Two unique patient identifiers are required on specimen containers.

Specimens with particulate matter or viscosity that would not allow the specimen to be pipetted will be rejected.

**Reported with rare specimen comment applied*

**Test Set up: Samples for Antigen testing received by 10:30AM will have same day test set up. Positive samples may require confirmation which could extend TAT. Serum & CSF specimens are pretreated to improve sensitivity. This adds one day to TAT.

ABBREVIATIONS

TAT: Turnaround time
BAL: Bronchial Lavage
CSF: Cerebral Spinal Fluid
RF: Refrigerated
FZ: Frozen
RT: Room Temperature, Ambient