

# **2-GROUP BELYI MAPS**

A Thesis

Submitted to the Faculty

in partial fulfillment of the requirements for the

degree of

Doctor of Philosophy

in

Mathematics

by Michael James Musty

Guarini School of Graduate and Advanced Studies

DARTMOUTH COLLEGE

Hanover, New Hampshire

June 14, 2019

Examining Committee:

---

John Voight, Chair

---

Thomas Shemanske

---

Carl Pomerance

---

David P. Roberts

---

F. Jon Kull, Ph.D.

Dean of the Guarini School of Graduate and Advanced Studies



# Abstract

This thesis concerns the explicit computation of Galois Belyi maps  $\phi: X \rightarrow \mathbb{P}^1$  with monodromy group a 2-group. The motivation behind computing these so-called *2-group Belyi maps* comes from Beckmann's theorem [1] which relates the primes of bad reduction of the algebraic curve  $X$  to the primes dividing the monodromy group of  $\phi$ . The computation has two parts. The first is a combinatorial computation to enumerate the isomorphism classes of 2-group Belyi maps. This computation provides evidence to a conjecture that every 2-group Belyi map is defined over an abelian extension of  $\mathbb{Q}$ , and motivates a partial proof to the conjecture for dihedral monodromy groups. The second part is an explicit algorithm to compute equations for the curve  $X$ . All computations are carried out using **Magma** [7] and the source code written by the author is available at [24].

# Contents

Abstract . . . . .	ii
<b>1 Introduction</b>	<b>1</b>
1.1 Motivation . . . . .	1
1.2 Main results . . . . .	4
1.3 Navigation . . . . .	6
<b>2 Background on Belyi maps</b>	<b>7</b>
2.1 Belyi maps and Galois Belyi maps . . . . .	7
2.2 Permutation triples and passports . . . . .	10
2.3 Triangle groups . . . . .	12
2.4 Fields of moduli and fields of definition . . . . .	14
<b>3 Group theory</b>	<b>16</b>
3.1 2-groups . . . . .	16
3.2 Examples of 2-groups . . . . .	19
3.3 Computing group extensions . . . . .	21
3.4 An iterative algorithm to produce generating triples . . . . .	32
3.5 Results of computations . . . . .	41

<b>4</b>	<b>Fields of definition of 2-group Belyi maps</b>	<b>50</b>
4.1	Refined passports . . . . .	50
4.2	A refined conjecture . . . . .	52
<b>5</b>	<b>Computing equations</b>	<b>59</b>
5.1	Quadratic extensions of number fields . . . . .	59
5.2	Curves and algebraic function fields . . . . .	61
5.3	Quadratic extensions of function fields . . . . .	65
5.4	An algorithm over $\mathbb{F}_q$ . . . . .	66
5.5	An implementation over $\mathbb{Q}^{\text{al}}$ . . . . .	76
5.6	Results of computations . . . . .	79
5.7	Naming conventions for database examples . . . . .	80
5.8	Degree 2 . . . . .	82
5.9	Degree 4 . . . . .	83
5.10	Degree 8 . . . . .	84
5.11	Degree 16 . . . . .	87
5.12	Degree 32 . . . . .	89
	<b>References</b>	<b>92</b>

# List of Figures

3.5.7 Distribution of genera up to degree 256 . . . . .	46
3.5.7 # nonhyperbolic and hyperbolic passports by degree (left), and # nonhyperbolic and hyperbolic lax passports by degree (right). . . . .	47
3.5.7 # permutation triples by degree with abelian and nonabelian mon- odromy groups (left) and # permutation triples by degree with mon- odromy groups of various nilpotency classes (right). . . . .	48
3.5.7 # passports of various sizes by degree . . . . .	49

---

## Chapter 1

---

# Introduction

### Section 1.1

### Motivation

A broad goal of arithmetic geometry is to use tools from algebraic geometry to study arithmetic questions that arise in number theory. One example of this connection is the theorem of Faltings, which bounds the number of rational points (an arithmetic quantity) according to the genus  $g \geq 2$  of a nonsingular algebraic curve (a geometric quantity). Another example of this connection is a theorem due to Belyi which states that an algebraic curve  $X$  over the complex numbers can be defined by equations with coefficients in a number field if and only if  $X$  admits a **Belyi map** a finite cover  $\phi: X \rightarrow \mathbb{P}^1$  unramified outside three points. The way in which Belyi maps capture precisely when a transcendental object is also an algebraic object is just one of the remarkable properties of these covers. The goal of this thesis is to exploit these properties in the particular setting which we now describe.

We begin with a motivating example. Let  $E$  be an elliptic curve over  $\mathbb{Q}$ , let  $\ell \in \mathbb{Z}$

## 1.1 MOTIVATION

---

be prime, and let  $G_{\mathbb{Q}} := \text{Gal}(\mathbb{Q}^{\text{al}} | \mathbb{Q})$  be the absolute Galois group of  $\mathbb{Q}$ . There is an action of  $G_{\mathbb{Q}}$  on the  $\ell$ -torsion points  $E[\ell](\mathbb{Q}^{\text{al}}) \cong \mathbb{Z}/\ell\mathbb{Z} \times \mathbb{Z}/\ell\mathbb{Z}$  of  $E$ , which determines a 2-dimensional mod- $\ell$  Galois representation

$$\rho: G_{\mathbb{Q}} \rightarrow \text{Aut}(E[\ell]) \cong \text{GL}_2(\mathbb{Z}/\ell\mathbb{Z}). \quad (1.1.1)$$

The kernel of this representation fixes the field  $\mathbb{Q}(E[\ell])$  the  $\ell$ -torsion field of  $E$  obtained by adjoining the coordinates of all  $\ell$ -torsion points of  $E$ . The geometry of  $E$  and the arithmetic of  $\rho$  are intimately related. For example, if  $E$  has good reduction at a prime  $p \neq \ell$ , then  $p$  will be unramified in the field  $\mathbb{Q}(E[\ell])$  by the criterion of Néron-Ogg-Shafarevich.

This relationship between curves and Galois representations extends to higher genus curves. Let  $X$  be an irreducible, smooth projective curve of genus  $g \geq 1$  over a number field  $K$ . The Jacobian variety of  $X$ ,  $J := \text{Jac}(X)$ , is an abelian variety over  $K$  of dimension  $2g$ . Again the  $\ell$ -torsion points  $J[\ell]$  of  $J$  define a mod- $\ell$  Galois representation and a number field  $K(J[\ell])$ . As was the case for elliptic curves, if  $X$  has good reduction at a prime  $\mathfrak{p}$  in  $K$ , then  $\mathfrak{p}$  is unramified in the  $\ell$ -torsion field  $K(J[\ell])$ .

The application of Belyi maps to this situation comes from Beckmann's theorem [1.1.2](#). To state Beckmann's theorem requires a bit more terminology. Associated to every Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  is the **monodromy group** of the covering obtained by lifting paths around the ramification points on  $\mathbb{P}^1$ . We say that a Belyi map is **Galois** if the covering is Galois (equivalently if the degree of the cover equals the size of the monodromy group). We can now state Beckmann's theorem.

**Theorem 1.1.2** (Beckmann [\[1\]](#)). *Let  $\phi: X \rightarrow \mathbb{P}^1$  be a Galois Belyi map with mon-*



## 1.1 MOTIVATION

---

odromy group  $G$  and suppose  $p$  does not divide  $\#G$ . Then there exists a number field  $M$  with the following properties:

- $p$  is unramified in  $M$
- the Belyi map  $\phi$  is defined over  $M$
- the Belyi curve  $X$  is defined over  $M$
- $X$  has good reduction at all primes  $\mathfrak{p}$  of  $M$  above  $p$

If we insist that  $G$  is a 2-group in Beckmann's theorem, then we can hope to find fields  $M$  and  $M(\text{Jac}(X)[2])$  unramified away from 2. The main interest in finding such a field comes from a conjecture (now theorem) of Gross.

**Conjecture 1.1.3** (Gross). *For every prime  $p$ , there exists a nonsolvable Galois number field ramified only at  $p$ .*

Although Gross's conjecture is now resolved, the only explicit example of such a field for  $p < 11$  is given in [27] for  $p = 5$ . A long-term application of the work in this thesis is to find an explicit nonsolvable number field ramified only at 2. The existence of such a field was proved in [11] using techniques unrelated to Belyi maps.

But first, to use Beckmann's theorem to construct interesting number fields, we must have explicit Galois Belyi maps with monodromy group a 2-group. The explicit construction of these objects is the focus of this thesis, and it is these objects we refer to as 2-group Belyi maps. We now summarize the results of this thesis concerning 2-group Belyi maps.

## Section 1.2

**Main results**

Motivated by the discussion in Section 1.1, this work aims to address the task of explicitly computing 2-group Belyi maps up to isomorphism. This computation consists of two main parts:

- enumerating isomorphism classes
- computing explicit equations for each isomorphism class

Let  $d \in \mathbb{Z}_{\geq 1}$ . Isomorphism classes of degree  $d$  Belyi maps are in bijection with the set of transitive permutation triples up to simultaneous conjugation in the symmetric group  $S_d$ . A **transitive permutation triple** is a triple of permutations  $\sigma := (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  that multiply to the identity and generate a transitive subgroup of  $S_d$ . We say that two permutation triples  $\sigma, \sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  such that

$$\sigma^\tau := (\tau^{-1}\sigma_0\tau, \tau^{-1}\sigma_1\tau, \tau^{-1}\sigma_\infty\tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty) = \sigma'. \quad (1.2.1)$$

The first main result of this thesis is an explicit algorithm (see Algorithm 3.4.28 in Section 3.4) to compute permutation triples corresponding to all 2-group Belyi maps up to a given degree. We use this algorithm (implemented in **Magma** [7]) all such permutation triples up to degree 256. The algorithms in Section ?? are used to prove results such as the following theorem.

**Theorem 1.2.2.** *The following table lists the number of isomorphism classes of per-*

## 1.2 MAIN RESULTS

---

*mutation triples corresponding to 2-group Belyi maps of degree  $d$  up to 256.*

$d$	1	2	4	8	16	32	64	128	256
$\#$ permutation triples	1	3	7	19	55	151	503	1799	7175

(1.2.3)

Other results of this type are detailed in Section 3.5. Having explicit permutation triples allows us to apply techniques from [25] to prove the following theorem.

**Theorem 1.2.4.** *Every 2-group Belyi map of degree  $d \leq 256$  is defined over an abelian number field.*

The beginning of Section 4.2 explains the above theorem and then pivots to the following conjecture.

**Conjecture 1.2.5.** *Every 2-group Belyi map is defined over an abelian number field.*

In addition to verifying this conjecture for  $d \leq 256$ , we were able to prove this conjecture for 2-group Belyi maps with monodromy group abelian (Lemma 4.2.9) or dihedral (Theorem 4.2.12).

The rest of the results of this thesis pertain to computing equations of 2-group Belyi maps. The first of these is an algorithm to compute 2-group Belyi maps over a finite field  $\mathbb{F}_q$  where  $q = p^k$  and  $p \neq 2$  (see Section 5.4). This algorithm has been implemented in **Magma** and used to construct a database of all 2-group Belyi maps (up to isomorphism) over  $\mathbb{F}_3^{\text{al}}$  up to degree 32.

The other main result is an implementation (similar to the algorithm over  $\mathbb{F}_q$ ) in characteristic zero (see Section 5.5). Although the characteristic zero implementation does not succeed in all cases, it does work well in practice. In particular, the **Magma** implementation succeeded to compute hundreds of 2-group Belyi maps over  $\mathbb{Q}^{\text{al}}$  up to degree 256.

The rest of Chapter 5 is devoted to describing the computations along with interesting examples encountered along the way.

### Section 1.3

## Navigation

Having motivated and stated the main results, we now provide some explanation of how this thesis is organized and where to find details pertaining to the main results.

Chapter 2 details some of the necessary background material related to Belyi maps, permutation triples, and function fields.

Chapter 3 describes an algorithm to enumerate the isomorphism classes of 2-group Belyi maps using permutation triples (see Algorithm 3.4.11 and Algorithm 3.4.28). These algorithms have been used to enumerate all isomorphism classes of 2-group Belyi maps with degree up to 256. The results of these computations are detailed in Section 3.5.

In Chapter 4 we apply the results of computations in Chapter 3 to provide evidence for a conjecture that all 2-group Belyi maps are defined over a cyclotomic field.

Chapter 5 discusses an algorithm to compute explicit equations for 2-group Belyi maps over finite fields with characteristic not 2 (see Algorithm 5.4.9, Algorithm 5.4.13, and Algorithm 5.4.15). Algorithm 5.5.1 describes the modifications to in characteristic zero.

The source code for the implementation used in this thesis can be found at [24].

---

## Chapter 2

---

# Background on Belyi maps

### Section 2.1

#### Belyi maps and Galois Belyi maps

We now set up the framework to discuss the main mathematical objects of interest in this work.

**Definition 2.1.1.** A Belyi map is a morphism  $\phi: X \rightarrow \mathbb{P}^1$  of smooth projective algebraic curves over  $\mathbb{C}$  that is unramified outside  $\{0, 1, \infty\}$ . We define the **genus** of  $\phi$  to be the genus of  $X$ .

**Definition 2.1.2.** Two Belyi maps  $\phi: X \rightarrow \mathbb{P}^1$  and  $\phi': X' \rightarrow \mathbb{P}^1$  are **isomorphic** if there exists an isomorphism of curves from  $X$  to  $X'$  such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{\sim} & X' \\ & \searrow \phi & \swarrow \phi' \\ & \mathbb{P}^1 & \end{array} \tag{2.1.3}$$

commutes. If instead we only insist that the isomorphism makes a diagram

$$\begin{array}{ccc} X & \xrightarrow{\sim} & X' \\ \phi \downarrow & & \downarrow \phi' \\ \mathbb{P}^1 & \xrightarrow[\beta]{\sim} & \mathbb{P}^1 \end{array} \quad (2.1.4)$$

commute, with the bottom map  $\beta$  satisfying  $\beta(\{0, 1, \infty\}) = \{0, 1, \infty\}$ , then we say that  $\phi$  and  $\phi'$  are **lax isomorphic**.

**Definition 2.1.5.** The triple of partitions  $(\lambda_0, \lambda_1, \lambda_\infty)$  encoding the ramification above 0, 1, and  $\infty$  is called the **ramification type** of  $\phi$ .

Let  $\phi: X \rightarrow \mathbb{P}^1$  be a Belyi map of degree  $d$ . Once we label the sheets of the cover and pick a basepoint  $\star \notin \{0, 1, \infty\}$ , we obtain a homomorphism

$$h: \pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \star) \rightarrow S_d \quad (2.1.6)$$

by lifting paths around the branch points of  $\phi$ .

**Definition 2.1.7.** The image of  $h$  in (2.1.6) is the **monodromy group** of  $\phi$  denoted  $\text{Mon}(\phi)$ .

**Definition 2.1.8.** A Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  is **defined over** a number field  $K \subseteq \mathbb{C}$  if the defining equations of  $\phi$  and  $X$  can be described by polynomials expressions with coefficients in  $K$ . We say that  $K$  is a **field of definition** for  $\phi$ .

**Theorem 2.1.9** (Belyi's theorem [2]). *An algebraic curve  $X$  over  $\mathbb{C}$  can be defined over a number field if and only if  $X$  admits a Belyi map.*

Belyi's theorem implies that every Belyi map can be described by a morphism  $\phi: X \rightarrow \mathbb{P}^1$  of algebraic curves defined over a number field  $K \subseteq \mathbb{C}$  (instead of over  $\mathbb{C}$ ).

Since maps of curves correspond to function field extensions, we can again rephrase a Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  (defined over  $K$ ) by an extension of function fields  $K(X) \supseteq K(\mathbb{P}^1)$ .  $K(\mathbb{P}^1)$  is isomorphic to the field of rational functions (referred to as the **rational function field** of  $K$ ) in one variable, say  $K(x)$ , and for  $K$  perfect  $K(X)$  can be written as  $K(x)(\alpha)$  for some primitive element  $\alpha$ .

The degree of a Belyi map in this setting is the degree of the corresponding function field extension  $K(X)$  over the rational function field. Ramification in this setting corresponds to the factorization of ideals  $(x)$ ,  $(x - 1)$ , and  $(1/x)$  in maximal orders of  $K(X)$ . The monodromy group in this setting corresponds to field automorphisms of the Galois closure of  $K(X)$  fixing  $K(x)$ .

Let  $K^{\text{al}}$  denote an algebraic closure of  $K$  in  $\mathbb{C}$ .

**Definition 2.1.10.** A Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  defined over  $K$  is (geometrically) Galois if the corresponding function field extension  $K^{\text{al}}(X)$  is a Galois field extension over the rational function field  $K^{\text{al}}(x)$ .

When  $\phi$  is Galois, the ramification type of  $\phi$  can more simply be encoded by a triple of integers  $(a, b, c) \in \mathbb{Z}_{\geq 1}^3$ . When  $\phi$  is a Galois Belyi map, we can identify  $\text{Mon}(\phi)$  in Definition 2.1.7 as the Galois group  $\text{Gal}(K^{\text{al}}(X) | K^{\text{al}}(\mathbb{P}^1))$ . For this reason, we may also write  $\text{Gal}(\phi)$  to denote  $\text{Mon}(\phi)$  when  $\phi$  is Galois.

We can now define the main object of interest in this thesis.

**Definition 2.1.11.** A 2-group Belyi map is a Galois Belyi map of degree  $d$  with monodromy group a 2-group of order  $d$ . Necessarily,  $\text{Mon}(\phi) = \text{Gal}(\phi) \subseteq S_d$  is the regular representation.

## Section 2.2

## Permutation triples and passports

**Definition 2.2.1.** A permutation triple of degree  $d \in \mathbb{Z}_{\geq 1}$  is a tuple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  such that  $\sigma_\infty \sigma_1 \sigma_0 = 1$ . A permutation triple is **transitive** if the subgroup  $\langle \sigma \rangle \leq S_d$  generated by  $\sigma$  is transitive. We say that two permutation triples  $\sigma, \sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  such that

$$\sigma^\tau := (\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty) = \sigma'. \quad (2.2.2)$$

An **automorphism** of a permutation triple  $\sigma$  is an element of  $S_d$  that simultaneously conjugates  $\sigma$  to itself, i.e.,  $\text{Aut}(\sigma) = C_{S_d}(\langle \sigma \rangle)$ , the centralizer inside  $S_d$ .

**Lemma 2.2.3.** *The set of transitive permutation triples of degree  $d$  up to simultaneous conjugation is in bijection with the set of Belyi maps of degree  $d$  up to isomorphism.*

*Proof.* The correspondence is via monodromy [20, Lemma 1.1]; in particular, the monodromy group of a Belyi map is (conjugate in  $S_d$  to) the group generated by  $\sigma$ .  $\square$

The group  $G_{\mathbb{Q}} := \text{Gal}(\mathbb{Q}^{\text{al}} | \mathbb{Q})$  acts on Belyi maps by acting on the coefficients of a set of defining equations; under the bijection of Lemma 2.2.3, it thereby acts on the set of transitive permutation triples, but this action is rather mysterious. We can cut this action down to size by identifying some basic invariants, as follows.

**Definition 2.2.4.** A **passport** consists of the data  $\mathcal{P} = (g, G, \lambda)$  where  $g \geq 0$  is an integer,  $G \leq S_d$  is a transitive subgroup, and  $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$  is a tuple of partitions  $\lambda_s$  of  $d$  for  $s = 0, 1, \infty$ . These partitions will be also be thought of as a tuple of conjugacy classes  $C = (C_0, C_1, C_\infty)$  by cycle type, so we will also write passports as



$(g, G, C)$ . Two passports  $(g, G, C)$  and  $(g', G', C')$  are **equal** if  $g = g'$ ,  $C = C'$ , and  $G$  is conjugate to  $G'$ .

**Definition 2.2.5.** The **passport** of a Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  is

$$\mathcal{P}(\phi) = (g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty)) \quad (2.2.6)$$

where  $g(X)$  is the genus of  $X$  and  $\lambda_s$  is the partition of  $d$  obtained by the ramification degrees above  $s = 0, 1, \infty$ , respectively.

**Definition 2.2.7.** The **passport** of a transitive permutation triple  $\sigma$  is

$$\mathcal{P}(\sigma) = (g(\sigma), \langle \sigma \rangle, \lambda(\sigma)) \quad (2.2.8)$$

where (by Riemann–Hurwitz)

$$g(\sigma) := 1 - d + (e(\sigma_0) + e(\sigma_1) + e(\sigma_\infty))/2 \quad (2.2.9)$$

and  $e$  is the index of a permutation ( $d$  minus the number of orbits), and  $\lambda(\sigma)$  is the cycle type of  $\sigma_s$  for  $s = 0, 1, \infty$ .

**Definition 2.2.10.** The **size** of a passport  $\mathcal{P}$  is the number of simultaneous conjugacy classes as in (2.2.2) of (necessarily transitive) permutation triples  $\sigma$  with passport  $\mathcal{P}$ .

The action of  $G_{\mathbb{Q}}$  on Belyi maps preserves passports. Therefore, after computing equations for all Belyi maps with a given passport, we can try to identify the Galois orbits of this action.

**Definition 2.2.11.** We say a passport is **irreducible** if it has one  $G_{\mathbb{Q}}$ -orbit and **reducible** otherwise.

We finish this section with an observation about ramification and the Riemann-Hurwitz formula in the case where we have a Galois Belyi map.

**Lemma 2.2.12.** *Let  $\sigma$  be a degree  $d$  permutation triple corresponding to  $\phi: X \rightarrow \mathbb{P}^1$  a Galois Belyi map with monodromy group  $G$  and  $m_s$  be the order of  $\sigma_s$  for  $s \in \{0, 1, \infty\}$ . Then  $\sigma_s$  consists of  $d/m_s$  many  $m_s$ -cycles. In particular, for a 2-group Belyi map,  $m_s$  and  $\#G$  are powers of 2.*

*Proof.* This follows from the condition that the field extension  $K(X)$  is Galois over the rational function field  $K(x)$ . The Galois action is transitive on primes above any prime of  $K(x)$  and in particular implies that the ramified primes all have the same ramification index if they lie above the same prime of  $K(x)$ .  $\square$

Lemma 2.2.12 allows for a simplified version of the Riemann-Hurwitz formula for Galois Belyi maps.

**Theorem 2.2.13** (Riemann-Hurwitz). *Let  $\sigma$  be a degree  $d$  permutation triple corresponding to  $\phi: X \rightarrow \mathbb{P}^1$  a Galois Belyi map with monodromy group  $G$ . Let  $a, b, c$  be the orders of  $\sigma_0, \sigma_1, \sigma_\infty$  respectively. Then*

$$g(X) = 1 + \frac{\#G}{2} \left( 1 - \frac{1}{a} - \frac{1}{b} - \frac{1}{c} \right). \quad (2.2.14)$$

### Section 2.3

## Triangle groups

**Definition 2.3.1.** Let  $(a, b, c) \in \mathbb{Z}_{\geq 1}^3$ . If  $1 \in (a, b, c)$ , then we say the triple is degenerate. Otherwise, we call the triple **spherical**, **Euclidean**, or **hyperbolic** according

to whether the value of

$$\chi(a, b, c) = \frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1 \quad (2.3.2)$$

is positive, zero, or negative. We call this the **geometry type** of the triple. We associate the geometry

$$H = \begin{cases} \mathbb{P}^1 & \chi(a, b, c) > 0 \\ \mathbb{C} & \chi(a, b, c) = 0 \\ \mathfrak{H} & \chi(a, b, c) < 0 \end{cases} \quad (2.3.3)$$

where  $\mathfrak{H}$  denotes the complex upper half-plane.

**Definition 2.3.4.** For each triple  $(a, b, c)$  in Definition 2.3.1 we define the **triangle group**

$$\Delta(a, b, c) = \langle \delta_a, \delta_b, \delta_c \mid \delta_a^a = \delta_b^b = \delta_c^c = \delta_c \delta_b \delta_a = 1 \rangle \quad (2.3.5)$$

The **geometry type** of a triangle group  $\Delta(a, b, c)$  is the geometry type of the triple  $(a, b, c)$ .

**Definition 2.3.6.** The **geometry type** of a Galois Belyi map with ramification type  $(a, b, c)$  is the geometry type of  $(a, b, c)$ .

**Definition 2.3.7.** Let  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty)$  be a transitive permutation triple. Let  $a, b, c$  be the orders of  $\sigma_0, \sigma_1, \sigma_\infty$  respectively. The **geometry type** of  $\sigma$  is the geometry type of  $(a, b, c)$ .

The connection between Belyi maps and triangle groups of various geometry types is explained by Lemma 2.3.8.

**Lemma 2.3.8.** *The set of isomorphism classes of degree  $d$  Belyi maps with ramification type  $(a, b, c)$  is in bijection with the set of index  $d$  subgroups  $\Gamma \leq \Delta(a, b, c)$  up*

to conjugation.

For a detailed explanation of this relationship see the first part of Section 1 in [\[20\]](#).

## Section 2.4

### Fields of moduli and fields of definition

We now discuss the background material necessary to describe the conjecture in Chapter 4. This section aims to define a canonical number field associated to a Belyi map which is well-defined on isomorphism classes, bound the degree of this number field, and discuss when a Belyi map can be defined over this field. To start let  $\text{Aut}(\mathbb{C})$  denote the field automorphisms of  $\mathbb{C}$ .

**Definition 2.4.1.** Let  $X$  be an algebraic curve over  $\mathbb{C}$ . The field of moduli of  $X$  is the fixed field of the field automorphisms

$$\{\tau \in \text{Aut}(\mathbb{C}) : \tau(X) \simeq X\} \quad (2.4.2)$$

where  $\tau \in \text{Aut}(\mathbb{C})$  acts on the defining equations of  $X$ . Denote this field as  $M(X)$ .

**Definition 2.4.3.** Let  $\phi: X \rightarrow \mathbb{P}^1$  be a Belyi map. The field of moduli of  $\phi$  is the fixed field of the field automorphisms

$$\{\tau \in \text{Aut}(\mathbb{C}) : \tau(\phi) \cong \phi\} \quad (2.4.4)$$

where  $\tau \in \text{Aut}(\mathbb{C})$  acts on the defining equations of  $\phi$  and isomorphism is determined by Definition 2.1.2. Denote this field as  $M(\phi)$ .

**Theorem 2.4.5.** *Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyi map with passport  $\mathcal{P}(\phi)$ . Then the degree of  $M(\phi)$  is bounded by the size of  $\mathcal{P}(\phi)$ .*

*Proof.* Let  $\tau \in G_{\mathbb{Q}}$  and consider the conjugated map  $\tau(\phi) : \tau(X) \rightarrow \mathbb{P}^1$ . By [19, Appendix]  $\tau(\phi)$  is a Belyi map with  $\mathcal{P}(\phi) = \mathcal{P}(\tau(\phi))$ . Thus  $G_{\mathbb{Q}}$  acts on the set of (isomorphism classes of) Belyi maps with a given passport. The degree of  $M(\phi)$  is bounded by the index of the stabilizer of  $\phi$  in  $G_{\mathbb{Q}}$  under this action, and this index is bounded by the size of  $\mathcal{P}(\phi)$ .  $\square$

Recall from Definition 2.1.8 that a Belyi map  $\phi : X \rightarrow \mathbb{P}^1$  is defined over a number field  $K$  if  $\phi$  and  $X$  can be defined with equations over  $K$ . We say that  $K$  is a **field of definition** for  $\phi$ . For a general Belyi map it may not be possible to define the Belyi map over its field of moduli. However, in the setting we are concerned with this is always possible. For a proof of the following theorem see [10, Proposition 2.5].

**Theorem 2.4.6.** *A Galois Belyi map can always be defined over its field of moduli.*

---

## Chapter 3

---

# Group theory

We begin this chapter with some background on 2-groups and group extensions which we use to explain the algorithms in Section 3.4 on computing explicit permutation triples corresponding to 2-group Belyi maps. We conclude the chapter with Section 3.5 where we summarize the computation of all permutation triples corresponding to 2-group Belyi maps up to degree 256. We also do some coarse data analysis of these results.

### Section 3.1

## 2-groups

In this section we set up some notation and summarize some background material on 2-groups all of which can be found in [13, §6.1].

Let  $G$  be a finite group. Denote the **centralizer** and **normalizer** of a subset  $S \subseteq G$  by  $C_G(S)$  and  $N_G(S)$  respectively. Let  $G$  act on a set  $X$ . For  $x \in X$  denote the **stabilizer** of  $x$  by  $\text{stab}_x(G)$  and the **orbit** of  $x$  by  $\text{orb}_x(G)$ .

### 3.1 2-GROUPS

---

**Definition 3.1.1.** Let  $p \in \mathbb{Z}$  be prime. A finite group  $G$  is a  $p$ -group if the cardinality of  $G$  is a power of  $p$ .

**Lemma 3.1.2.** *The center of a nontrivial  $p$ -group is nontrivial.*

**Lemma 3.1.3.** *Let  $H$  be a normal subgroup of a  $p$ -group  $G$ . Let  $C$  be a conjugacy class of  $G$ . Then either  $C \subseteq H$  or  $C \cap H = \emptyset$ .*

**Lemma 3.1.4.** *Let  $G$  be a  $p$ -group. Let  $H$  be a nontrivial normal subgroup of  $G$ . Then  $H$  intersects the center  $Z(G)$  nontrivially.*

**Corollary 3.1.5.** *Let  $H$  be a normal subgroup of order  $p$  of a  $p$ -group  $G$ . Then  $H$  is central.*

**Lemma 3.1.6.** *Let  $H$  be a normal subgroup of a  $p$ -group  $G$ . Let  $\#G = p^\alpha$ . Then  $H$  contains a subgroup  $H_\beta$  of order  $p^\beta$  for every divisor  $p^\beta$  of  $\#H$  with the property that  $H_\beta$  is normal in  $G$  for every  $\beta$ .*

**Lemma 3.1.7.** *Every maximal subgroup  $H$  of a  $p$ -group  $G$  has  $[G : H] = p$  and  $H \trianglelefteq G$ .*

**Definition 3.1.8.** Let  $G$  be a finite group. We define a sequence of subgroups of  $G$  iteratively as follows. Let  $Z_0(G) = \{1\}$  and let  $Z_1(G) = Z(G)$ . For  $i \geq 2$  consider the map

$$\pi: G \rightarrow G/Z_i(G),$$

and define  $Z_{i+1}(G)$  to be the preimage of the center of  $G/Z_i(G)$  under  $\pi$  as follows.

$$Z_{i+1}(G) := \pi^{-1} \left( Z \left( \frac{G}{Z_i(G)} \right) \right)$$

### 3.1 2-GROUPS

---

Continuing this process produces a sequence of characteristic subgroups of  $G$

$$Z_0(G) \trianglelefteq Z_1(G) \trianglelefteq \cdots \trianglelefteq Z_i(G) \trianglelefteq \cdots$$

called the **upper central series** of  $G$ .

**Definition 3.1.9.** For  $x, y \in G$  a finite group, define the **commutator** of  $x$  and  $y$  by  $[x, y] := x^{-1}y^{-1}xy$ . For subgroups  $H, K$  of  $G$  define  $[H, K] := \langle [h, k] : h \in H \text{ and } k \in K \rangle$ . We define the **lower central series** of  $G$  iteratively as follows. Let  $G_0 = G$ , let  $G_1 = [G, G]$ , and for  $i \geq 1$  define  $G_{i+1} = [G, G_i]$ .

**Definition 3.1.10.** A finite group  $G$  is **nilpotent** if the upper central series

$$Z_0(G) \trianglelefteq Z_1(G) \trianglelefteq \cdots \trianglelefteq Z_i(G) \trianglelefteq \cdots$$

has  $Z_c(G) = G$  for some nonnegative integer  $c$ . The integer  $c$  is called the **nilpotency class** of the nilpotent group  $G$ .

**Lemma 3.1.11.** *A finite group  $G$  is nilpotent if and only if  $G^c = \{1\}$  for some nonnegative integer  $c$ . Moreover, the smallest  $c$  such that  $G^c = \{1\}$  is the nilpotency class of  $G$  and*

$$Z_i(G) \leq G^{c-i-1} \leq Z_{i+1}(G)$$

*for all  $i \in \{0, 1, \dots, c-1\}$ .*

**Lemma 3.1.12.** *Every  $p$ -group is nilpotent.*



Section 3.2

## Examples of 2-groups

In this section we define some families of nonabelian 2-groups that we refer to later in the partial proof of Conjecture 4.2.8. These examples are all 2-groups with an index 2 cyclic subgroup. According to Berkovich and Janko ([3, Theorem 1.2]), these groups all have order 2 center, order 4 abelianization, and highest possible nilpotency class.

*Example 3.2.1 (Dihedral).* For  $n \geq 2$  define

$$D_{2^{n+1}} := \langle a, b \mid a^{2^n} = b^2 = 1, bab = a^{-1} \rangle. \quad (3.2.2)$$

We summarize some properties of  $D_{2^{n+1}}$ :

- $D_{2^{n+1}}$  has  $2^{n+1}$  elements which can be written as

$$\{1, a, a^2, \dots, a^{2^n-1}, b, ab, a^2b, \dots, a^{2^n-1}b\}. \quad (3.2.3)$$

- The center  $Z(D_{2^{n+1}}) = \{1, a^{2^{n-1}}\}$  is characteristic.
- $D_{2^{n+1}}$  is a split extension of cyclic groups. There exists an exact sequence

$$1 \longrightarrow Z(D_{2^{n+1}}) \cong \mathbb{Z}/2\mathbb{Z} \longrightarrow D_{2^{n+1}} \longrightarrow \mathbb{Z}/2^n\mathbb{Z} \longrightarrow 1. \quad (3.2.4)$$

- All elements in  $D_{2^{n+1}} \setminus \langle a \rangle$  are involutions.
- The conjugacy classes of  $D_{2^{n+1}}$  are as follows. There are 2 conjugacy classes of

### 3.2 EXAMPLES OF 2-GROUPS

---

size 1, namely  $\{1\}$  and  $\{a^{2^{n-1}}\}$ . There are  $2^{n-1} - 1$  conjugacy classes of size 2,

$$\left\{ \{a^i, a^{-i}\} \right\}_{i=1}^{2^{n-1}-1}. \quad (3.2.5)$$

Finally, there are 2 conjugacy classes of size  $2^{n-1}$ ,

$$\{a^{2^i}b : 0 \leq i \leq 2^{n-1} - 1\} \text{ and } \{a^{2^{i+1}}b : 0 \leq i \leq 2^{n-1} - 1\}. \quad (3.2.6)$$

*Example 3.2.7* (Generalized Quaternion). For  $n \geq 2$  define

$$Q_{2^{n+1}} := \langle a, b \mid a^{2^n} = 1, b^2 = a^{2^{n-1}}, b^{-1}ab = a^{-1} \rangle. \quad (3.2.8)$$

*Example 3.2.9* (Semi-dihedral). For  $n \geq 3$  define

$$SD_{2^{n+1}} := \langle a, b \mid a^{2^n} = b^2 = 1, bab = a^{-1+2^{n-1}} \rangle. \quad (3.2.10)$$

**Lemma 3.2.11.** *Let  $G$  be one of the groups  $D_{2^{n+1}}$ ,  $Q_{2^{n+1}}$ ,  $SD_{2^{n+1}}$  discussed in the previous examples with center  $Z(G)$ . Then  $\#Z(G) = 2$  and  $G/Z(G)$  is a dihedral group.*

*Proof.* This is proved in Berkovich and Janko [3, Theorem 1.2] where they attribute a stronger result to Burnside. □

## Section 3.3

## Computing group extensions

In Section 3.4, we will be interested in constructing 2-groups as (central) extensions of other 2-groups. The computations we rely on are implemented in **Magma** and described in [6]. We now describe the broad strokes of this implementation emphasizing the particular setting we are interested in. The background material concerning group extensions in this section is summarized from [13, §17.4].

**Definition 3.3.1.** Let  $G$  be a finite group and  $A$  a finite abelian group. An **extension** of  $A$  by  $G$  is a group  $\tilde{G}$  such that the sequence

$$1 \longrightarrow A \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} G \longrightarrow 1 \quad (3.3.2)$$

is exact. An extension (3.3.2) is **central** if  $\iota(A)$  is contained in the center of  $\tilde{G}$ .

Note that for a group extension (3.3.2) there is an action of  $G$  on  $\iota(A)$  by conjugation. This action is obtained by choosing a lift in  $\tilde{G}$  and conjugating. Conjugating  $\iota(A)$  by this lift is well-defined since  $A$  is abelian. From now on we identify  $A$  with its image  $\iota(A)$  in  $\tilde{G}$  to ease notation. To keep track of the action of  $G$  on  $A$  we make the following definition.

**Definition 3.3.3.** Let  $G$  be a finite group. A  $G$ -**module** is a finite abelian group  $A$  and a group homomorphism  $\phi: G \rightarrow \text{Aut}(A)$ .

**Proposition 3.3.4.** *The extension in (3.3.2) is central if and only if  $A$  (identified with its image  $\iota(A)$  in  $\tilde{G}$ ) has trivial  $G$ -module structure.*

### 3.3 COMPUTING GROUP EXTENSIONS

---

*Proof.* Let  $a \in A$ , let  $g \in G$ , and let  $\tilde{g} \in \pi^{-1}(g)$ . Then  $g$  acts on  $a$  by

$$g \cdot a = \tilde{g}a\tilde{g}^{-1}, \quad (3.3.5)$$

so the action is trivial if and only if  $a = \tilde{g}a\tilde{g}^{-1}$  for all  $\tilde{g} \in \tilde{G}$  if and only if  $a \in Z(\tilde{G})$ .  $\square$

**Definition 3.3.6.** Two extensions of  $A$  by  $G$  are **equivalent** if there exists an isomorphism of groups  $\phi$  making the diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & A & \longrightarrow & \tilde{G}_1 & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow \text{id} & & \downarrow \phi & & \downarrow \text{id} \\ 1 & \longrightarrow & A & \longrightarrow & \tilde{G}_2 & \longrightarrow & G \longrightarrow 1 \end{array} \quad (3.3.7)$$

commute.

*Remark 3.3.8.* The notion of equivalence from Definition 3.3.6 requires an isomorphism  $\phi$  inducing the identity map on  $A$  and  $G$ . This definition comes from the  $G$ -module structure of  $A$  in the sense that equivalent extensions induce (by conjugation) the same  $G$ -module structure on  $A$ . A weaker notion of equivalence (where we only require  $\phi$  to be any isomorphism from  $A$  to  $A$ ) is useful to characterize the groups  $\tilde{G}$  up to group isomorphism, but will not be used in our situation.

We now look at a motivating example.

*Example 3.3.9.* Let  $A$  be a  $G$ -module with  $\phi: G \rightarrow \text{Aut}(A)$  defining the action of  $G$  on  $A$ . Then we can construct the **(external) semidirect product**  $A \rtimes G$  which is the set  $A \times G$  equipped with multiplication defined by

$$(a_1, g_1)(a_2, g_2) := (a_1 + \phi(g_1)(a_2), g_1g_2).$$

### 3.3 COMPUTING GROUP EXTENSIONS

---

Then  $A \rtimes G$  is an extension of  $A$  by  $G$

$$1 \longrightarrow A \xrightarrow{\iota} A \rtimes G \xrightarrow{\pi} G \longrightarrow 1$$

where the conjugation action of  $\pi^{-1}(G)$  on  $A$  (identified with  $\iota(A)$ ) coincides with the original  $G$ -module action of  $A$ .

We now explain the bijection between equivalence classes of extensions (of  $A$  by  $G$ ) and elements of the group  $H^2(G, A)$ . The latter can be efficiently computed in **Magma** [6], and is a crucial part of the algorithms in Section 3.4.

**Definition 3.3.10.** A function  $s: G \rightarrow \tilde{G}$  such that  $\pi \circ s = \text{id}_G$  is called a **section** of  $\pi$ . A section is **normalized** if it maps  $\text{id}_G$  to  $\text{id}_{\tilde{G}}$ . An extension is **split** if there exists a section  $s$  such that  $s$  is a homomorphism.

**Proposition 3.3.11.** *The extension in (3.3.2) is split if and only if it is equivalent to*

$$1 \longrightarrow A \xrightarrow{\iota'} A \rtimes G \xrightarrow{\pi'} G \longrightarrow 1$$

where  $A \rtimes G$  is the semidirect product of  $G$  and  $A$  relative to the given action described in Example 3.3.9.

*Proof.* Suppose  $\phi: \tilde{G} \rightarrow A \rtimes G$  is an isomorphism inducing the identity maps on  $A$  and  $G$ . Let  $s': G \rightarrow A \rtimes G$  be the section  $g \mapsto (\text{id}_A, g)$ . Then the section  $s := \phi^{-1}s'$  is a group homomorphism  $s: G \rightarrow \tilde{G}$  showing the extension is split. Conversely, assume there exists a section  $s: G \rightarrow \tilde{G}$  which is a group homomorphism. Then the map  $\phi: A \rtimes G \rightarrow \tilde{G}$  defined by

$$(a, g) \mapsto \iota(a)s(g)$$

### 3.3 COMPUTING GROUP EXTENSIONS

---

is a bijection. We now show that this map is a group isomorphism by analyzing the multiplication of two elements in the image of  $\phi$ . Let  $\iota(a)s(g)$  and  $\iota(a')s(g')$  in the image of  $\phi$ . Then from the  $G$ -module structure of  $A$  we have

$$s(g)\iota(a') = \iota(ga')s(g). \quad (3.3.12)$$

(3.3.12) then implies

$$\iota(a)s(g)\iota(a')s(g') = \iota(a)\iota(ga')s(g)s(g') = \iota(a + ga')s(gg')$$

which is precisely the semidirect product multiplication rule on  $A \times G$ .  $\square$

Proposition 3.3.11 completely describes split extensions. For nonsplit extensions, we must analyze sections that are not homomorphisms. To measure the failure of  $s$  to be a homomorphism, we make the following definition.

**Definition 3.3.13.** Let  $s$  be a section of an extension (3.3.2). Let  $f: G \times G \rightarrow A$  be defined by the equation

$$s(g)s(h) = \iota(f(g, h))s(gh). \quad (3.3.14)$$

In other words,  $\pi(s(gh)) = \pi(s(g)s(h)) = gh$ , so we know that  $s(gh)$  and  $s(g)s(h)$  differ by an element of  $\iota(A)$ . We define  $f(g, h)$  to be the element  $a \in A$  such that (3.3.14) is satisfied. The function  $f$  is called the **factor set** for the extension and the section  $s$ . A factor set is **normalized** if  $s$  is normalized. A normalized factor set  $f$  satisfies

$$f(g, 1) = f(1, g) = 0$$

for all  $g \in G$ .

### 3.3 COMPUTING GROUP EXTENSIONS

---

In Lemma 3.3.20 we will see that a factor set for an extension with a section is a special case of a 2-cocycle which we now define.

**Definition 3.3.15.** A 2-cocycle is a map  $f: G \times G \rightarrow A$  satisfying

$$f(g, h) + f(gh, k) = gf(h, k) + f(g, hk) \quad (3.3.16)$$

for all  $g, h, k \in G$ . A 2-cocycle  $f$  is normalized if

$$f(g, 1) = f(1, g) = 0$$

for all  $g \in G$ .

**Definition 3.3.17.** A 2-coboundary is a map  $f: G \times G \rightarrow A$  such that there exists  $f_1: G \rightarrow A$  satisfying

$$f(g, h) = gf_1(h) - f_1(gh) + f_1(g) \quad (3.3.18)$$

for all  $g, h \in G$ .

**Definition 3.3.19.** Let  $Z^2(G, A)$  denote the set of 2-cocycles and  $B^2(G, A)$  denote the set of all 2-coboundaries. The second cohomology group  $H^2(G, A)$  is defined by the quotient  $Z^2(G, A)/B^2(G, A)$ .

**Lemma 3.3.20.** The factor set  $f$  of an extension as in (3.3.2) and a section  $s$  is a 2-cocycle.

**Lemma 3.3.21.** Consider an extension as in (3.3.2). Let  $s$  and  $s'$  be sections of this extension with corresponding factor sets  $f$  and  $f'$  respectively. Then  $f' - f$  is a 2-coboundary.

### 3.3 COMPUTING GROUP EXTENSIONS

---

Lemma 3.3.20 and Lemma 3.3.21 are explained on page 825 and 826 of [13].

**Lemma 3.3.22.** *An equivalence class of extensions of  $A$  by  $G$  determine a unique element of  $H^2(G, A)$ .*

*Proof.* Let  $f$  be the factor set for any section of the extension. Lemma 3.3.20 shows that  $f \in Z^2(G, A)$ . Lemma 3.3.21 shows that any other choice of  $f$  corresponding to another choice of section differs from  $f$  by an element of  $B^2(G, A)$ . Thus, any single extension of  $A$  by  $G$  determines a unique cohomology class in  $H^2(G, A)$ . It remains to show that equivalent extensions determine the same element of  $H^2(G, A)$ . Consider the equivalent extensions

$$\begin{array}{ccccccc}
 1 & \longrightarrow & A & \longrightarrow & \tilde{G}_1 & \xrightarrow{\pi_1} & G \longrightarrow 1 \\
 & & \downarrow \text{id} & & \downarrow \phi & & \downarrow \text{id} \\
 1 & \longrightarrow & A & \longrightarrow & \tilde{G}_2 & \xrightarrow{\pi_2} & G \longrightarrow 1.
 \end{array} \tag{3.3.23}$$

and let  $s_1: G \rightarrow \tilde{G}_1$  be a section of  $\pi_1$ . From (3.3.23) we have that  $s_2 := \phi \circ s_1$  is a section of  $\pi_2$ . Let  $f_1$  and  $f_2$  be the factor sets corresponding to  $s_1$  and  $s_2$  respectively defined by

$$\begin{aligned}
 s_1(g)s_1(h) &= f_1(g, h)s_1(gh) \\
 s_2(g)s_2(h) &= f_2(g, h)s_2(gh)
 \end{aligned} \tag{3.3.24}$$



### 3.3 COMPUTING GROUP EXTENSIONS

---

for all  $g, h \in G$ . Chasing through the diagram in (3.3.23) we have

$$\begin{aligned}
 s_2(g)s_2(h) &= \phi(s_1(g))\phi(s_1(h)) \\
 &= \phi(s_1(g)s_1(h)) \\
 &= \phi(f_1(g, h)s_1(gh)) \\
 &= \phi(f_1(g, h))\phi(s_1(gh)) \\
 &= f_1(g, h)s_2(gh)
 \end{aligned} \tag{3.3.25}$$

where the last equality in (3.3.25) follows from chasing the diagram through the identity map  $\text{id}: A \rightarrow A$ . This shows if two extensions are equivalent, then we can define sections for both extensions such that the corresponding factor sets are the same 2-cocycle. In particular, equivalent extensions define the same element of  $H^2(G, A)$ , which completes the proof.  $\square$

Lemma 3.3.22 proves that any factor set for an extension of  $A$  by  $G$  defines a unique class in  $H^2(G, A)$ . We now discuss the reverse process of constructing an extension of  $A$  by  $G$  from a 2-cocycle.

**Lemma 3.3.26.** *Let  $f \in H^2(G, A)$  for some finite group  $G$  and  $G$ -module  $A$ . Then there is an extension*

$$1 \longrightarrow A \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} G \longrightarrow 1 \tag{3.3.27}$$

*whose factor set is equivalent to  $f$  in  $H^2(G, A)$ .*

*Proof.* Let  $\tilde{G}$  be defined by the set  $A \times G$  equipped with the operation

$$(a_1, g_1)(a_2, g_2) = (a_1 + g_1 a_2 + f(g_1, g_2), g_1 g_2). \tag{3.3.28}$$

### 3.3 COMPUTING GROUP EXTENSIONS

---

$A \times G$  with this operation is a group with identity element  $(-f(1, 1), 1)$  and the inverse of  $(a, g) \in A \times G$  given by

$$(a, g)^{-1} = (-g^{-1}a - f(g^{-1}, g) - f(1, 1), g^{-1}). \quad (3.3.29)$$

We now construct the rest of the extension. Let  $A^*$  be defined by

$$A^* := \{(a - f(1, 1), 1) : a \in A\}. \quad (3.3.30)$$

$A^*$  is a normal subgroup of  $\tilde{G}$  with the inverses given by

$$(a - f(1, 1), 1)^{-1} = (-a - f(1, 1), 1) \quad (3.3.31)$$

The isomorphism  $\iota: A \rightarrow A^*$  is defined by

$$a \mapsto (a - f(1, 1), 1). \quad (3.3.32)$$

Define  $\pi: \tilde{G} \rightarrow G$  by the projection  $(a, g) \mapsto g$ . Now  $A^*$ , the image of  $\iota$ , is contained in  $\ker \pi$  since the second coordinate is  $1 \in G$  for every element of  $A^*$ . Thus (3.3.27) is an extension of  $A$  by  $G$ . Lastly, let  $s: G \rightarrow \tilde{G}$  be a section of  $\pi$  and let  $f_s$  be the factor set of the extension in (3.3.27). One can show that  $f_s$  and  $f$  are equal in  $H^2(G, A)$ .

For more of the details of this proof see page 827 of [13] and page 92 of [8].  $\square$

*Remark 3.3.33.* The construction in Lemma 3.3.26 generalizes the semidirect product construction in Example 3.3.9.

**Theorem 3.3.34.** *There is a bijection between equivalence classes of extensions of  $A$  by  $G$  as in (3.3.2) and elements of  $H^2(G, A)$ .*

*Proof.* The least technical proof is by reducing to the normalized setting and is described in detail on page 826 and page 827 in [13]. Below we give a summary of the proof.

Every 2-cocycle  $f$  has a normalized 2-cocycle in its cohomology class, so without loss of generality we can assume  $f$  is normalized. One then shows that extension constructed from  $f$  using Lemma 3.3.26 has normalized factor set equal to  $f$ . The last step is showing that this procedure does not depend on the choice of normalized 2-cocycle by showing that as long as the normalized 2-cocycles are in the same cohomology class then the corresponding extensions will be equivalent.  $\square$

Having established Theorem 3.3.34, we are interested in computing representatives of  $H^2(G, A)$ . To do this we use the implementation in **Magma** described in [6, Cohomology and group extensions]. Describing this implementation in detail is beyond the scope of this work. Instead, we provide Example 3.3.37 at the end of this section detailing how we use these implementations in practice. In our computation of permutation triples corresponding to 2-group Belyi maps in the next section, we will first be concerned with computing extensions of  $A$  by  $G$  where  $G$  is a finite 2-group and  $A \simeq \mathbb{Z}/2\mathbb{Z}$ . The first consideration in producing these extensions is the possible  $G$ -module structures on  $A$ . Fortunately, the only  $G$ -module structure on  $A$  is the trivial action corresponding to the only homomorphism

$$G \rightarrow \text{Aut}(\mathbb{Z}/2\mathbb{Z}). \tag{3.3.35}$$

### 3.3 COMPUTING GROUP EXTENSIONS

---

According to Theorem 3.3.34, the equivalent extensions of  $A$  by  $G$  correspond to elements of  $H^2(G, A)$  which can be computed efficiently in **Magma** and explicitly converted to group extensions as in Example 3.3.37.

*Remark 3.3.36.* Modifications are required to compute extensions when  $A$  is cyclic of prime order  $p$ . All possible homomorphisms  $G \rightarrow \text{Aut}(\mathbb{Z}/p\mathbb{Z}) \cong (\mathbb{Z}/p\mathbb{Z})^\times$  must be computed, and for each  $G$ -module  $A$ , the corresponding group  $H^2(G, A)$  must also be computed. When  $A$  has more than one cyclic factor, the situation becomes more complicated. For example, the possible  $G$ -module structures on  $A \cong \underbrace{\mathbb{Z}/p\mathbb{Z} \times \cdots \times \mathbb{Z}/p\mathbb{Z}}_{d \text{ times}}$  correspond to irreducible  $\mathbb{F}_p[G]$ -modules of dimension  $d$ . Although **Magma** is capable of computing these modules, we do not require this level of generality for the computations in the next section.

We conclude this section with an example of how we compute group extensions in **Magma**.

*Example 3.3.37.* A file with the source code for this example can be found in the repository [24] and can be run from a shell in the repository as follows.

Shell
<code>magma thesis_examples/group_extensions.m</code>

Let  $\sigma$  be the permutation triple representing the size 1 passport  $(41, G, (16, 2, 8))$  where  $G$  is the permutation group generated by  $\sigma$  of order 256 and small group database label  $(256, 100)$ . Suppressing the permutations in  $\sigma$ , the source code is as follows.

Magma
<code>...</code>
<code>G := sub&lt;Sym(256) sigma&gt;;</code>

### 3.3 COMPUTING GROUP EXTENSIONS

---

```
assert IsTransitive(G);
assert #G eq 256;
A := TrivialModule(G, GF(2));
CM := CohomologyModule(G, A);
H2 := CohomologyGroup(CM, 2);
extensions := [* *];
for h in H2 do
    E_fp, pi_fp, iota_fp := Extension(CM, h);
    iso, E, K := CosetAction(E_fp, sub<E_fp|Id(E_fp)>);
    iotaE := iota_fp*iso;
    piE := (iso^-1)*pi_fp;
    assert Image(iotaE) eq Kernel(piE);
    assert Image(iotaE).1 in Center(E);
    Append(~extensions, [* E, iotaE, piE , h *]);
end for;
```

We first construct  $A$  as a  $G$ -module and construct  $H^2(G, A)$  using the *Cohomology module* functionality in **Magma**. In this example  $\#H^2(G, A) = 32$ . For each cohomology class, we compute the corresponding extension (as a finitely presented group) along with mappings defining the extension. Lastly, we act on the identity coset to obtain the extension as a permutation group along with the appropriate mappings.

Section 3.4

## An iterative algorithm to produce generating triples

The aim of this section is to use techniques to compute group extensions from Section 3.3 to iteratively compute *p-group permutation triples* which we define below.

**Definition 3.4.1.** Let  $p$  be prime. Let  $d \in \mathbb{Z}_{\geq 1}$ . A *p-group permutation triple* of degree  $d$  is a triple of permutations  $\sigma := (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  satisfying

- $\sigma_\infty \sigma_1 \sigma_0 = 1$ ;
- $G := \langle \sigma \rangle$  is a transitive subgroup of  $S_d$ ; and
- $G$  is a *p-group* of order  $d$  embedded in  $S_d$  via its left regular representation.

The group  $G$  is called the **monodromy group** of  $\sigma$ . We say that two *p-group permutation triples*  $\sigma, \sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  such that

$$\sigma^\tau := (\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty) = \sigma'. \quad (3.4.2)$$

*Remark 3.4.3.* In the process of computing extensions of monodromy groups of *p-group Belyi maps* we must pass back and forth between permutation groups and abstract groups given by a presentation. Insisting that  $G$  embeds into  $S_d$  via its regular representation eliminates the ambiguity in embedding a finitely presented group into  $S_d$ . This explains the last property in Definition 3.4.1.

*Example 3.4.4.* When  $d = 1$  we define the triple  $(\text{id}, \text{id}, \text{id}) \in S_1^3$  to be a *p-group permutation triple* for every  $p$ . This is the unique *p-group permutation triple* of

degree 1.

*Example 3.4.5.* Let  $d = p$  and let  $\sigma_s$  be any  $p$ -cycle in  $S_p$ . Then we can write 3 distinct  $p$ -group permutation triples of degree  $p$ :

$$\left(\sigma_s, \sigma_s^{-1}, \text{id}\right), \left(\sigma_s, \text{id}, \sigma_s^{-1}\right), \left(\text{id}, \sigma_s, \sigma_s^{-1}\right). \quad (3.4.6)$$

These are the only  $p$ -group permutation triples of degree  $p$  up to simultaneous conjugation.

We will describe the algorithms in this section in this slightly more general setting even though the  $p = 2$  case is our primary concern.

**Notation 3.4.7.** Let  $\sigma$  be a  $p$ -group permutation triple with monodromy group  $G$  and let  $A \cong \mathbb{Z}/p\mathbb{Z}$  cyclic of prime order. Let  $\tilde{G}$  be an extension of  $A$  by  $G$  sitting in the exact sequence

$$1 \longrightarrow A \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} G \longrightarrow 1. \quad (3.4.8)$$

By Corollary 3.1.5 the image of  $\iota$  is a central subgroup of  $\tilde{G}$ . The algorithm discussed in this section is iterative, and the base case for this iteration is described in Example 3.4.4.

**Definition 3.4.9.** We say that a  $p$ -group permutation triple  $\tilde{\sigma}$  is a **degree  $p$  lift** (or simply a **lift**) of a  $p$ -group permutation triple  $\sigma$  of degree  $d$  if  $\tilde{\sigma}$  is a  $p$ -group permutation triple of degree  $pd$  with monodromy group  $\tilde{G}$  sitting in the exact sequence in (3.4.8) where  $G$  is the monodromy group of  $\sigma$  and  $A \cong \mathbb{Z}/p\mathbb{Z}$ .

**Notation 3.4.10.** In Algorithm 3.4.11, the objective will be to lift a  $p$ -group permutation triple  $\sigma$  of degree  $d$  to  $p$ -group permutation triples  $\tilde{\sigma}$  of degree  $pd$ . We will denote the set of lifts of  $\sigma$  by  $\text{Lifts}(\sigma)$  and write  $\text{Lifts}(\sigma)/\sim$  to denote the equivalence classes of lifts up to simultaneous conjugation in  $S_{pd}$ .

Once we can compute  $\text{Lifts}(\sigma)$ , the next objective is to enumerate all  $p$ -group permutation triples up to a given degree along with the bipartite graph structure determined by lifting triples. More precisely, let  $\mathcal{G}_{p^i}$  denote the bipartite graph with the following node sets.

- $\mathcal{G}_{p^i}^{\text{above}}$  : the set of isomorphism classes of  $p$ -group permutation triples of degree  $p^i$  indexed by permutation triples  $\tilde{\sigma}$  up to simultaneous conjugation in  $S_{p^i}$
- $\mathcal{G}_{p^i}^{\text{below}}$  : the set of isomorphism classes of  $p$ -group permutation triples of degree  $p^{i-1}$  indexed by permutation triples  $\sigma$  up to simultaneous conjugation in  $S_{p^{i-1}}$

The edge set of  $\mathcal{G}_{p^i}$  is defined as follows. For every pair of nodes  $(\tilde{\sigma}, \sigma) \in \mathcal{G}_{p^i}^{\text{above}} \times \mathcal{G}_{p^i}^{\text{below}}$  there is an edge between  $\tilde{\sigma}$  and  $\sigma$  if and only if  $\tilde{\sigma}$  is simultaneously conjugate to a lift of  $\sigma$ .

Now that we have set up some notation and definitions, we now describe the algorithms.

**Algorithm 3.4.11.** Let  $p$  be prime and let  $d \in \mathbb{Z}_{\geq 1}$ .

**Input:**

- $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  a  $p$ -group permutation triple with monodromy group  $G$
- $A$  a  $G$ -module



**Output:**

All degree  $p$  lifts  $\tilde{\sigma}$  of  $\sigma$  up to simultaneous conjugation in  $S_{pd}$  where the induced  $G$ -module structure on  $A$  from the extension in (3.4.8) matches the  $G$ -module structure of  $A$  given as input.

1. Let  $G = \langle \sigma \rangle$  and compute representatives of  $H^2(G, A)$ .
2. For each  $f \in H^2(G, A)$  compute the corresponding extension

$$1 \longrightarrow A \xrightarrow{\iota_f} \tilde{G}_f \xrightarrow{\pi_f} G \longrightarrow 1 \quad (3.4.12)$$

3. For each extension  $\tilde{G}_f$  in (3.4.12) compute the set

$$\text{Lifts}(\sigma, f) := \left\{ \tilde{\sigma} : \tilde{\sigma}_s \in \pi_f^{-1}(\sigma_s) \text{ for } s \in \{0, 1, \infty\}, \tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1, \langle \tilde{\sigma} \rangle = \tilde{G}_f \right\} \quad (3.4.13)$$

4. Let

$$\text{Lifts}(\sigma) := \bigcup_{f \in H^2(G, A)} \text{Lifts}(\sigma, f) \quad (3.4.14)$$

5. Quotient  $\text{Lifts}(\sigma)$  by the equivalence relation  $\sim$  identifying triples in  $\text{Lifts}(\sigma)$  that are simultaneously conjugate, as in (3.4.2), to obtain representatives of  $\text{Lifts}(\sigma)/\sim$ .

*Proof of correctness.* The computation of  $H^2(G, A)$  is described in [6] and implemented in [7]. Theorem 3.3.34 in Section 3.3 implies the following.

- The elements of  $H^2(G, A)$  are in bijection with extensions  $\tilde{G}_f$  as in (3.4.12).
- Any lift of  $\sigma$  inducing the  $G$ -module structure of  $A$  on  $\mathbb{Z}/p\mathbb{Z}$  must have monodromy group sitting in an exact sequence obtained in Step 2.

### 3.4 AN ITERATIVE ALGORITHM TO PRODUCE GENERATING TRIPLES

---

In Step 3 all possible lifts of  $\sigma$  for a single extension  $\tilde{G}_f$  are computed. This is done by computing all  $(\#A)^3$  triples mapping to  $\sigma$  under  $\pi_f$  and checking which satisfy the conditions to be a lift of  $\sigma$ . After collecting all the lifts together in Step 4 it is possible there are simultaneously conjugate  $p$ -group permutation triples in  $\text{Lifts}(\sigma)$ . In Step 5 we quotient by simultaneous conjugation to obtain the desired set of lifts as output.  $\square$

Algorithm 3.4.11 reduces the problem of finding all lifts of a given  $p$ -group permutation triple  $\sigma$  to determining all possible  $\langle\sigma\rangle$ -module structures on  $\mathbb{Z}/p\mathbb{Z}$ . Although computations of this sort are implemented in [7], it is especially easy to do when  $p = 2$ .

**Lemma 3.4.15.** *Let  $G$  be a finite group. The only  $G$ -module structure on  $\mathbb{Z}/2\mathbb{Z}$  is trivial.*

*Proof.* A  $G$ -module structure on  $\mathbb{Z}/2\mathbb{Z}$  is a homomorphism from  $G$  to  $\text{Aut}(\mathbb{Z}/2\mathbb{Z})$ . But  $\text{Aut}(\mathbb{Z}/2\mathbb{Z}) \cong (\mathbb{Z}/2\mathbb{Z})^\times$  which is the trivial group, so there is only one such homomorphism.  $\square$

For the rest of this section we suppose that  $p = 2$ . In this special case, Algorithm 3.4.11 does not require a  $G$ -module as input since (by Lemma 3.4.15) the trivial  $G$ -module structure on  $\mathbb{Z}/2\mathbb{Z}$  can be assumed.

*Remark 3.4.16.* Suppose  $p = 2$  using Notation 3.4.7. Then  $\iota(A)$  is an order 2 normal subgroup of  $\tilde{G}$ . Let  $\alpha$  denote the generator of  $\iota(A)$ . From the perspective of branched covers,  $\alpha$  is identifying  $2d$  sheets in a degree  $2d$  cover down to  $d$  sheets in a degree  $d$  cover. To relate the degree  $2d$  cover corresponding to  $\tilde{G}$  with the degree  $d$  cover corresponding to  $G$  it is convenient to choose  $\alpha$  to be the following product of  $d$

transpositions.

$$\alpha := (1 \ d + 1)(2 \ d + 2) \dots (d - 1 \ 2d - 1)(d \ 2d) \quad (3.4.17)$$

The benefit of following this convention can be seen in Example 3.4.18 where we illustrate Algorithm 3.4.11.

*Example 3.4.18.* In this example we carry out Algorithm 3.4.11 for the degree 2 permutation triple  $\sigma = ((1 \ 2), \text{id}, (1 \ 2))$ . Here  $G = \langle \sigma \rangle \cong \mathbb{Z}/2\mathbb{Z}$ . In Algorithm 3.4.11 Step 2, we obtain two group extensions  $\tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$  and  $\tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}$  sitting in the following exact sequences.

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{Z}/2\mathbb{Z} & \xrightarrow{\iota_1} & \tilde{G}_1 & \xrightarrow{\pi_1} & G \longrightarrow 1 \\ & & & & & & \\ 1 & \longrightarrow & \mathbb{Z}/2\mathbb{Z} & \xrightarrow{\iota_2} & \tilde{G}_2 & \xrightarrow{\pi_2} & G \longrightarrow 1 \end{array} \quad (3.4.19)$$

We will consider the two extensions separately.

- For  $\tilde{G}_1$ , we can look at preimages of  $\sigma_s$  under the map  $\pi_1$  to obtain 4 triples that multiply to the identity:

$$\begin{aligned} & \left\{ ((1 \ 2)(3 \ 4), \text{id}, (1 \ 2)(3 \ 4)), ((1 \ 2)(3 \ 4), (1 \ 3)(2 \ 4), (1 \ 4)(2 \ 3)), \right. \\ & \left. ((1 \ 4)(2 \ 3), \text{id}, (1 \ 4)(2 \ 3)), ((1 \ 4)(2 \ 3), (1 \ 3)(2 \ 4), (1 \ 2)(3 \ 4)) \right\} \end{aligned} \quad (3.4.20)$$

Before we continue with the algorithm, let us take a moment to analyze these triples more closely. The generator  $\alpha$  of  $\iota(\mathbb{Z}/2\mathbb{Z})$  in  $\tilde{G}_1$  is  $(1 \ 3)(2 \ 4)$ . Each triple in (3.4.20) must act on the blocks  $\left\{ \boxed{1 \ 3}, \boxed{2 \ 4} \right\}$  so that the induced permutations of these blocks is the same as the corresponding permutation in  $\sigma$ . For

$$(\tilde{\sigma}_0, \tilde{\sigma}_1, \tilde{\sigma}_\infty) = ((1 \ 2)(3 \ 4), (1 \ 3)(2 \ 4), (1 \ 4)(2 \ 4)) \quad (3.4.21)$$

we have  $\tilde{\sigma}_0(\boxed{13}) = \boxed{24}$  and  $\tilde{\sigma}_0(\boxed{24}) = \boxed{13}$  so that the induced permutation of blocks is

$$(\boxed{13}, \boxed{24}) \quad (3.4.22)$$

which is the same as the permutation  $\sigma_0 = (12)$  (as long as we identify  $\boxed{13}$  with 1 and  $\boxed{24}$  with 2). Insisting  $\alpha$  has the form in Remark 3.4.16 allows us to label blocks by reducing modulo  $d$  as in (3.4.22). The last requirement for a triple  $\tilde{\sigma}$  in Equaiton 3.4.20 to be in  $\text{Lifts}(\sigma, \tilde{G}_1)$  is that  $\tilde{\sigma}$  generates  $\tilde{G}_1$ . We obtain  $\text{Lifts}(\sigma, \tilde{G}_1)$  to be

$$\left\{ ((12)(34), (13)(24), (14)(23)), ((14)(23), (13)(24), (12)(34)) \right\} \quad (3.4.23)$$

- For  $\tilde{G}_2$ , we obtain  $\text{Lifts}(\sigma, \tilde{G}_2)$  to be

$$\begin{aligned} & \left\{ ((1432), \text{id}, (1234)), ((1234), (13)(24), (1234)), \right. \\ & \left. ((1234), \text{id}, (1432)), ((1432), (13)(24), (1432)) \right\} \end{aligned} \quad (3.4.24)$$

At the end of Step 4 we have that  $\text{Lifts}(\sigma)$  contains the 2 triples in (3.4.23) and the 4 triples in (3.4.24). Lastly, in Step 5 we quotient by simultaneous conjugation to obtain the 3 triples

$$\begin{aligned} \text{Lifts}(\sigma)/\sim = & \left\{ ((12)(34), (13)(24), (14)(23)), \right. \\ & ((1432), \text{id}, (1234)), \\ & \left. ((1234), (13)(24), (1234)) \right\} \end{aligned} \quad (3.4.25)$$

as output.

### 3.4 AN ITERATIVE ALGORITHM TO PRODUCE GENERATING TRIPLES

---

Now that we have an algorithm to find all lifts of a single permutation triple, we now describe how to use this to compute all isomorphism classes of 2-group permutation triples up to a given degree. In the algorithms to follow, we are concerned with constructing the bipartite graphs  $\mathcal{G}_{2^i}$  defined in Notation 3.4.10.

**Algorithm 3.4.26.** Let  $p = 2$  and the notation be as in 3.4.7 and 3.4.10. Then we can construct  $\mathcal{G}_2$  as follows.

- The set of nodes  $\mathcal{G}_2^{\text{below}}$  consists of a single triple  $(\text{id}, \text{id}, \text{id}) \in S_1^3$
- The set of nodes  $\mathcal{G}_2^{\text{above}}$  consists of 3 triples described in Example 3.4.5.
- The edge set of  $\mathcal{G}_2$  consists of 3 edges (i.e. it is the complete bipartite graph for the sets  $\mathcal{G}_2^{\text{below}}$  and  $\mathcal{G}_2^{\text{above}}$  )

*Proof of correctness.* By definition, the 3 degree 2 permutation triples from Example 3.4.5 are the only 2-group permutation triples of degree 2. These are all lifts of the unique 2-group permutation triple (in Example 3.4.4) via the extension

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\text{id}} \mathbb{Z}/2\mathbb{Z} \xrightarrow{\pi} \{\text{id}\} \longrightarrow 1 \quad (3.4.27)$$

□

Having constructed  $\mathcal{G}_2$ , we now describe the iterative process to compute  $\mathcal{G}_{2^i}$  from  $\mathcal{G}_{2^{i-1}}$ .

**Algorithm 3.4.28.** Let  $p = 2$  and the notation be as in 3.4.7 and 3.4.10. This algorithm describes the process of computing  $\mathcal{G}_{2^i}$  given  $\mathcal{G}_{2^{i-1}}$ .

**Input:** The bipartite graph  $\mathcal{G}_{2^{i-1}}$

**Output:** The bipartite graph  $\mathcal{G}_{2^i}$

### 3.4 AN ITERATIVE ALGORITHM TO PRODUCE GENERATING TRIPLES

---

1. For every  $\sigma \in \mathcal{G}_{2^{i-1}}^{\text{above}}$  apply Algorithm 3.4.11 to obtain the set  $\text{Lifts}(\sigma)/\sim$  for each  $\sigma$ . Combine these lifts into a single set

$$\text{Lifts}(\mathcal{G}_{2^{i-1}}) := \bigcup_{\sigma \in \mathcal{G}_{2^{i-1}}^{\text{above}}} \text{Lifts}(\sigma) \quad (3.4.29)$$

2. Compute  $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$  which we define to be the equivalence classes of  $\text{Lifts}(\mathcal{G}_{2^{i-1}})$  where two triples  $\tilde{\sigma}$  and  $\tilde{\sigma}'$  in  $\text{Lifts}(\mathcal{G}_{2^{i-1}})$  are equivalent if and only if they are simultaneously conjugate in  $S_{2^i}$ . Denote the equivalence class of  $\tilde{\sigma} \in \text{Lifts}(\mathcal{G}_{2^{i-1}})$  by  $[\tilde{\sigma}] \in \text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$ .
3. Define  $\mathcal{G}_{2^i}^{\text{below}} := \mathcal{G}_{2^{i-1}}^{\text{above}}$ . Define  $\mathcal{G}_{2^i}^{\text{above}}$  by choosing a single representative for each equivalence class of  $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$ . This defines the nodes of  $\mathcal{G}_{2^i}$ .
4. For every pair  $(\tilde{\sigma}, \sigma) \in \mathcal{G}_{2^i}^{\text{above}} \times \mathcal{G}_{2^i}^{\text{below}}$  place an edge between  $\tilde{\sigma}$  and  $\sigma$  if and only if there is a triple in the equivalence class  $[\tilde{\sigma}] \in \text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$  that is a lift of  $\sigma$ .
5. Return  $\mathcal{G}_{2^i}$  as output.

*Proof of correctness.* Since 2-groups are nilpotent, every 2-group permutation triple of degree  $2^i$  is the lift of at least one 2-group permutation triple of degree  $2^{i-1}$ . Let  $\tilde{\sigma} \in \mathcal{G}_{2^i}^{\text{above}}$  be an arbitrary representative of an isomorphism class of 2-group permutation triples of degree  $2^i$  contained in  $\text{Lifts}(\sigma)$  for some degree  $2^{i-1}$  triple  $\sigma$ . Let  $\sigma'$  denote the representative in  $\mathcal{G}_{2^{i-1}}^{\text{above}}$  that is simultaneously conjugate to  $\sigma$ . Algorithm 3.4.11 ensures that there is a 2-group permutation triple  $\tilde{\sigma}'$  of degree  $2^i$  in  $\text{Lifts}(\sigma')$  that is simultaneously conjugate to  $\tilde{\sigma}$ . Thus,  $\text{Lifts}(\mathcal{G}_{2^{i-1}})$  computed in Step 1 contains at least one triple for every isomorphism class of 2-group permutation triples of degree  $2^i$ . It is,

### 3.5 RESULTS OF COMPUTATIONS

---

however, possible for  $\text{Lifts}(\mathcal{G}_{2^{i-1}})$  to contain simultaneously conjugate triples arising as lifts of different triples in  $\mathcal{G}_{2^{i-1}}^{\text{above}}$ . Step 2 quotients  $\text{Lifts}(\mathcal{G}_{2^{i-1}})$  by simultaneous conjugation and Steps 3 and 4 define the desired graph  $\mathcal{G}_{2^i}$  in such a way that the edge structure of the lifts is preserved.  $\square$

Algorithm 3.4.26 combined with Algorithm 3.4.28 allows us to compute

$$\mathcal{G}_2, \mathcal{G}_4, \dots, \mathcal{G}_{2^i}, \dots, \mathcal{G}_{2^m} \quad (3.4.30)$$

up to any degree  $d = 2^m$ . A Magma implementation of Algorithms 3.4.11, 3.4.26, and 3.4.28 can be found at [24]. In the next section we discuss the results of these computations.

#### Section 3.5

### Results of computations

In this section we discuss the Magma implementation of Algorithms 3.4.11, 3.4.26, and 3.4.28 available at [24] where the techniques of this chapter are used to tabulate a database of 2-group permutation triples up to degree 256. This computation took roughly 50 CPU hours on a standart desktop. The majority of this time is spent checking conjugacy of degree 256 permutation triples. This database consists of roughly 340MB worth of text files. We devote the rest of this section to summarizing the results of these computations.

**Theorem 3.5.1.** *The following table lists the number of isomorphism classes of 2-*

### 3.5 RESULTS OF COMPUTATIONS

---

*group permutation triples of degree  $d$  up to 256.*

$d$	1	2	4	8	16	32	64	128	256
$\#$ permutation triples	1	3	7	19	55	151	503	1799	7175

(3.5.2)

**Theorem 3.5.3.** *The following table lists the number of passports of 2-group permutation triples of degree  $d$  up to 256.*

$d$	1	2	4	8	16	32	64	128	256
$\#$ passports	1	3	7	16	41	96	267	834	2893

(3.5.4)

**Theorem 3.5.5.** *The following table lists the number of lax passports of 2-group permutation triples of degree  $d$  up to 256.*

$d$	1	2	4	8	16	32	64	128	256
$\#$ lax passports	1	1	3	6	14	31	85	257	882

(3.5.6)

**Theorem 3.5.7.** *The following table lists the number of 2-group permutation triples up to degree 256 with  $\{\text{order}(\sigma_s) : s \in \{0, 1, \infty\}\}$  equal to  $\{a, b, c\}$  as sets.*

$(a, b, c)$	$\#$ permutation triples
(1, 1, 1)	1
(1, 2, 2)	3
(1, 4, 4)	3
(1, 8, 8)	3
(1, 16, 16)	3
(1, 32, 32)	3



### 3.5 RESULTS OF COMPUTATIONS

---

$(1, 64, 64)$	3
$(1, 128, 128)$	3
$(1, 256, 256)$	3
$(2, 2, 2)$	1
$(2, 2, 4)$	24
$(2, 2, 8)$	132
$(2, 2, 16)$	144
$(2, 2, 32)$	60
$(2, 2, 64)$	24
$(2, 2, 128)$	12
$(2, 4, 4)$	24
$(2, 4, 8)$	78
$(2, 4, 16)$	78
$(2, 4, 32)$	30
$(2, 4, 64)$	18
$(2, 4, 128)$	6
$(2, 8, 8)$	132
$(2, 8, 16)$	156
$(2, 8, 32)$	60
$(2, 8, 64)$	12
$(2, 16, 16)$	144
$(2, 16, 32)$	36
$(2, 32, 32)$	60

### 3.5 RESULTS OF COMPUTATIONS

---

(2, 64, 64)	24
(2, 128, 128)	12
(2, 256, 256)	3
(4, 4, 4)	65
(4, 4, 8)	1581
(4, 4, 16)	969
(4, 4, 32)	225
(4, 4, 64)	69
(4, 4, 128)	15
(4, 8, 8)	1581
(4, 8, 16)	960
(4, 8, 32)	168
(4, 8, 64)	24
(4, 16, 16)	969
(4, 16, 32)	84
(4, 32, 32)	225
(4, 64, 64)	69
(4, 128, 128)	15
(4, 256, 256)	6
(8, 8, 8)	726
(8, 8, 16)	1542
(8, 8, 32)	378
(8, 8, 64)	78

### 3.5 RESULTS OF COMPUTATIONS

---

(8, 16, 16)	1542
(8, 16, 32)	72
(8, 32, 32)	378
(8, 64, 64)	78
(8, 128, 128)	24
(8, 256, 256)	12
(16, 16, 16)	136
(16, 16, 32)	552
(16, 32, 32)	552
(16, 64, 64)	144
(16, 128, 128)	48
(16, 256, 256)	24
(32, 64, 64)	288
(32, 128, 128)	96
(32, 256, 256)	48
(64, 128, 128)	192
(64, 256, 256)	96
(128, 256, 256)	192

### 3.5 RESULTS OF COMPUTATIONS

---




Figure 3.5.7: Distribution of genera up to degree 256

### 3.5 RESULTS OF COMPUTATIONS

---




Figure 3.5.7: # nonhyperbolic and hyperbolic passports by degree (left), and # nonhyperbolic and hyperbolic lax passports by degree (right).

### 3.5 RESULTS OF COMPUTATIONS




Figure 3.5.7: # permutation triples by degree with abelian and nonabelian monodromy groups (left) and # permutation triples by degree with monodromy groups of various nilpotency classes (right).

### 3.5 RESULTS OF COMPUTATIONS

---




Figure 3.5.7: # passports of various sizes by degree

---

## Chapter 4

---

# Fields of definition of 2-group Belyi maps

Recall from Definition 2.1.8, that a field of definition of a Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  is a number field  $K \subseteq \mathbb{C}$  such that  $X$  and  $\phi$  are defined using algebraic equations having coefficients in  $K$ . Using data from Chapter 3, we formulate a conjecture about the possible fields of definition of 2-group Belyi maps.

### Section 4.1

#### Refined passports

Let  $\sigma$  be a 2-group permutation triple. Recall, from Definition 2.2.4, that the passport of  $\sigma$  consists of the data  $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$  where  $g(\sigma)$  is the genus,  $\langle \sigma \rangle$  is the monodromy group (2-group in its regular representation) as a subgroup of  $S_d$ , and  $\lambda(\sigma)$  is the triple of partitions specifying the three ordered  $S_d$  conjugacy classes  $C_0, C_1, C_\infty$  of  $\sigma_0, \sigma_1, \sigma_\infty$  respectively. Let  $\mathcal{P}$  be the passport of  $\sigma$ . The size of  $\mathcal{P}$  is the cardinality



of the set

$$\Sigma_{\mathcal{P}} = \{(\sigma_0, \sigma_1, \sigma_\infty) \in C_0 \times C_1 \times C_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1, \sigma_\infty \rangle = G\} / \sim \quad (4.1.1)$$

where  $(\sigma_0, \sigma_1, \sigma_\infty) \sim (\sigma'_0, \sigma'_1, \sigma'_\infty)$  if the triples are simultaneously conjugate by an element of  $S_d$ . By Theorem 2.4.5, the cardinality of  $\Sigma_{\mathcal{P}}$  bounds the field of moduli of the Belyi map corresponding to  $\sigma$ .

Let  $G$  be a transitive subgroup of  $S_d$  and let  $C$  be a conjugacy class of  $S_d$ . Then  $C$  can be partitioned into conjugacy classes of  $G$ . To analyze conjugacy in  $G$  we make the following definition.

**Definition 4.1.2.** A refined passport  $\mathcal{P}$  consists of the data  $(g, G, c)$  where  $g \geq 0$  is an integer,  $G \leq S_d$  is a transitive subgroup, and  $c = (c_0, c_1, c_\infty)$  is a triple of conjugacy classes of  $G$ . For a refined passport  $\mathcal{P}$  consider the set

$$\Sigma_{\mathcal{P}} = \{(\sigma_0, \sigma_1, \sigma_\infty) \in c_0 \times c_1 \times c_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1, \sigma_\infty \rangle = G\} / \sim \quad (4.1.3)$$

where  $(\sigma_0, \sigma_1, \sigma_\infty) \sim (\sigma'_0, \sigma'_1, \sigma'_\infty)$  if and only if there exists  $\alpha \in \text{Aut}(G)$  with  $\alpha(\sigma_s) = \sigma'_s$  for  $s \in \{0, 1, \infty\}$ .

Let  $\sigma$  be a 2-group permutation triple and let  $c_s$  denote the conjugacy class of  $\langle \sigma \rangle$  containing  $\sigma_s$  for  $s \in \{0, 1, \infty\}$ . We define the refined passport of  $\sigma$  to be

$$\mathcal{P}(\sigma) = (g(\sigma), \langle \sigma \rangle, (c_0, c_1, c_\infty)). \quad (4.1.4)$$

Section 4.2

## A refined conjecture

Let  $\sigma$  be a 2-group permutation triple. Let  $\mathcal{P}$  and  $\mathcal{P}$  denote the passport and refined passport of  $\sigma$  respectively. Let  $\Sigma_{\mathcal{P}}$  and  $\Sigma_{\mathcal{P}}$  denote the sets in (4.1.1) and (4.1.3) respectively. Let  $\tilde{\sigma}$  be a lift of  $\sigma$  with passport and refined passport  $\tilde{\mathcal{P}}$  and  $\tilde{\mathcal{P}}$ .

Chapter 3 provides us with an explicit list of all 2-group permutation triples (up to simultaneous conjugation in  $S_d$ ) for fixed degree. Using explicit techniques from Musty, Schiavone, Sijlsing, and Voight in [25], we computed  $\Sigma_{\mathcal{P}(\sigma)}$  for every 2-group permutation triple  $\sigma$  of degree  $d$  for  $d \leq 256$ .

*We observed that  $\#\Sigma_{\mathcal{P}(\sigma)} = 1$  for every such permutation triple  $\sigma$ !* (4.2.1)

This observation provides motivation to take a closer look at the behavior of refined passports with respect to the iterative structure of 2-group permutation triples.

**Lemma 4.2.2.** *Let  $\sigma$  be a 2-group permutation triple with passport  $\mathcal{P} = (g, G, C)$  where  $C = (C_0, C_1, C_\infty)$ . Let  $\mathcal{P} = (g, G, c)$  be the refined passport of  $\sigma$  where  $c = (c_0, c_1, c_\infty)$ . Let  $\sigma'$  be a 2-group permutation triple simultaneously conjugate to  $\sigma$  with refined passport  $\mathcal{P}' = (g, G, c')$  where  $c' = (c'_0, c'_1, c'_\infty)$ . Then  $\#\Sigma_{\mathcal{P}} = \#\Sigma_{\mathcal{P}'}$ .*

*Proof.* Let  $S_d$  denote the ambient symmetric group. By hypothesis, there exists a permutation  $\tau \in S_d$  such that  $\sigma^\tau = \sigma'$ . Now define a map  $\tau: \Sigma_{\mathcal{P}} \rightarrow \Sigma_{\mathcal{P}'}$  by simultaneously conjugating a representative triple in  $\Sigma_{\mathcal{P}}$  by  $\tau$ . Since conjugation by  $\tau$  in  $S_d$  is an automorphism of  $G$ , simultaneous conjugation by  $\tau$  is a bijection from  $\Sigma_{\mathcal{P}}$  to  $\Sigma_{\mathcal{P}'}$ . □

**Definition 4.2.3.** Let  $\tilde{\sigma}$  be a lift of a 2-group permutation triple  $\sigma$  obtained from the exact sequence

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} G \longrightarrow 1 \quad (4.2.4)$$

with  $G = \langle \sigma \rangle$  and  $\tilde{G} = \langle \tilde{\sigma} \rangle$ . We say that  $\tilde{\sigma}$  is a **characteristic lift** of  $\sigma$  if  $\iota(\mathbb{Z}/2\mathbb{Z})$  is a characteristic subgroup of  $\tilde{G}$ .

**Lemma 4.2.5.** *Let  $\sigma$  and  $\sigma'$  be 2-group permutation triples with distinct refined passports  $(g, G, c)$  and  $(g, G, c')$  respectively. Then the refined passport of any characteristic lift of  $\sigma$  is not equal to the refined passport of any characteristic lift of  $\sigma'$ .*

*Proof.* Assume for contradiction that we have lifts  $\tilde{\sigma}$  and  $\tilde{\sigma}'$  of  $\sigma$  and  $\sigma'$  with the same refined passport  $(\tilde{g}, \tilde{G}, \tilde{C})$ . Let

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} G \longrightarrow 1 \quad (4.2.6)$$

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota'} \tilde{G} \xrightarrow{\pi'} G \longrightarrow 1$$

be extensions of  $\mathbb{Z}/2\mathbb{Z}$  by  $G$  such that  $\tilde{\sigma} \in \pi^{-1}(\sigma)$  and  $\tilde{\sigma}' \in \pi'^{-1}(\sigma')$ . Since  $\tilde{\sigma}$  and  $\tilde{\sigma}'$  have the same refined passport, there exists an automorphism  $\tilde{\alpha} \in \text{Aut}(\tilde{G})$  with  $\tilde{\alpha}(\tilde{\sigma}_s) = \tilde{\sigma}'_s$  for each  $s \in \{0, 1, \infty\}$ .

Define  $\alpha: G \rightarrow G$  by projecting  $\tilde{\alpha}$ , that is,  $\alpha := \pi' \circ \tilde{\alpha} \circ \pi^{-1}$ . Since  $\alpha(\sigma_s) = \sigma'_s$  by definition, we will have a contradiction if we can show that  $\alpha \in \text{Aut}(G)$ . Let  $\tau$  be the generator of the characteristic subgroup  $\iota(\mathbb{Z}/2\mathbb{Z}) = \iota'(\mathbb{Z}/2\mathbb{Z})$  and note that  $\tilde{\alpha}(\tau) = \tau$ .

First we show that  $\alpha$  is well-defined. For  $\sigma_s \in \sigma$ ,  $\pi^{-1}(\sigma_s) \in \{\tilde{\sigma}_s, \tau\tilde{\sigma}_s\}$ , and the

calculation

$$\pi'(\tilde{\alpha}(\tau\tilde{\sigma}_s)) = \pi'(\tau\tilde{\alpha}(\tilde{\sigma}_s)) = \pi'(\tau\tilde{\sigma}'_s) \quad (4.2.7)$$

shows that  $\alpha$  is well-defined homomorphism.

To finish the proof we now show that  $\alpha$  is injective. Let  $g \in \ker \alpha$  so that  $\pi'(\tilde{\alpha}(\pi^{-1}(g))) = \text{id}_G$ . Then  $\tilde{\alpha}(\pi^{-1}(g)) \in \iota'(\mathbb{Z}/2\mathbb{Z})$  which implies  $\pi^{-1}(g) \in \iota(\mathbb{Z}/2\mathbb{Z})$  so that  $g = \text{id}_G$ .  $\square$

**Conjecture 4.2.8.** *Every 2-group permutation triple  $\sigma$  has refined passport size 1.*

**Lemma 4.2.9.** *Conjecture 4.2.8 is true for  $\langle \sigma \rangle$  abelian.*

*Proof.* In an abelian group the conjugacy classes consist of sets of size 1. Therefore every refined passport  $\mathcal{P} = (g, G, c)$  with  $G$  abelian has

$$\Sigma_{\mathcal{P}} \leq 1. \quad (4.2.10)$$

Since  $\sigma$  is a 2-group permutation triple with refined passport  $\mathcal{P}(\sigma)$ , we have that

$$\Sigma_{\mathcal{P}(\sigma)} \geq 1. \quad (4.2.11)$$

(4.2.10) and (4.2.11) imply  $\Sigma_{\mathcal{P}(\sigma)} = 1$  for  $\langle \sigma \rangle$  abelian.  $\square$

**Theorem 4.2.12.** *Conjecture 4.2.8 is true for  $\langle \sigma \rangle$  dihedral.*

*Proof.* The proof is by induction on the degree of  $\sigma$ . There is a unique 2-group permutation triple of degree 1 which therefore has refined passport size 1. For induction, assume that every 2-group permutation triple of degree  $d$  with  $\langle \sigma \rangle$  dihedral has refined passport size 1. Let  $\tilde{\sigma}$  be a 2-group permutation triple of degree  $2d$  with  $\langle \tilde{\sigma} \rangle$  dihedral. We are required to show that the refined passport of  $\tilde{\sigma}$  has size 1.

## 4.2 A REFINED CONJECTURE

---

Let  $\Sigma_{\mathcal{P}} := \Sigma_{\mathcal{P}(\tilde{\sigma})}$  be a set of refined passport representatives defined in (4.1.3). By Algorithm 3.4.11 there exists a 2-group permutation triple  $\sigma$  such that the following sequence is exact.

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 \quad (4.2.13)$$

In other words,  $\tilde{\sigma}$  is a lift of  $\sigma$ . By Lemma 3.2.11, quotients of dihedral groups are dihedral, and therefore  $\langle \sigma \rangle$  is dihedral. Thus, by induction, the refined passport of  $\sigma$  has size 1.

First note that for  $\langle \tilde{\sigma} \rangle$  dihedral we have  $Z(\langle \tilde{\sigma} \rangle) = \mathbb{Z}/2\mathbb{Z}$ . Now, since  $\iota(\mathbb{Z}/2\mathbb{Z})$  is contained in the center of  $\langle \tilde{\sigma} \rangle$ , we have that  $\iota(\mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}$ . Thus, we can apply Lemma 4.2.5 to get that every element of  $\Sigma_{\mathcal{P}}$  must also be a lift of  $\sigma$ . Thus, it is sufficient to prove that every lift  $\tilde{\sigma}'$  satisfies one of the following.

1. The  $\langle \tilde{\sigma} \rangle$  conjugacy class of  $\tilde{\sigma}'_s$  differs from the  $\langle \tilde{\sigma} \rangle$  conjugacy class of  $\tilde{\sigma}_s$  for some  $s \in \{0, 1, \infty\}$
2. There exists an automorphism  $\alpha \in \text{Aut}(\langle \tilde{\sigma} \rangle)$  with  $\alpha(\tilde{\sigma}'_s) = \tilde{\sigma}_s$  for all  $s \in \{0, 1, \infty\}$

Let  $\tau \in \iota(\mathbb{Z}/2\mathbb{Z})$  be the generator of the image. There are  $2^3 = 8$  preimages of  $\sigma$  under the map  $\pi$ . Since  $\tilde{\sigma}_{\infty}\tilde{\sigma}_1\tilde{\sigma}_0 = 1$  and  $\tau$  is central, there are exactly 4 preimages that multiply to 1. They are as follows.

$$\{(\tilde{\sigma}_0, \tilde{\sigma}_1, \tilde{\sigma}_{\infty}), (\tilde{\sigma}_0, \tau\tilde{\sigma}_1, \tau\tilde{\sigma}_{\infty}), (\tau\tilde{\sigma}_0, \tilde{\sigma}_1, \tau\tilde{\sigma}_{\infty}), (\tau\tilde{\sigma}_0, \alpha\tilde{\sigma}_1, \tilde{\sigma}_{\infty})\} \quad (4.2.14)$$

Using the notation for dihedral groups from Example 3.2.1 with  $2^{n+1}$  replaced with

## 4.2 A REFINED CONJECTURE

---

$2d$  we have the following conjugacy classes of  $\langle \tilde{\sigma} \rangle$ .

- $\{1\}, \{a^{\frac{d}{2}}\}$
- $c_i := \{a^i, a^{-i}\}$  for  $i \in \{1, \dots, \frac{d}{2} - 1\}$
- $c_{\text{even}} := \{a^{2i}b : 0 \leq i \leq \frac{d}{2} - 1\}$
- $c_{\text{odd}} := \{a^{2i+1}b : 0 \leq i \leq \frac{d}{2} - 1\}$

Note that  $c_{\text{even}}$  and  $c_{\text{odd}}$  consist of all the involutions of the group. We will say that an involution has **even parity** if it is in  $c_{\text{even}}$  and **odd parity** if it is in  $c_{\text{odd}}$ .

If  $\tilde{\sigma}_s$  is central, then the conjugacy class of  $\tau\tilde{\sigma}_s$  cannot be equal to the conjugacy class of  $\tilde{\sigma}_s$ . Thus we can assume the conjugacy class of  $\tilde{\sigma}_s$  is  $c_i$  for some  $i$ ,  $c_{\text{even}}$ , or  $c_{\text{odd}}$  for every  $s \in \{0, 1, \infty\}$ . If  $\tilde{\sigma}_s \in c_i$ , then

$$\tau\tilde{\sigma}_s = a^{d/2}a^{\pm i} = a^{\frac{d}{2} \pm i}. \quad (4.2.15)$$

Now  $a^{\frac{d}{2} \pm i} \in c_i$  if and only if  $i = \pm d/4$ . Thus we can assume the conjugacy class of  $\tilde{\sigma}_s$  is  $c_{d/4}$ ,  $c_{\text{even}}$ , or  $c_{\text{odd}}$  for every  $s \in \{0, 1, \infty\}$ .

Now let us focus on the condition that  $\tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1$ . First note that every element of  $c_{\text{even}} \cup c_{\text{odd}}$  is an involution, so we need at least one of  $\tilde{\sigma}_s \in c_{d/4}$  to satisfy  $\tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1$ . Without loss of generality assume that  $\tilde{\sigma}_\infty \in c_{d/4}$ . Then since  $\tilde{\sigma}_\infty$  has order 4, we must have  $\tilde{\sigma}_0, \tilde{\sigma}_1 \in c_{\text{even}} \cup c_{\text{odd}}$  (again to have a chance of satisfying their product equals 1). Let  $\tilde{\sigma}_1 = a^k b \in c_{\text{even}} \cup c_{\text{odd}}$  be an involution. Then

$$\tilde{\sigma}_\infty \tilde{\sigma}_1 = a^{\pm d/4} a^k b = a^{\pm(d/4) + k} b \quad (4.2.16)$$

## 4.2 A REFINED CONJECTURE

---

The parity of the involution  $\tilde{\sigma}_1$  is the same as the parity of the involution  $\tilde{\sigma}_\infty \tilde{\sigma}_1$  when  $d \geq 8$  and the parity is different for  $d = 4$ .

*Claim.* Two involutions of the same parity do not generate  $D_{2d}$  for  $d = 2^n$  and  $n \geq 2$ .

*Proof of Claim.* Let  $a^k b$  and  $a^{k'} b$  be involutions of  $D_{2d}$  with  $k$  and  $k'$  the same parity. Since  $d \geq 4$  is a power of 2,  $\pm k, \pm k'$  all have the same parity. Now since  $\langle a^k b, a^{k'} b \rangle$  is generated by involutions we have

$$\begin{aligned} \langle a^k b, a^{k'} b \rangle &= \{a^k b, a^{k'} b\} \cup \langle a^k b a^{k'} b \rangle \\ &= \{a^k b, a^{k'} b\} \cup \langle a^{k-k'} \rangle \\ &= \{a^k b, a^{k'} b\} \cup \langle a^{k'-k} \rangle \end{aligned} \tag{4.2.17}$$

which never contains  $a$  since  $k$  and  $k'$  have the same parity.  $\square$

The claim finishes the proof for  $d \geq 8$ . To summarize, the condition that  $\tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1$  with each  $\tilde{\sigma}_s \in c_{d/4} \cup c_{\text{even}} \cup c_{\text{odd}}$  implies that the triple  $\tilde{\sigma}$  consists of exactly one element from  $c_{d/4}$  and the other two elements are involutions with the same parity. By the claim, no such triple can generate  $D_{2d}$  which completes the proof for  $d \geq 8$ .

It remains to consider the case  $d = 4$  (i.e.  $D_{2d} = D_8$ ) when the 2 involutions have opposite parity. At this point we could simply appeal to the explicit computation of this refined passport in observation (4.2.1), but we also provide a proof in this specific case for completeness.

In this case,  $c_{d/4} = c_1 = \{a, a^{-1}\}$ ,  $c_{\text{even}} = \{b, a^2 b\}$ ,  $c_{\text{odd}} = \{ab, a^3 b\}$ , and we are considering triples  $\tilde{\sigma} \in c^3$  where  $c = c_1 \cup c_{\text{even}} \cup c_{\text{odd}}$ . As discussed above, to satisfy  $\tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1$  we must have exactly one of  $\tilde{\sigma}_s \in c_1$ . Without loss of generality assume  $\tilde{\sigma}_\infty \in c_1$ . Then from (4.2.16) we have that  $\tilde{\sigma}_1$  and  $\tilde{\sigma}_0$  are involutions with opposite

## 4.2 A REFINED CONJECTURE

---

parity. Without loss of generality let  $\tilde{\sigma}_1 \in c_{\text{odd}}$  and  $\tilde{\sigma}_0 \in c_{\text{even}}$ . We then have the following triples  $\tilde{\sigma}$  that generate  $D_8$  and satisfy  $\tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1$ .

$$(a^2b, ab, a), (b, a^3b, a), (b, ab, a^{-1}), (a^2b, a^3b, a^{-1}) \quad (4.2.18)$$

To show that the refined passport of a  $\tilde{\sigma}$  taken from (4.2.18) has size 1, we are required to find elements of  $\text{Aut}(D_8)$  showing all 4 of these triples are equivalent. Let  $f_1 \in \text{Aut}(D_8)$  be defined by  $a \mapsto a$ , and  $b \mapsto a^2b$ . Let  $f_2 \in \text{Aut}(D_8)$  be defined by  $a \mapsto a^{-1}$ , and  $b \mapsto b$ . Then  $f_1$  identifies  $(a^2b, ab, a)$  and  $(b, a^3b, a)$ ,  $f_2$  identifies  $(b, a^3b, a)$  and  $(b, ab, a^{-1})$ , and  $f_1$  identifies  $(b, ab, a^{-1})$  and  $(a^2b, a^3b, a^{-1})$ . This completes the proof for  $D_8$  and thus for the entire dihedral case.  $\square$

*Remark 4.2.19.* It is reasonable to expect that a similar direct approach in the proof of Theorem 4.2.12 could succeed for other families of 2-groups that satisfy Theorem 3.2.11, namely generalized quaternion groups and semi-dihedral groups. These families also have characteristic centers so that Lemma 4.2.5 can be applied. However, a different technique is required for a general proof, since experimental evidence in Section 3.5 suggests that 2-groups of all nilpotency classes appear as monodromy groups of 2-group permutation triples.

Conjecture 4.2.8, together with refined field of definition equal to field of moduli for Galois Belyi maps and a strong version of Beckmann's theorem would imply the following.

**Conjecture 4.2.20.** *Every 2-group Belyi map is defined over an abelian extension of  $\mathbb{Q}$ .*



---

## Chapter 5

---

# Computing equations

In this chapter we discuss how to compute equations for 2-group Belyi maps corresponding to the 2-group permutation triples computed in Chapter 3. As was the case for computing the permutation triples, the algorithm to compute equations follows an iterative approach. In this chapter we construct the 2-group Belyi maps as towers of quadratic extensions of function fields. We begin in Section 5.1 by discussing the analogous situation over number fields. In Section 5.2 and Section 5.3 we discuss the relevant background about algebraic function fields. The algorithms to compute equations for 2-group Belyi maps (over  $\mathbb{F}_q$ ) are described in Section 5.4, the implementation in characteristic zero is detailed in Section 5.5, and the results of these computations can be found in Section 5.6.

### Section 5.1

## Quadratic extensions of number fields

By way of motivation, let  $F$  be a number field and let  $\mathbb{Z}_F$  denote the ring of integers of  $F$ . Kummer theory tells us that quadratic extensions of  $F$  are in bijection with

nontrivial cosets  $dF^{\times 2}$  in the quotient  $F^{\times}/F^{\times 2}$ ; such a coset defines a quadratic extension  $F(\sqrt{d})$ . Conversely, let  $F(\alpha)$  be a quadratic extension of  $F$ . The discriminant of the minimal polynomial of  $\alpha$  defines the bijection in the other direction.

Let  $\text{Pl}(F)$  denote the set of places of  $F$  and let  $S_{\infty}$  denote the archimedean places. For  $v \in \text{Pl}(F) \setminus S_{\infty}$  let  $\mathfrak{p}_v$  be the prime ideal of  $\mathbb{Z}_F$  corresponding to  $v$ . Let  $S \subset \text{Pl}(F) \setminus S_{\infty}$  be a finite set of nonarchimedean places, and further suppose that each place of  $S$  has odd residue field. We aim to answer the following question.

*Question 5.1.1.* How do we construct a quadratic extension of  $F$  ramified at  $\mathfrak{p}_v$  for all  $v \in S$  and unramified at all nonarchimedean places outside of  $S$ ? If so, then how *unique* is the construction?

To formulate this question more clearly, let  $\mathfrak{a} := \prod_{v \in S} \mathfrak{p}_v$  encode the primes we want to ramify in this quadratic extension. There are three possibilities.

- It is possible that no such extension exists.
- $\mathfrak{a} = (d)$  is principal and the extension  $F(\sqrt{d})$  is a quadratic extension ramified exactly at each  $\mathfrak{p}_v$ , and the generator  $d$  is unique up to multiplication by a unit in  $\mathbb{Z}_F^{\times}$ .
- If  $\mathfrak{a}$  is not principal, it is possible there exists a fractional ideal  $\mathfrak{b}$  such that  $\mathfrak{a}\mathfrak{b}^2 = (d)$ . In this case, we can again construct the extension  $F(\sqrt{d})$  with the prescribed ramification, but  $d$  is only unique up to the ideal  $\mathfrak{b}$  used to construct it.

Let us consider the last case more closely. Let  $\text{Cl}_F$  denote the class group of  $F$  and for a fractional ideal  $\mathfrak{c}$  let  $[\mathfrak{c}]$  denote its ideal class in  $\text{Cl}_F$ . The equation  $\mathfrak{a}\mathfrak{b}^2 = (d)$

means that  $[\mathfrak{a}] = [\mathfrak{b}^{-2}]$  so that  $[\mathfrak{a}] \in \text{Cl}_F^2$ . Moreover, if we take an element  $[\mathfrak{c}]$  of order 2 in  $\text{Cl}_F$ , then

$$[\mathfrak{a}\mathfrak{b}^2] = [\mathfrak{a}\mathfrak{b}^2] = [\mathfrak{a}\mathfrak{b}^2][\mathfrak{c}^2] = [\mathfrak{a}(\mathfrak{b}\mathfrak{c})^2]. \quad (5.1.2)$$

Thus, in the case where  $\mathfrak{a}$  is not principal, but there exists  $\mathfrak{b}$  with  $\mathfrak{a}\mathfrak{b}^2$  principal, we have  $[\mathfrak{a}] \in \text{Cl}_F^2$  and  $[\mathfrak{b}]$  is unique up to multiplication by  $[\mathfrak{c}] \in \text{Cl}_F[2]$ .

We can now formulate our precise goal. Given  $\mathfrak{a}$  (encoding ramification data), find  $\mathfrak{b}^2$  and  $d$  such that  $\mathfrak{a}\mathfrak{b}^2 = (d)$ . In the following sections we will rephrase this problem in the function field setting.

## Section 5.2

# Curves and algebraic function fields

In this section we summarize the setting in which the algorithms of this chapter are stated. There are many comprehensive resources on this topic such as [16, I.6], [35], and [28].

First, let  $K$  be a perfect field.

**Definition 5.2.1.** An algebraic function field in one variable over  $K$  is a field extension  $F$  over  $K$  of transcendence degree 1. That is, there exists  $x \in F$  such that  $x$  is transcendental over  $K$  and  $[F : K(x)]$  is finite.

**Definition 5.2.2.** We say the  $K$  is the constant field of  $F$ . The exact constant field of  $F$  is the algebraic closure of  $K$  in  $F$ .

*Remark 5.2.3.* In theory we can assume that  $K$  is the exact constant field of  $F$ , but in practice for  $K$  a number field or  $\mathbb{F}_q$  we try to work with constant fields that are as simple as possible.

*Example 5.2.4.* Let  $X$  be an irreducible affine plane curve defined by the defining equation  $f(x, y) = 0$  with  $f \in K[x, y]$ . Then the function field of  $X$ , denoted by  $K(X)$ , is defined to be the field of fractions of the coordinate ring  $\frac{K[x, y]}{(f(x, y))}$  of  $X$ .

**Definition 5.2.5.** A place  $P$  of  $F$  is the maximal ideal of some discrete valuation ring  $\mathcal{O}_P$  of  $F$ . We denote the valuation on  $F$  corresponding to  $P$  by  $\text{ord}_P$ . The set of places of  $F$  is denoted  $\text{Pl}(F)$ . The degree of a place  $P$ , denoted  $\deg(P)$ , is the index  $[\mathcal{O}_P/P : K]$  of the residue class field as an extension of  $K$ .

**Definition 5.2.6.** The divisor class group of  $F$ , denoted  $\text{Div}(F)$ , is the free abelian group generated by the places of  $F$ . A divisor  $D \in \text{Div}(F)$  is represented by a formal sum of places  $D = \sum_{P \in \text{Pl}(F)} a_P P$  with  $a_P \in \mathbb{Z}$  for all  $P$  and  $a_P = 0$  for all but finitely many  $P$ . We define  $\text{ord}_P(D)$  to be the coefficient  $a_P$  in the representation of  $D$ .

**Definition 5.2.7.** The support of a divisor  $D = \sum_{P \in \text{Pl}(F)} a_P P$ , denoted  $\text{supp}(D)$ , is  $\{P \in \text{Pl}(F) : a_P \neq 0\}$ . The degree of  $D$  is defined to be  $\deg(D) := \sum_{P \in \text{Pl}(F)} a_P \deg(P)$ . The subgroup of  $\text{Div}(F)$  consisting of the set of degree zero divisors of  $F$  is denoted by  $\text{Div}^0(F)$ .

**Definition 5.2.8.** The image of the map  $\text{div} : F^\times \rightarrow \text{Div}(F)$  defined by

$$\text{div}(f) = \sum_{P \in \text{Pl}(F)} \text{ord}_P(f) P \quad (5.2.9)$$

is the subgroup of principal divisors of  $F$  and denoted  $\text{Princ}(F)$ . Two divisors  $D_1, D_2 \in \text{Div}(F)$  are linearly equivalent if  $D_1 - D_2 \in \text{Princ}(F)$ .

**Definition 5.2.10.** The Picard group of  $F$  is defined by  $\text{Pic}(F) := \text{Div}(F) / \text{Princ}(F)$ . The Jacobian of  $F$  is defined by  $\text{Pic}^0(F) := \text{Div}^0(F) / \text{Princ}(F)$ .

**Definition 5.2.11.** There is a partial order on  $\text{Div}(F)$  defined by  $D_1 \geq D_2$  if and only if  $\text{ord}_P(D_1) \geq \text{ord}_P(D_2)$  for all  $P \in \text{Pl}(F)$ . We say that  $D \in \text{Div}(F)$  is **effective** if  $D \geq 0$ .

**Definition 5.2.12.** The Riemann-Roch space of a divisor  $D \in \text{Div}(F)$  is defined by  $\mathcal{L}(D) := \{f \in F : \text{div}(f) + D \geq 0\} \cup \{0\}$ .

Now that we have some of the basic definitions of algebraic function fields, we also need to introduce some terminology concerning extensions of algebraic function fields.

**Definition 5.2.13.** Let  $F, F'$  be algebraic function fields over constant fields  $K, K'$  respectively and suppose that  $F \subseteq F'$  and  $K \subseteq K'$ . When these conditions are satisfied we say that  $F'$  is an **algebraic function field extension** of  $F$ .

Every place  $P'$  of  $F'$  lies **above** a unique place  $P = F \cap P'$  of  $F$ . Every place  $P$  of  $F$  lies **below** finitely many places  $P'$  of  $F'$ . We denote a place  $P'$  above  $P$  by  $P'|P$ . When  $P'|P$  we can view  $\mathcal{O}_{P'}$  as a free  $\mathcal{O}_P$ -module of rank  $[F' : F]$  and  $\mathcal{O}_P = \mathcal{O}_{P'} \cap F$ .

We now summarize the fundamental identity from algebraic number theory in the function field setting. Let  $F'$  over  $K'$  be an extension of  $F$  over  $K$  and let  $P'$  be a place of  $F'$  above  $P \in \text{Pl}(F)$ . There is a unique positive integer denoted  $e(P'|P)$  such that  $\text{ord}_{P'}(f) = e(P'|P) \text{ord}_P(f)$  for all  $f \in F$ . The positive integer  $e(P'|P)$  is called the **ramification index** of  $P'|P$ . The **residue degree**, denoted  $f(P'|P)$  is defined to be the index  $[\mathcal{O}_{P'}/P' : \mathcal{O}_P/P]$  which makes sense after embedding  $\mathcal{O}_P/P$  into  $\mathcal{O}_{P'}/P'$ . The **fundamental identity** is then given by the equation

$$[F' : F] = \sum_{P'|P} e(P'|P) f(P'|P). \quad (5.2.14)$$

We now summarize some necessary facts about extending the field of constants of and algebraic function field.

**Definition 5.2.15.** An extension  $F'$  (with constants  $K'$ ) of  $F$  (with constants  $K$ ) is a constant field extension if  $F' = FK'$ .

Constant field extensions are one way in which the function field setting differs from the number field setting. When  $F' = FK'$  is a constant field extension of  $F$  over  $K$ , there are several observations to make. First, the relative degree over the rational function field does not change, that is,  $[F : K(x)] = [F' : K'(x)]$  for all  $x \in F \setminus K$ . Second, no places of  $F$  ramify in  $F'$ . Lastly, define the **conorm map** by

$$\text{con}_{F'|F}(P) := \sum_{P'|P} e(P'|P)P' \in \text{Div}(F'). \quad (5.2.16)$$

The conorm map extends to a homomorphism on divisor, principal divisors, and hence on divisor classes. Since constant field extensions are unramified, the conorm map induces an injection  $\text{Pic}(F) \hookrightarrow \text{Pic}(F')$ .

We conclude this section by proving a lemma we will need later in this chapter.

**Lemma 5.2.17.** *Let  $aF^{\times 2}$  be a nontrivial coset of  $F^{\times}/F^{\times 2}$  and consider the extension  $K := F(\sqrt{a})$ . Then a prime  $P$  of  $F$  is ramified in  $K$  if and only if  $\text{ord}_P(a)$  is odd.*

*Proof.* Since  $a$  is not a square in  $F$ , the extension is quadratic. Suppose  $\text{ord}_P(a)$  is odd and let  $\mathfrak{p}$  be a place above  $P$  in  $K$ . Then we have

$$2 \text{ord}_{\mathfrak{p}}(\sqrt{a}) = \text{ord}_{\mathfrak{p}}(a) = e(\mathfrak{p}/P) \text{ord}_P(a). \quad (5.2.18)$$

Since  $\text{ord}_P(a)$  is odd, (5.2.18) implies that 2 divides  $e(\mathfrak{p}/P)$  so that  $P$  is ramified in

$K$ . Moreover, this says that  $e(\mathfrak{p}/P) = 2$ .

For the converse we check ramification locally at the place  $P$ . Suppose  $e := \text{ord}_P(a)$  is even. Choose a uniformizer  $t$  at  $P$  and let  $b = a/t^{(e/2)}$ . Then  $F(\sqrt{a}) = F(\sqrt{b})$ , and  $\text{ord}_P(b) = 0$  implies that  $F(\sqrt{b}) = F(\sqrt{a})$  is unramified at  $P$ .

For a more general proof of this in arbitrary Kummer extensions see [28, Proposition 10.3].  $\square$

### Section 5.3

## Quadratic extensions of function fields

We now address two tasks concerning quadratic extensions of function fields that we need for the algorithms in Section 5.4.

The first task is the problem (analogous to the problem in Section 5.1) of finding a quadratic extension  $F(\sqrt{f})/F$  with ramification (in the relative extension) prescribed by  $R \in \text{Div}(F)$ . By Proposition 5.2.17, we can take all nonzero coefficients of  $R$  to have absolute value 1. As was the case for number fields, there are three possibilities.

First, it could be the case that no such extension exists in which case there is nothing to do. The other easy case occurs when  $R$  is a principal divisor so that  $R = \text{div}(f)$  for some  $f \in F^\times$ . In this case, the extension  $F(\sqrt{f})$  has the desired ramification determined by  $R$ .

The last case occurs when  $R$  is not principal, but there exists  $D \in \text{Div}(F)$  with  $R - 2D = \text{div}(f)$  for some  $f \in F$ . By Proposition 5.2.17, the extension  $F(\sqrt{f})/F$  will be ramified precisely at the places in the support of  $R$ . For  $D \in \text{Div}(F)$ , let  $[D]$  denote the class of  $D$  in  $\text{Pic}(F)$ . Since  $[R - 2D] = 0 \in \text{Pic}^0(F)$ , we have that

$R \in 2\text{Pic}(F)$ . Moreover, if we let  $[T] \in \text{Pic}^0(F)[2]$ , then

$$[R - 2D] = [R - 2D] - [2T] = [R - 2(D + T)]. \quad (5.3.1)$$

Thus, in the case where  $R - 2D$  is principal, we have  $R \in 2\text{Div}(F)$  and  $D$  is unique up to an order 2 element of  $\text{Pic}(F)$ . The fact that we cannot determine  $D$  exactly requires us to compute  $\text{Pic}(F)$  to carry out the desired computations. This forces us to work over  $\mathbb{F}_q$  where Picard group computations are implemented.

The other task is to determine when the quadratic extension  $F(\sqrt{f})$  over  $F$  is Galois (as an absolute extension of  $\mathbb{F}_q(x)$ ) given that  $F$  is Galois over  $\mathbb{F}_q(x)$ . Kummer theory tells us precisely when such an extension is Galois in the following Lemma.

**Lemma 5.3.2.** *Let  $F$  be a Galois extension of  $\mathbb{F}_q(x)$  with Galois group  $G$  and let  $f \in F^\times/F^{\times 2}$ . Then the quadratic extension  $F(\sqrt{f})$  is Galois as an absolute extension of  $\mathbb{F}_q(x)$  if and only if  $\sigma(f)/f$  is a square in  $F$  for every  $\sigma \in G$ .*

We can now formulate these concepts into an algorithm over  $\mathbb{F}_q$ .

## Section 5.4

### An algorithm over $\mathbb{F}_q$

Let  $F$  be function field with field of constants  $\mathbb{F}_q$  with  $q = p^r$  and  $p \neq 2$ . Let  $\mathbb{F}_q(x)$  denote the rational function field in the variable  $x$ .

**Definition 5.4.1.** A 2-group Belyi map modulo  $q$  is a Galois extension of function fields  $\mathbb{F}_q(x) \hookrightarrow F$  with  $[F : \mathbb{F}_q(x)] = 2^m$  unramified outside of all places above  $\{0, 1, \infty\}$ .



*Remark 5.4.2.* A Belyi map  $\phi: X \rightarrow \mathbb{P}^1$  over a field of characteristic  $p$  where  $p \nmid \text{Mon}(\phi)$  is called **tame**. The theory of tame Belyi maps is the same as in characteristic zero.

**Definition 5.4.3.** Two Belyi maps  $\mathbb{F}_q(x) \hookrightarrow F_1$  and  $\mathbb{F}_q(x) \hookrightarrow F_2$ , are **isomorphic** if there exists an isomorphism  $\varphi: F_1 \rightarrow F_2$  of function fields over  $\mathbb{F}_q(x)$  such that the diagram

$$\begin{array}{ccc} F_1 & \xrightarrow{\varphi} & F_2 \\ & \searrow & \swarrow \\ & \mathbb{F}_q(x) & \end{array} \quad (5.4.4)$$

commutes.

We now describe the algorithms to iteratively compute 2-group Belyi maps modulo  $q$ . The basic idea is to compute a tower of quadratic extensions by extracting square roots to work our way up the tower. Since we are concerned with Galois Belyi maps, we want to make sure that each intermediate field is Galois as an absolute extension of  $\mathbb{F}_q(x)$ . To start, we describe the degree 2 Belyi maps.

**Lemma 5.4.5.** *The three degree 2 Belyi maps modulo  $q$  up to isomorphism are a*

$$F_{(1,2,2)} = \frac{\mathbb{F}_q(x)[y]}{(y^2 + x - 1)}, \quad F_{(2,1,2)} = \frac{\mathbb{F}_q(x)[y]}{(y^2 - x)}, \quad \text{and} \quad F_{(2,2,1)} = \frac{\mathbb{F}_q(x)[y]}{(y^2 - x^2 + x)}. \quad (5.4.6)$$

*Proof.* All three degree 2 passports  $(0, \mathbb{Z}/2\mathbb{Z}, (1, 2, 2))$ ,  $(0, \mathbb{Z}/2\mathbb{Z}, (2, 1, 2))$ , and  $(0, \mathbb{Z}/2\mathbb{Z}, (2, 2, 1))$  have size 1 by Theorem 3.5.1 and Theorem 3.5.3. Since each field is a 2-group Belyi map with passport specified by its subscript with no two isomorphic, this is an exhaustive list.  $\square$

Next, we discuss the algorithms to test when a quadratic extension is Galois (over

the rational function field).

**Algorithm 5.4.7** (IsGalois).

**Input:**

- $F$  a Galois extension of  $\mathbb{F}_q(x)$
- $\text{Gal}(F | \mathbb{F}_q(x))$  explicitly given as automorphisms of  $F$
- $f \in F$

**Output:** True if the quadratic extension  $F(\sqrt{f})$  of  $F$  is a Galois extension over  $\mathbb{F}_q(x)$  and False otherwise

1. For each generator  $\sigma$  of  $\text{Gal}(F | \mathbb{F}_q(x))$  test if  $\sigma(f)/f$  is a square in  $F$ .
2. If  $\sigma(f)/f$  is a square in  $F$  for all generators  $\sigma$ , then return True otherwise return False.

*Proof of correctness.* The correctness of this algorithm follows from Kummer theory as discussed in Lemma 5.3.2. It suffices to test on generators since the property of being a square is multiplicative.  $\square$

**Algorithm 5.4.8** (IsGaloisOverExtension).

**Input:**

- $F$  a Galois extension of  $\mathbb{F}_q(x)$
- $\text{Gal}(F | \mathbb{F}_q(x))$  explicitly given as automorphisms of  $F$
- $f \in F$

#### 5.4 AN ALGORITHM OVER $\mathbb{F}_q$

---

Let  $F'$  be the function field  $F$  with the constant field extended from  $\mathbb{F}_q$  to  $\mathbb{F}_{q^2}$ .

**Output:** **True** if the quadratic extension  $F(\sqrt{f})$  of  $F$  is a Galois extension over  $\mathbb{F}_{q^m}(x)$  after extending the field of constants from  $\mathbb{F}_q$  to  $\mathbb{F}_{q^m}$  (for some positive integer  $m$ ) and **False** otherwise

1. For each generator  $\sigma$  of  $\text{Gal}(F' | \mathbb{F}_{q^2}(x))$  test if  $\sigma(f)/f$  is a square in  $F'$ .
2. If  $\sigma(f)/f$  is a square in  $F'$  for all generators  $\sigma$ , then return **True** otherwise return **False**.

*Proof of correctness.* The proof is similar to the previous algorithm. The proof that it is sufficient to check if elements are square over  $\mathbb{F}_{q^2}$  can be found in [35, Corollary 3.7.4]. □

The next algorithm details the process of finding the appropriate candidate function to obtain a quadratic extension by extracting a square root.

**Algorithm 5.4.9** (GetCandidateFunctions).

**Input:**

- $F$  a 2-group Belyi map modulo  $q$  of degree  $d = 2^m$  corresponding to a 2-group permutation triple  $\sigma$
- A passport  $\mathcal{P} = (\tilde{G}, (a, b, c))$  with  $\tilde{G}$  a 2-group of order  $2d$  such that there exists a 2-group permutation triple  $\tilde{\sigma}$  with passport  $\mathcal{P}$  that is a lift of  $\sigma$
- $\text{Gal}(F | \mathbb{F}_q(x)) \cong \langle \sigma \rangle$  explicitly given as automorphisms of  $F$

**Output:** A list of candidate functions  $\{f_i\}$  with each  $f_i \in F$  such that  $F(\sqrt{f_i})$  is a 2-group Belyi map modulo  $q$  with passport  $\mathcal{P}$ .

1. For  $s \in \{0, 1, \infty\}$  compute

$$r_s := \begin{cases} 0 & \text{if } \text{order}(\sigma_s) = \text{order}(\tilde{\sigma}_s) \\ 1 & \text{if } \text{order}(\sigma_s) < \text{order}(\tilde{\sigma}_s) \end{cases} \quad (5.4.10)$$

2. Compute

$$R := \sum_{s \in \{0, 1, \infty\}} r_s R_s \in \text{Div}(F) \quad (5.4.11)$$

where  $R_0, R_1, R_\infty$  are defined to be the supports of  $\text{div}(x)$ ,  $\text{div}(x - 1)$ , and  $\text{div}(1/x)$  respectively.

3. Compute the abelian group  $\text{Pic}(F) = T \oplus \mathbb{Z}$  (with  $T$  a finite abelian group) along with a map  $\psi: \text{Pic}(F) \rightarrow \text{Div}(F)$ .
4. Compute  $[R] := \psi^{-1}(R)$ .
5. For each  $a \in \text{Pic}(F)[2]$  compute the following:

- (a) Let  $D_a := \psi(a + [R]/2) \in \text{Div}(F)$ .
- (b) Compute  $\mathcal{L}(R - 2D_a)$ .
- (c) If  $\mathcal{L}(R - 2D_a)$  has dimension 1, then compute  $f_a \in F$  with  $\text{div}(f_a)$  generating  $\mathcal{L}(R - 2D_a)$  and go to Step 5d. Otherwise go to the next  $a \in \text{Pic}(F)[2]$ .
- (d) Apply Algorithm 5.4.7 to  $F$ ,  $\text{Gal}(F | \mathbb{F}_q(x))$ , and  $f_a$  from Step 5c to see if  $F(\sqrt{f_a})$  generates a Galois extension. If  $F(\sqrt{f_a})$  is Galois over  $\mathbb{F}_q(x)$  then save  $f_a$  and go to the next  $a \in \text{Pic}(F)[2]$ . If  $F(\sqrt{f_a})$  is not Galois over  $\mathbb{F}_q(x)$ , then go to Step 5e.

- (e) Let  $F'$  be the function field  $F$  after extending the field of constants  $\mathbb{F}_q$  to  $\mathbb{F}_{q^2}$ . Apply Algorithm 5.4.8 to  $F'$ ,  $\text{Gal}(F' | \mathbb{F}_{q^2}(x))$ , and  $f_a$  (viewed as an element of  $F'$ ) from Step 5c to see if  $F'(\sqrt{f_a})$  generates a Galois extension. If  $F(\sqrt{f_a})$  is Galois over  $\mathbb{F}_{q^2}(x)$  then save  $f_a$ . Go to the next  $a \in \text{Pic}(F)[2]$ .
6. Let  $S$  be the set of  $f_a$  saved in Step 5d. Let  $S'$  be the set of  $f_a$  saved in Step 5e.
7. • If  $S$  is nonempty, then for each  $f_a \in S$  compute  $F(\sqrt{f_a})$ ,

$$G_a \cong \text{Gal}(F(\sqrt{f_a}) | \mathbb{F}_q(x)),$$

and let  $S'' = \{f_a \in S : G_a \cong \tilde{G}\}$ .

- If  $S$  is empty, then for each  $f_a \in S'$  compute  $F'(\sqrt{f_a})$ ,

$$G_a \cong \text{Gal}(F'(\sqrt{f_a}) | \mathbb{F}_{q^2}(x)),$$

and let  $S'' = \{f_a \in S' : G_a \cong \tilde{G}\}$ .

8. Return the list  $S''$  from Step 7.

*Proof of correctness.* First, note that since we enumerated the isomorphism classes of 2-group Belyi maps in Chapter 3, we know the size of each passport  $\mathcal{P}$  as input to this algorithm. The divisor  $R$  computed in Step 2 encodes the ramification required to obtain a 2-group Belyi map with ramification matching the passport  $\mathcal{P}$ . From the discussion in Section 5.3,  $R \in 2\text{Div}(F)$ , and we can find all solutions to the equation

$$[R - 2D] = [0] \tag{5.4.12}$$

in  $\text{Pic}(F)$ . For every element  $a \in \text{Pic}(F)[2]$  we get a solution to (5.4.12). More precisely, the divisor  $D_a$  computed in Step 5a satisfies  $[R - 2D_a] = [0]$ , and all solutions to (5.4.12) are of the form  $D_a$  for some  $a \in \text{Pic}(F)[2]$ . Now, since  $R - 2D_a$  is principal for each  $a$ , we can find a candidate function  $f_a \in F$  with  $\text{div}(f_a) = R - 2D_a$ . After collecting the candidate functions  $f_a$ , we first use Algorithm 5.4.7 and Algorithm 5.4.8 to eliminate  $f_a$  that do not generate Galois extensions. Lastly, in Step 7, we only keep candidate functions  $f_a$  that generate extensions with Galois group isomorphic to the group  $\tilde{G}$  specified by the passport  $\mathcal{P}$ . Algorithm 5.4.7 and Algorithm 5.4.8 guarantee that that no further constant field extension is required.  $\square$

The next algorithm details the process of extracting a square root of a candidate function (obtained from the output of Algorithm 5.4.9) and lifting automorphisms.

**Algorithm 5.4.13** (LiftBelyiMap).

**Input:**

- The same input as in Algorithm 5.4.9
- Additionally, a specific  $f_a$  from the output of Algorithm 5.4.9

**Output:** A 2-group Belyi map modulo  $q$  with passport  $\mathcal{P}$  and explicit automorphisms identified with its Galois group  $\tilde{G}$

1. Compute  $m_{f_a, \mathbb{F}_q(x)} \in \mathbb{F}_q(x)[y]$  the minimal polynomial of  $f_a$  over  $\mathbb{F}_q(x)$  and let  $\alpha$  be a root of  $m_{f_a, \mathbb{F}_q(x)}(y^2)$ . Let  $\tilde{F}$  denote the extension  $\mathbb{F}_q(x)(\alpha)$ .
2. Let  $m_{\alpha, \mathbb{F}_q(x)}$  be the minimal polynomial of  $\alpha$  over  $\mathbb{F}_q(x)$  and compute the set

$$R := \{r : r \text{ is a root of } m_{\alpha, \mathbb{F}_q(x)} \text{ in } \tilde{F}\}. \quad (5.4.14)$$

3. Return the following:

- The absolute extension  $\tilde{F}$  of  $\mathbb{F}_q(x)$
- The set of field automorphisms  $\{\tau_r : r \in R\}$  where  $\tau_r : \tilde{F} \rightarrow \tilde{F}$  is defined by  $\alpha \mapsto r$ .

*Proof of correctness.* Since  $f_a$  is obtained from the output of Algorithm 5.4.9, the extension  $\tilde{F}$  is Galois so that  $m_{\alpha, \mathbb{F}_q(x)}$  has exactly  $\deg(\tilde{F})$  roots in  $\tilde{F}$ . Again by Algorithm 5.4.9, the extension  $\tilde{F}$  defines a 2-group Belyi map modulo  $q$  with passport  $\mathcal{P}$ . The maps  $\tau_r : \alpha \mapsto r$  define  $\deg(\tilde{F})$  automorphisms of  $\tilde{F}$  over  $\mathbb{F}_q(x)$ .  $\square$

With Algorithm 5.4.9 and Algorithm 5.4.13 at our disposal, we can now explain how to compute *all* 2-group Belyi maps with a given passport.

**Algorithm 5.4.15** (ComputePassport).

**Input:**

- A passport  $\mathcal{P}_{\text{above}} = (\tilde{G}, (a, b, c))$
- A list of passports  $\mathcal{P}_1, \dots, \mathcal{P}_k$
- For each  $\mathcal{P}_i$  a list of triples of data  $(\sigma_i^1, F_i^1, G_i^1), \dots, (\sigma_i^{\#P_i}, F_i^{\#P_i}, G_i^{\#P_i})$  with the  $F_i^j$  pairwise non-isomorphic and each  $(\sigma_i^j, F_i^j, G_i^j)$  satisfying the following:
  - $\sigma_i^j$  is a 2-group permutation triple with passport  $\mathcal{P}_i$
  - There exists a 2-group permutation triple  $\tilde{\sigma}_i^j$  with passport  $\mathcal{P}_{\text{above}}$  that is a lift of  $\sigma_i^j$
  - $F_i^j$  is a 2-group Belyi map modulo  $q$

- $G_i^j$  is the Galois group of  $F_i^j$  over  $\mathbb{F}_q(x)$  explicitly given as automorphisms of  $F_i^j$

**Output:** A list of triples of data  $(\tilde{F}^1, \tilde{G}^1), \dots, (\tilde{F}^{\#\mathcal{P}_{\text{above}}}, \tilde{G}^{\#\mathcal{P}_{\text{above}}})$  with  $\tilde{F}^j$  a 2-group Belyi map modulo  $q'$  (with  $q'$  a power of  $q$ ),  $\tilde{G}^j$  the Galois group of  $F_i^j$  explicitly given as automorphisms of  $\tilde{F}^j$ , and the  $\tilde{F}^j$  pairwise non-isomorphic.

1. Apply Algorithm 5.4.9 to every triple of data  $(\sigma_i^j, F_i^j, G_i^j)$  downstairs (along with the passport  $\mathcal{P}_{\text{above}}$ ) to obtain a list of candidate functions  $\text{CFS}_q := \{f_i^{j,k}\}$  with  $f_i^{j,k} \in F_i^j$  for each  $k$ .
2. For each  $f_i^{j,k} \in \text{CFS}_q$ , apply Algorithm 5.4.13 with input  $(\sigma_i^j, F_i^j, G_i^j)$  and  $f_i^{j,k}$  to obtain  $\widetilde{F_i^{j,k}}$  a 2-group Belyi map modulo  $q$  with passport  $\mathcal{P}$  and Galois group  $\widetilde{G_i^{j,k}}$ . Let  $\text{BELYI}_q$  denote the list of all pairs  $(\widetilde{F_i^{j,k}}, \widetilde{G_i^{j,k}})$  obtained in this step.
3. Test isomorphism of fields  $\widetilde{F_i^{j,k}}$  and  $\widetilde{F_i^{j,k'}}$  over  $\mathbb{F}_q(x)$  for each pair of fields in  $\text{BELYI}_q$ . Keep exactly one representative of each isomorphism class from  $\text{BELYI}_q$  and store this data (including the Galois group) in  $\text{BELYIISO}_q$ .
4. If  $\#\text{BELYIISO}_q = \#\mathcal{P}_{\text{above}}$  then return  $\text{BELYIISO}_q$ . Otherwise, extend the constant field from  $\mathbb{F}_q$  to  $\mathbb{F}_{q^2}$  and repeat Steps 1, 2, and 3 to obtain lists  $\text{CFS}_{q^2}$ ,  $\text{BELYI}_{q^2}$ , and  $\text{BELYIISO}_{q^2}$ . Continue this process for  $q, q^2, q^3, \dots$  and return  $\text{BELYIISO}_{q^m}$  for the first  $m \in \{1, 2, \dots\}$  with the same cardinality as  $\mathcal{P}_{\text{above}}$ .

*Proof of correctness.* This algorithm is largely bookkeeping and applying Algorithm 5.4.9 and Algorithm 5.4.13. The triples of data downstairs enumerate the isomorphism classes of 2-group Belyi maps modulo  $q$  in all passports that have a representative with a lift that has passport  $\mathcal{P}_{\text{above}}$ .



It is important to note at this point that for every downstairs passport  $\mathcal{P}_i$ , we need *all*  $\#\mathcal{P}_i$  triples of data  $(\sigma_i^j, F_i^j, G_i^j)$ . This is because Algorithm 5.4.9 only identifies candidate functions that produce 2-group Belyi maps with the correct passport. It does not provide a way to identify the precise isomorphism class. That is why, in this algorithm, we must be content with a list of pairwise non-isomorphic Belyi maps. By testing isomorphisms we can ensure that we have a representative from every isomorphism class, but we cannot identify the permutation triple corresponding to a Belyi map. In this algorithm the permutation triples (obtained from the algorithms in Section 3.4) are simply a bookkeeping tool.

The one subtle point is explaining how to obtain the  $q'$ . After each round of computing the lists  $\text{CFS}_{q^i}$ ,  $\text{BELYI}_{q^i}$ , and  $\text{BELYIISO}_{q^i}$  it is possible that we failed to find all candidate functions over the constant field  $\mathbb{F}_{q^i}$ . The enumeration of isomorphism classes in Section 3.4 ensures that this process of extending the constant field will terminate, but does not provide an a priori bound on the  $q'$  required.  $\square$

Applying Algorithm 5.4.15 to every degree  $d$  passport allows us to enumerate all 2-group Belyi maps modulo  $q$  *one degree at a time*. Section 5.6 details how far we were able to push these computations in practice using [24].

*Remark 5.4.16.* The algorithms in this section rely on the **Magma** implementations to compute class groups  $\text{Pic}(F)$  for global function fields, and the implementations to compute Riemann-Roch spaces  $\mathcal{L}(D)$ .

We conclude this section with an example of how these algorithms are applied in a specific example.

*Example 5.4.17.* In this example we use the algorithms in this section to compute all three 2-group Belyi maps modulo 3 with passport  $(3, G, (4, 4, 4))$  where  $G = (\mathbb{Z}/4\mathbb{Z}) :$

## 5.5 AN IMPLEMENTATION OVER $\mathbb{Q}^{\text{al}}$

---

$(\mathbb{Z}/4\mathbb{Z})$  is the Galois group described at the following link.

<http://www.lmfdb.org/GaloisGroup/16T8>

The Magma script can be run using the following command from the repository [24].

Shell

```
magma thesis_examples/compute_passport_16T8_444_g3.m
```

The source code in this file is as follows.

Magma

```
load "config.m";
SetVerbose("TwoDBPassport", 3);
SetVerbose("TwoDB", 1);
objs := GetPassportObjects(16);
s := objs[#objs-2];
ComputeBelyiMaps(s : optimized := false);
```

This example has 7 size 1 passports  $\mathcal{P}_i$  as in Algorithm 5.4.15. Each isomorphism class downstairs yields a candidate Belyi map upstairs, and the isomorphism checking in Algorithm 5.4.15 Step 3 correctly identifies the 3 distinct isomorphism classes out of 7.

### Section 5.5

## An implementation over $\mathbb{Q}^{\text{al}}$

We now discuss the situation in characteristic zero. The procedure has the same broad strokes as that in characteristic  $p \neq 2$ , but there is a key difference in the technique to get candidate functions to extract a square root of in Algorithm 5.4.9.

In characteristic zero there is no implementation to compute  $\text{Pic}(F)$  (for general  $F$ ). To show how we can get around this (in some cases), we now rewrite Algorithm 5.4.9 in the characteristic zero setting.

**Algorithm 5.5.1.**

**Input:**

- $K(x) \hookrightarrow F := K(X)$  a 2-group Belyi map of degree  $d = 2^m$  corresponding to a 2-group permutation triple  $\sigma$
- A passport  $\mathcal{P} = (\tilde{G}, (a, b, c))$  with  $\tilde{G}$  a 2-group of order  $2d$  such that there exists a 2-group permutation triple  $\tilde{\sigma}$  with passport  $\mathcal{P}$  that is a lift of  $\sigma$
- $\text{Gal}(F | K(x)) \cong \langle \sigma \rangle$  explicitly given as automorphisms of  $F$

**Output:** A list of candidate functions  $\{f_i\}$  with each  $f_i \in F$  such that  $K(x) \hookrightarrow F(\sqrt{f_i})$  is a 2-group Belyi map with passport  $\mathcal{P}$ .

1. For  $s \in \{0, 1, \infty\}$  compute

$$r_s := \begin{cases} 0 & \text{if } \text{order}(\sigma_s) = \text{order}(\tilde{\sigma}_s) \\ 1 & \text{if } \text{order}(\sigma_s) < \text{order}(\tilde{\sigma}_s) \end{cases} \quad (5.5.2)$$

2. Compute

$$R := \sum_{s \in \{0, 1, \infty\}} r_s R_s \in \text{Div}(F) \quad (5.5.3)$$

where  $R_0, R_1, R_\infty$  are defined to be the supports of  $\text{div}(x)$ ,  $\text{div}(x - 1)$ , and  $\text{div}(1/x)$  respectively.

3. Let  $M$  denote the set  $(R+2\mathbb{Z}R) \cap \text{Div}^0(F)$  and for  $B \in \mathbb{Z}_{\geq 1}$  let  $M_B = \left\{ R+2nR : n \in \{-B, -B+1, \dots, B-1, B\} \right\} \cap \text{Div}^0(F)$ .
4. For each  $D \in M$  compute the following:
  - (a) Compute  $\mathcal{L}(D)$ .
  - (b) If  $\mathcal{L}(D)$  has dimension 1, then compute  $f_D \in F$  with  $\text{div}(f_D)$  generating  $\mathcal{L}(D)$  and go to the next step. Otherwise, go to the next  $D \in M$ .
  - (c) Check to see if  $F(\sqrt{f_D})$  is Galois. If  $F(\sqrt{f_D})$  is Galois, then save  $f_D$ . and go to the next  $D \in M$ . If  $F(\sqrt{f_D})$  is not Galois, then go to the next Step
  - (d) Let  $F'$  be the function field  $F$  after extending the field of constants to the compositum of the residue fields of all places in the support of  $D$ . Check to see if  $F'(\sqrt{f_D})$  is Galois. If  $F'(\sqrt{f_D})$  is Galois, then save  $f_D$ . Go to the next  $D \in M$ .
5. Let  $S$  be the set of  $f_D$  saved in Step 4c. Let  $S'$  be the set of  $f_a$  saved in Step 4d.
6.
  - If  $S$  is nonempty, then for each  $f_D \in S$  compute  $F(\sqrt{f_D})$ ,

$$G_D \cong \text{Gal}(F(\sqrt{f_D}) \mid K(x)),$$

and let  $S'' = \{f_D \in S : G_D \cong \tilde{G}\}$ .

- If  $S$  is empty, then for each  $f_D \in S'$  compute  $F'(\sqrt{f_D})$ ,

$$G_D \cong \text{Gal}(F'(\sqrt{f_D}) \mid K(x)),$$

and let  $S'' = \{f_D \in S' : G_D \cong \tilde{G}\}$ .

7. Return the list  $S''$  from Step 6.

Although Algorithm 5.5.1 in characteristic zero resembles Algorithm 5.4.9 over  $\mathbb{F}_q$ , the characteristic zero algorithm is unfortunately not guaranteed to find any candidate functions! This is due to the fact that we are not computing representatives of  $\text{Pic}^0(F)[2]$ .

Without enumerating representatives of  $\text{Pic}^0(F)[2]$ , the approach in characteristic zero will always be ad hoc in the sense that in Step 3 we are blindly looking at all combinations of points that yield the desired ramification. This process is guaranteed to succeed when  $F$  has class number 1, but this condition is not often satisfied (in fact there are only 8 such non-rational function fields over  $\mathbb{F}_q$ ).

This ad hoc approach does, however, allow us to compute some 2-group Belyi maps in characteristic zero. We conclude this section by describing the results of these computations along with those in positive characteristic.

## Section 5.6

### Results of computations

In this section we summarize the computations carried out in [24] based on the algorithms discussed in Section 5.4 and Section 5.5.

In characteristic 3 we were able to compute all 2-group Belyi maps modulo 3 up to degree 32. The results of these computations can be accessed in **Magma** (with working directory the repository [24]) using the following code.

**Magma**

```
load "config.m";  
  
d := 16;  
  
objs := [ReadTwoDBPassport(f) : f in PassportFileNames(d)];
```

The information for a given passport can be accessed using the following code.

**Magma**

```
s := Random(objs);  
  
FunctionFields(s);  
  
BelyiMaps(s);  
  
FunctionFieldAutomorphisms(s);
```

In addition to the systematic computation of 2-group Belyi maps modulo 3, we were also able to apply the implementation in Section 5.5 to compute hundreds of 2-group Belyi maps in characteristic zero with degrees up to 256.

We conclude this chapter with interesting examples encountered during these computations. First, in Section 5.7 we set up some notation to help with the description of these examples.

## Section 5.7

### **Naming conventions for database examples**

In this section we explain the conventions used for filenames in [24]. This will be useful for the rest of this chapter when referring to specific examples.

The first point to explain is how we deal with isomorphism classes of 2-group Belyi maps as opposed to passports. As discussed in Chapter 3, isomorphism classes of 2-group Belyi maps correspond to 2-group permutation triples each with a unique

filename of the form

$$\text{DNG-a, b, c-gE-H} \quad (5.7.1)$$

where

- D : degree in  $\{2, 4, 8, 16, 32, 64, 128, 256\}$
- N : either T or S identifying group database
- G : a positive integer identifying the group
- a : ramification index of 0 in  $\{2, 4, 8, 16, 32, 64, 128, 256\}$
- b : ramification index of 1 in  $\{2, 4, 8, 16, 32, 64, 128, 256\}$  (5.7.2)
- c : ramification index of  $\infty$  in  $\{2, 4, 8, 16, 32, 64, 128, 256\}$
- g : just the letter g
- E : the genus in  $\mathbb{Z}_{\geq 0}$
- H : the hash of the 2-group permutation triple a positive integer

For example, the filename 16T12-4,8,2-g2-1396531181 corresponds to a degree 64 Belyi map with monodromy group identified in the transitive group database (T for transitive)

<http://magma.maths.usyd.edu.au/magma/handbook/text/753>

by 16T12. The 4,8,2 encodes the ramification above 0, 1,  $\infty$  respectively, the g2 indicates that this Belyi map has genus 2, and the 1396531181 is the hash of the permutation triple corresponding to this Belyi map.

Another example, 64S7-8,8,4-g17-1653847134 corresponds to a degree 16 Belyi map with monodromy group identified in the small group database (S for small)

<http://magma.maths.usyd.edu.au/magma/handbook/text/748>

by 64S7. The 8,8,4 encodes the ramification above 0, 1,  $\infty$  respectively, the g17 indicates that this Belyi map has genus 17, and the 1653847134 is the hash of the permutation triple corresponding to this Belyi map. In this example, the hash is necessary to distinguish this example from 64S7-8,8,4-g17-2483683244, 64S7-8,8,4-g17-623082418, and 64S7-8,8,4-g17-964508325.

A passport is given a similar filename. The only difference for a passport is that the hash is no longer required. For example, the Belyi maps with filenames 64S7-8,8,4-g17-1653847134 64S7-8,8,4-g17-2483683244, 64S7-8,8,4-g17-623082418, and 64S7-8,8,4-g17-964508325 all have passport the same passport 64S7-8,8,4-g17.

The reason for this distinction is that the algorithms for computing equations in this chapter are designed for passports. Now that we have a concise way of talking about example, we use this in the rest of the chapter to discuss examples.

## Section 5.8

### Degree 2

In degree 2, there are 3 isomorphism classes of 2-group Belyi maps. In characteristic zero they are represented by

$$\frac{\mathbb{Q}(x)[y]}{(y^2 + x - 1)}, \quad \frac{\mathbb{Q}(x)[y]}{(y^2 - x)}, \quad \text{and} \quad \frac{\mathbb{Q}(x)[y]}{(y^2 - x^2 + x)} \quad (5.8.1)$$

and each is the unique Belyi map with passport 2T1-1,2,2-g0, 2T1-2,1,2-g0, and 2T1-2,2,1-g0, respectively.



## Section 5.9

**Degree 4**

In degree 4 there are 7 isomorphism classes of 2-group Belyi maps and each has passport size 1.

All function fields are of the form  $\mathbb{Q}(x)[y]/(f(x, y))$  with  $f$  one of the polynomials in (5.9.1). The subscripts indicate the passport.

$$\begin{aligned}
 f_{4T1-1,4,4-g0} &= y^4 + x - 1 \\
 f_{4T1-2,4,4-g1} &= y^4 + x^3 - x^2 \\
 f_{4T1-4,1,4-g0} &= y^4 - x \\
 f_{4T1-4,2,4-g1} &= y^4 - x^3 + 2x^2 - x \\
 f_{4T1-4,4,1-g0} &= y^4 - x^4 + x^3 \\
 f_{4T1-4,4,2-g1} &= y^4 - x^2 + x \\
 f_{4T2-2,2,2-g0} &= y^4 + (4x - 2)y^2 + 1
 \end{aligned} \tag{5.9.1}$$

Although every function field has constant field  $\mathbb{Q}$ , lifting the automorphisms requires extending the constant field to  $\mathbb{Q}(\zeta_4)$  in all examples except for the Belyi map defined by  $f_{4T2-2,2,2-g0}$ .

Use the following code from [24] to obtain a list of these function fields in **Magma**.

**Magma**

```
load "config.m";
objs := [ReadTwoDBPassportChar0(f) : f in PassportFileNames(4)];
fields := [FunctionFields(s)[1] : s in objs];
```

## Section 5.10

**Degree 8**

In degree 8 there are 13 isomorphism classes of 2-group Belyi maps that have size 1 passports and there are 3 size 2 passports.

All function fields are of the form  $\mathbb{Q}(x)[y]/(f(x, y))$  with  $f$  one of the polynomials in (5.10.1), (5.10.2), (5.10.5), (5.10.3), (5.10.4), for size 1 passports, or (5.10.6), for size 2 passports. The subscripts indicate the passport.

$$\begin{aligned} f_{8T1-1,8,8-g0} &= y^8 + x - 1 \\ f_{8T1-8,1,8-g0} &= y^8 - x \\ f_{8T1-8,8,1-g0} &= y^8 - x^8 + x^7 \end{aligned} \tag{5.10.1}$$

$$\begin{aligned} f_{8T1-2,8,8-g2} &= y^8 + x^5 - x^4 \\ f_{8T1-8,2,8-g2} &= y^8 - x^5 + 4x^4 - 6x^3 + 4x^2 - x \\ f_{8T1-8,8,2-g2} &= y^8 - x^4 + 3x^3 - 3x^2 + x \end{aligned} \tag{5.10.2}$$

$$\begin{aligned} f_{8T4-2,2,4-g0} &= y^8 + (4x - 2)y^4 + 1 \\ f_{8T4-2,4,2-g0} &= y^8 + (8x^4 - 16x^3 + 16x - 8)y^4 + 16x^8 - 128x^7 + 448x^6 \\ &\quad - 896x^5 + 1120x^4 - 896x^3 + 448x^2 - 128x + 16 \\ f_{8T4-4,2,2-g0} &= y^8 + (-8x^4 + 16x^3)y^4 + 16x^8 \end{aligned} \tag{5.10.3}$$

$$f_{8T5-4,4,4-g_2} = y^8 + (1/2x^3 - 3/2x^2 + x)y^4 + 1/16x^6 - 1/8x^5 + 1/16x^4 \quad (5.10.4)$$

$$\begin{aligned} f_{8T2-2,4,4-g_1} &= y^8 + (8x - 4)y^6 + (4x - 4)y^5 + (16x^2 - 31/2x + 11/2)y^4 \\ &\quad + (-8x + 8)y^3 + (2x^2 + x + 1)y^2 + (-15x^2 + 18x - 3)y \\ &\quad + 4x^3 - 127/16x^2 + 35/8x + 9/16 \end{aligned}$$

$$f_{8T2-4,2,4-g_1} = y^8 + (1/16x^3 - 1/16x^2 + 1/128x)y^4 + 1/65536x^2 \quad (5.10.5)$$

$$\begin{aligned} f_{8T2-4,4,2-g_1} &= y^8 + (8x - 4)y^6 + (-1/2x^4 - 23/2x^3 + 28x^2 - 16x + 6)y^4 \\ &\quad + (6x^5 - 25x^4 + 27x^3 - 8x^2 + 8x - 4)y^2 + 1/16x^8 - 9/8x^7 \\ &\quad + 97/16x^6 - 9x^5 + 7/2x^4 + 9/2x^3 - 4x^2 + 1 \end{aligned}$$

$$\begin{aligned} f_{8T1-4,8,8-g_3} &= y^8 + x^5 - 3x^4 + 3x^3 - x^2 \\ f_{8T1-8,4,8-g_3} &= y^8 - x^5 + 2x^4 - x^3 \\ f_{8T1-8,8,4-g_3} &= y^8 - x^2 + x \end{aligned} \quad (5.10.6)$$

Although every function field has constant field  $\mathbb{Q}$ , lifting the automorphisms requires extending the constant field to  $\mathbb{Q}(\zeta_8)$  in all examples except for the those that are covers of the Belyi map defined by  $f_{4T2-2,2,2-g_0}$  in Section 5.9.

Use the following code from [24] to obtain a list of these function fields in **Magma**.

**Magma**

```
load "config.m";

objs := [ReadTwoDBPassportChar0(f) : f in PassportFileNames(8)];

fields := [FunctionFields(s)[1] : s in objs];
```

Perhaps the most interesting note to make in degree 8 is that it is the lowest degree where the characteristic zero approach appears to fail. Indeed, the genus 3 passports represented in (5.10.6) have size 2, but only a single representative Belyi map for each. In these examples, all candidate functions obtained from the adhoc approach in Algorithm 5.5.1 yield isomorphic function fields over  $\mathbb{Q}(x)$ , so we know we are missing a Belyi map in each one of these passports.

The techniques in Section 5.4, however, do manage to succeed for these size 2 passports. Working over  $\mathbb{F}_3$ , using Algorithm 5.4.15, we obtain the function fields

$$\frac{\mathbb{F}_3(x)[y]}{(y^8 + 2x^6 + 2x^5 + 2x^4 + x^3 + x^2 + x)} \quad \text{and} \quad \frac{\mathbb{F}_3(x)[y]}{(y^8 + x^6 + 2x^3)} \quad (5.10.7)$$

for the passport 8T1-8,8,4-g3, and one can check that the fields in (5.10.7) are not isomorphic.

To obtain a list of degree 8 function fields corresponding to 2-group Belyi maps in characteristic 3, use the following code from [24].

Magma

```
load "config.m";
objs := [ReadTwoDBPassport(f) : f in PassportFileNames(8)];
fields := [* *];
for s in objs do
    fields cat:= FunctionFields(s);
end for;
```

## Section 5.11

**Degree 16**

In degree 16 there are 55 isomorphism classes of 2-group Belyi maps with 41 distinct passports. All passports are of size 1 except for Passport 16T8-4,4,4-g3 which has size 3, Passports 16T1-4,16,16-g6, 16T1-16,4,16-g6, 16T1-16,16,4-g6 which have size 2, and Passports 16T1-8,16,16-g7, 16T1-16,8,16-g7, 16T1-16,16,8-g7 which have size 4.

Algorithm 5.4.15 succeeds in characteristic 3, but the equations are too large to write here. For an illustrative example of Algorithm 5.4.15 in action see Example 5.4.17. To obtain a list of degree 16 function fields corresponding to 2-group Belyi maps in characteristic 3, use the following code from [24].

**Magma**

```
load "config.m";
objs := [ReadTwoDBPassport(f) : f in PassportFileNames(16)];
fields := [* *];
for s in objs do
    fields cat:= FunctionFields(s);
end for;
```

Below are some of the characteristic zero function fields corresponding to 2-group Belyi maps computed using Algorithm 5.5.1. At this point the equations tend to be too big to fit on a page.

$$f_{16T1-1,16,16-g0} = y^{16} + x - 1 \quad (5.11.1)$$

$$f_{16T13-2,2,8-g0} = y^{16} + (4x - 2)y^8 + 1 \quad (5.11.2)$$

$$f_{16T1-8,16,16-g7} = y^{16} + x^9 - 7x^8 + 21x^7 - 35x^6 + 35x^5 - 21x^4 + 7x^3 - x^2 \quad (5.11.3)$$

$$\begin{aligned} f_{16T12-4,2,8-g2} = & y^{16} + (-1/8x^4 + 1/8x^3)y^{12} + (1/1024x^9 - 1/2048x^8 \\ & - 1/256x^7 + 1/256x^6)y^8 + (1/32768x^{12} - 1/32768x^{11})y^4 \\ & + 1/16777216x^{16} \end{aligned} \quad (5.11.4)$$

$$\begin{aligned} f_{16T10-4,2,4-g1} = & y^{16} + (-1/4x^4 + 1/2x^3 + 6x^2 + 8x - 4)y^{12} + (-5x^4 - 15x^3 + 20x^2)y^{10} \\ & + (3/128x^8 - 1/16x^7 + 19/16x^6 + 15x^5 - 83/4x^4 \\ & + 15/2x^3 + 34x^2 - 16x + 6)y^8 \\ & + (1/8x^8 - 47/8x^7 + 31/4x^6 + 6x^5 - 46x^4 + 46x^3 - 8x^2)y^6 \\ & + (-1/1024x^{12} + 1/512x^{11} - 11/128x^{10} + 17/16x^9 - 63/64x^8 \\ & - 29/8x^7 + 113/8x^6 - 13x^5 - 63/4x^4 + 79/2x^3 - 22x^2 + 8x - 4)y^4 \\ & + (3/256x^{12} - 23/256x^{11} + 1/64x^{10} + 9/16x^9 - 11/8x^8 \\ & + 9/8x^7 + 7/4x^6 - 6x^5 - x^4 + 17x^3 - 12x^2)y^2 \\ & + 1/65536x^{16} - 1/2048x^{14} + 1/512x^{13} + 3/1024x^{12} - 15/512x^{11} \\ & + 9/128x^{10} - 3/32x^9 + 35/128x^8 - 9/16x^7 + 3/16x^6 + 3/4x^4 \\ & + 1/2x^3 - 2x^2 + 1 \end{aligned} \quad (5.11.5)$$

$$\begin{aligned} f_{16T8-4,4,4-g3} = & \\ & y^{16} + (-1/4x^4 + 3/4x^3 - 3/4x^2 + 33/4x - 4)y^{12} \\ & + (-20x^3 + 40x^2 - 20x)y^{10} \\ & + (3/128x^8 - 9/64x^7 + 45/128x^6 + 481/32x^5 - 6899/128x^4 \\ & + 4455/64x^3 - 2909/128x^2 - 33/4x + 6)y^8 \\ & + (-11/2x^7 + 55/2x^6 - 55x^5 + 39x^4 + 25/2x^3 - 53/2x^2 + 8x)y^6 \\ & + (-1/1024x^{12} + 9/1024x^{11} - 9/256x^{10} + 269/256x^9 - 3287/512x^8 \\ & + 8991/512x^7 - 5305/256x^6 + 65/256x^5 + 15527/1024x^4 \\ & + 3201/1024x^3 - 1135/64x^2 + 63/4x - 4)y^4 \\ & + (-5/64x^{11} + 5/8x^{10} - 35/16x^9 + 27/8x^8 + 1/32x^7 - 65/8x^6 \\ & + 205/16x^5 - 75/8x^4 + 987/64x^3 - 49/2x^2 + 12x)y^2 \\ & + 1/65536x^{16} - 3/16384x^{15} + 33/32768x^{14} - 23/16384x^{13} \\ & - 721/65536x^{12} + 549/8192x^{11} - 2009/16384x^{10} - 995/8192x^9 \\ & + 63855/65536x^8 - 33495/16384x^7 + 78241/32768x^6 - 20131/16384x^5 \\ & - 63679/65536x^4 + 2097/1024x^3 - 157/128x^2 + 1/4x + 1 \end{aligned} \tag{5.11.6}$$

---

**Section 5.12****Degree 32**

In degree 32 there are 151 isomorphism classes of 2-group Belyi maps with 96 distinct passports. All passports are of size 1 except for the following. All size 8 passports are listed in (5.12.1), the size 6 passport is listed in (5.12.2), all size 4 passports are

## 5.12 DEGREE 32

---

listed in (5.12.3), the size 3 passport is listed in (5.12.4), and all size 2 passports are listed in (5.12.5).

$$32S1-16,32,32-g15, \quad 32S1-32,16,32-g15, \quad 32S1-32,32,16-g15 \quad (5.12.1)$$

$$32S15-8,8,8-g11 \quad (5.12.2)$$

$$32S1-8,32,32-g14, \quad 32S1-32,8,32-g14, \quad 32S1-32,32,8-g14 \quad (5.12.3)$$

$$32S6-4,4,4-g5 \quad (5.12.4)$$

$$\begin{aligned} &32S1-4,32,32-g12, \quad 32S1-32,4,32-g12, \quad 32S1-32,32,4-g12, \\ &32S10-4,4,8-g7, \quad 32S10-4,8,4-g7, \quad 32S10-8,4,4-g7, \\ &32S11-4,4,8-g7, \quad 32S11-4,8,4-g7, \quad 32S11-8,4,4-g7, \\ &32S12-4,8,8-g9, \quad 32S12-8,4,8-g9, \quad 32S12-8,8,4-g9 \\ &32S16-8,16,16-g13, \quad 32S16-16,8,16-g13, \quad 32S16-16,16,8-g13 \\ &32S17-8,16,16-g13, \quad 32S17-16,8,16-g13, \quad 32S17-16,16,8-g13 \end{aligned} \quad (5.12.5)$$

To obtain a list of degree 16 function fields corresponding to 2-group Belyi maps in characteristic 3, use the following code from [24].



**Magma**

```
load "config.m";  
  
objs := [ReadTwoDBPassport(f) : f in PassportFileNames(32)];  
  
fields := [* *];  
  
for s in objs do  
    fields cat:= FunctionFields(s);  
end for;
```

---

# Bibliography

- [1] Sybilla Beckmann, *Ramified primes in the field of moduli of branched coverings of curves*, Journal of Algebra **125** (1989), no. 1, 236–255.
- [2] Gennadii Vladimirovich Belyi, *On galois extensions of a maximal cyclotomic field*, Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya **43** (1979), no. 2, 267–276.
- [3] Yakov Berkovich and Zvonimir Janko, *Groups of prime power order volume 1*, De Gruyter, 2008.
- [4] ———, *Groups of prime power order volume 2*, De Gruyter, 2008.
- [5] ———, *Groups of prime power order volume 3*, De Gruyter, 2008.
- [6] Wieb Bosma and John Cannon, *Discovering mathematics with magma*, Springer, 2006.
- [7] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, Computational algebra and number theory (London, 1993). MR MR1484478

## BIBLIOGRAPHY

---

- [8] Kenneth S Brown, *Cohomology of groups*, vol. 87, Springer Science & Business Media, 2012.
- [9] Pete L Clark and John Voight, *Algebraic curves uniformized by congruence subgroups of triangle groups*, arXiv preprint arXiv:1506.01371 (2015).
- [10] Kevin Coombes, David Harbater, et al., *Hurwitz families and arithmetic galois groups*, Duke mathematical journal **52** (1985), no. 4, 821–839.
- [11] Lassina Dembélé, *A non-solvable galois extension of  $q$  ramified at 2 only*, Comptes Rendus Mathématique **347** (2009), no. 3-4, 111–116.
- [12] I Dolgachev, *Mckay correspondence. winter 2006/07*, Lecture notes (2009).
- [13] David Steven Dummit and Richard M Foote, *Abstract algebra*, vol. 3, Wiley Hoboken, 2004.
- [14] Hershel M Farkas and Irwin Kra, *Riemann surfaces*, Riemann surfaces, Springer, 1992, pp. 9–31.
- [15] Alexandre Grothendieck, *Esquisse d'un programme*, London Mathematical Society Lecture Note Series (1997), 5–48.
- [16] Robin Hartshorne, *Algebraic geometry*, vol. 52, Springer Science & Business Media, 2013.
- [17] Derek F Holt, *A computer program for the calculation of a covering group of a finite group*, Journal of Pure and Applied Algebra **35** (1985), 287–295.
- [18] Derek F. Holt, *The mechanical computation of first and second cohomology groups*, Journal of Symbolic Computation **1** (1985), no. 4, 351–361.

## BIBLIOGRAPHY

---

- [19] G Jones and Manfred Streit, *Galois groups, monodromy groups and cartographic groups*, LONDON MATHEMATICAL SOCIETY LECTURE NOTE SERIES (1997), 25–66.
- [20] Michael Klug, Michael Musty, Sam Schiavone, and John Voight, *Numerical calculation of three-point branched covers of the projective line*, LMS Journal of Computation and Mathematics **17** (2014), no. 01, 379–430.
- [21] Cemile Kürkoğlu, *Exceptional belyi coverings*, Ph.D. thesis, bilkent university, 2015.
- [22] Gunter Malle and Bernd Heinrich Matzat, *Inverse galois theory*, Springer, 1999.
- [23] Rick Miranda, *Algebraic curves and riemann surfaces*, vol. 5, American Mathematical Soc., 1995.
- [24] Michael Musty, *2 group dessins*, <https://github.com/michaelmusty/2GroupDessins>, 2019.
- [25] Michael Musty, Sam Schiavone, Jeroen Sijsling, and John Voight, *A database of belyi maps*, arXiv preprint arXiv:1805.07751 (2018).
- [26] David Roberts, *Fractalized cyclotomic polynomials*, Proceedings of the American Mathematical Society **135** (2007), no. 7, 1959–1967.
- [27] David P Roberts, *Nonsolvable polynomials with field discriminant 5a*, International Journal of Number Theory **7** (2011), no. 02, 289–322.
- [28] Michael Rosen, *Number theory in function fields*, vol. 210, Springer Science & Business Media, 2013.

## BIBLIOGRAPHY

---

- [29] Jean-Pierre Serre, *Topics in galois theory*, AK Peters/CRC Press, 2016.
- [30] Tanush Shaska, *Determining the automorphism group of a hyperelliptic curve*, Proceedings of the 2003 International Symposium on Symbolic and Algebraic Computation, ACM, 2003, pp. 248–254.
- [31] Jeroen Sijsling and John Voight, *On computing belyi maps, numéro consacré au trimestre “méthodes arithmétiques et applications”, automne 2013*, Publ. Math. Besançon Algèbre Théorie Nr **2014/1** (2014), 73–131.
- [32] Joseph H Silverman, *The arithmetic of elliptic curves*, vol. 106, Springer Science & Business Media, 2009.
- [33] ———, *Advanced topics in the arithmetic of elliptic curves*, vol. 151, Springer Science & Business Media, 2013.
- [34] David Singerman, *Finitely maximal fuchsian groups*, Journal of the London Mathematical Society **2** (1972), no. 1, 29–38.
- [35] Henning Stichtenoth, *Algebraic function fields and codes*, vol. 254, Springer Science & Business Media, 2009.