



2-solvable Belyĭ maps




Michael Musty
Algebra and Number Theory Seminar
Dartmouth College
May 8, 2018

2-solvable Belyĭ maps




Michael Musty
Algebra and Number Theory Seminar
Dartmouth College
May 8, 2018

2-solvable Belyĭ maps




Michael Musty
Algebra and Number Theory Seminar
Dartmouth College
May 8, 2018

2-solvable Belyĭ maps



Michael Musty
Algebra and Number Theory Seminar
Dartmouth College
May 8, 2018

2-solvable Belyĭ maps



Michael Musty
Algebra and Number Theory Seminar
Dartmouth College
May 8, 2018



1. What is a 2-solvable Belyĭ map?
2. Motivation: Beckmann's Theorem
3. An algorithm to compute 2-solvable Belyĭ maps
 - (a) Computing permutation triples
 - (b) Computing equations
4. Examples
5. Application: Number fields obtained from 2-torsion points





Theorem (G.V. Belyĭ 1979)

A smooth projective curve X over \mathbb{C} can be defined over $\overline{\mathbb{Q}}$ if and only if there exists a branched covering of compact connected Riemann surfaces $\phi : X \rightarrow \mathbb{P}^1$ unramified (unbranched) above $\mathbb{P}^1 \setminus \{0, 1, \infty\}$.



Theorem (G.V. Belyĭ 1979)

A smooth projective curve X over \mathbb{C} can be defined over $\overline{\mathbb{Q}}$ if and only if there exists a branched covering of compact connected Riemann surfaces $\phi : X \rightarrow \mathbb{P}^1$ unramified (unbranched) above $\mathbb{P}^1 \setminus \{0, 1, \infty\}$.

Such a map is called a **Belyĭ map**.



Theorem (G.V. Belyĭ 1979)

A smooth projective curve X over \mathbb{C} can be defined over $\overline{\mathbb{Q}}$ if and only if there exists a branched covering of compact connected Riemann surfaces $\phi : X \rightarrow \mathbb{P}^1$ unramified (unbranched) above $\mathbb{P}^1 \setminus \{0, 1, \infty\}$.

Such a map is called a **Belyĭ map**.

Two Belyĭ maps $\phi : X \rightarrow \mathbb{P}^1$ and $\phi' : X' \rightarrow \mathbb{P}^1$ are **isomorphic** if there is an isomorphism $\iota : X \rightarrow X'$ such that $\phi' \iota = \phi$.





A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .



A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

The passport of a Belyĭ map $\phi : X \rightarrow \mathbb{P}^1$ is $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$ with $g(X)$ the genus of X , $\text{Mon}(\phi)$ the monodromy group of ϕ , and the partitions specified by ramification.



A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

The passport of a Belyĭ map $\phi : X \rightarrow \mathbb{P}^1$ is $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$ with $g(X)$ the genus of X , $\text{Mon}(\phi)$ the monodromy group of ϕ , and the partitions specified by ramification.

There is an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on Belyĭ maps.



A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

The passport of a Belyĭ map $\phi : X \rightarrow \mathbb{P}^1$ is $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$ with $g(X)$ the genus of X , $\text{Mon}(\phi)$ the monodromy group of ϕ , and the partitions specified by ramification.

There is an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on Belyĭ maps. This action preserves passports.



Passports of permutation triples



A **transitive permutation triple** is a triple

$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.



A **transitive permutation triple** is a triple $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.

Two such triples σ and σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$



A **transitive permutation triple** is a triple

$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.

Two such triples σ and σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$



A **transitive permutation triple** is a triple $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.

Two such triples σ and σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$ where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$



A **transitive permutation triple** is a triple

$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.

Two such triples σ and σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$ where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \# \text{cycles of } \tau,$$



A **transitive permutation triple** is a triple $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.

Two such triples σ and σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$ where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \# \text{cycles of } \tau,$$

and $\lambda(\sigma)$ is specified by cycle structures.



A **transitive permutation triple** is a triple

$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ with $\langle \sigma \rangle$ a transitive subgroup of S_d and $\sigma_\infty \sigma_1 \sigma_0 = 1$.

Two such triples σ and σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$ where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \# \text{cycles of } \tau,$$

and $\lambda(\sigma)$ is specified by cycle structures.

The **size** of a passport \mathcal{P} is the number of simultaneous conjugacy classes of transitive permutation triples with passport \mathcal{P} .

A group-theoretic description of Belyĭ maps







Lemma

The set of transitive permutation triples of degree d up to simultaneous conjugation is in bijection with the set of Belyĭ maps of degree d up to isomorphism.











All up to the appropriate version of equivalence in each category.

Example 2T1-2, 1, 2-g0













Theorem (Beckmann-Kazez 1989)

Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map with monodromy group G .



Theorem (Beckmann-Kazez 1989)

*Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map with monodromy group G .
Suppose p does not divide $\#G$.*



Theorem (Beckmann-Kazez 1989)

Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map with monodromy group G . Suppose p does not divide $\#G$. Then there exists a number field M such that p is unramified in M and



Theorem (Beckmann-Kazez 1989)

Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map with monodromy group G . Suppose p does not divide $\#G$. Then there exists a number field M such that p is unramified in M and ϕ is defined over M with good reduction at all primes \mathfrak{p} of M above p .



Theorem (Beckmann-Kazez 1989)

Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map with monodromy group G . Suppose p does not divide $\#G$. Then there exists a number field M such that p is unramified in M and ϕ is defined over M with good reduction at all primes \mathfrak{p} of M above p .

Upshot:



Theorem (Beckmann-Kazez 1989)

Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map with monodromy group G . Suppose p does not divide $\#G$. Then there exists a number field M such that p is unramified in M and ϕ is defined over M with good reduction at all primes \mathfrak{p} of M above p .

Upshot: Every 2-solvable Belyĭ curve has a model with good reduction away from $p = 2$.

Computing 2-solvable permutation triples



Computing 2-solvable permutation triples



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$.



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. We want to find $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$ corresponding to $\tilde{\sigma} \in S_{2d}^3$ such that



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. We want to find $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$ corresponding to $\tilde{\sigma} \in S_{2d}^3$ such that

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow \tilde{\phi} & \searrow 2 & \\ & X & \\ & \swarrow \phi & \\ & \mathbb{P}^1 & \end{array}$$



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. We want to find $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$ corresponding to $\tilde{\sigma} \in S_{2d}^3$ such that

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow \tilde{\phi} & \searrow 2 & \\ & X & \\ & \swarrow \phi & \\ & \mathbb{P}^1 & \end{array}$$

Such a $\tilde{\sigma}$ sits in the following exact sequence of groups:



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. We want to find $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$ corresponding to $\tilde{\sigma} \in S_{2d}^3$ such that

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow \tilde{\phi} & \searrow 2 & \\ & X & \\ & \swarrow \phi & \\ & \mathbb{P}^1 & \end{array}$$

Such a $\tilde{\sigma}$ sits in the following exact sequence of groups:

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1.$$

Computing 2-solvable permutation triples





Theorem

Given all 2-solvable permutation triples of degree 2^ℓ , there is an effective algorithm to compute all 2-solvable permutation triples of degree $2^{\ell+1}$.



Theorem

Given all 2-solvable permutation triples of degree 2^ℓ , there is an effective algorithm to compute all 2-solvable permutation triples of degree $2^{\ell+1}$.

Moreover, these computations have been explicitly carried out up to degree 128.



Theorem

Given all 2-solvable permutation triples of degree 2^ℓ , there is an effective algorithm to compute all 2-solvable permutation triples of degree $2^{\ell+1}$.

Moreover, these computations have been explicitly carried out up to degree 128.

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.



Theorem

Given all 2-solvable permutation triples of degree 2^ℓ , there is an effective algorithm to compute all 2-solvable permutation triples of degree $2^{\ell+1}$.

Moreover, these computations have been explicitly carried out up to degree 128.

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.

This allows us to efficiently compute all equivalence classes of extensions for a given permutation triple.



Theorem

Given all 2-solvable permutation triples of degree 2^ℓ , there is an effective algorithm to compute all 2-solvable permutation triples of degree $2^{\ell+1}$.

Moreover, these computations have been explicitly carried out up to degree 128.

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.

This allows us to efficiently compute all equivalence classes of extensions for a given permutation triple.

For each extension, we get 8 possible $\tilde{\sigma}$.



Theorem

Given all 2-solvable permutation triples of degree 2^ℓ , there is an effective algorithm to compute all 2-solvable permutation triples of degree $2^{\ell+1}$.

Moreover, these computations have been explicitly carried out up to degree 128.

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.

This allows us to efficiently compute all equivalence classes of extensions for a given permutation triple.

For each extension, we get 8 possible $\tilde{\sigma}$. We then check the necessary conditions and do some bookkeeping.

Passport counts



degree	2	4	8	16	32	64	128
# genus 0 passports	3	4	6	6	6	6	6
# genus 1 passports		3	3	3	3	3	3
# genus 2 passports			4	6	0	0	0
# genus 3 passports			3	8	12	0	0
# genus 4 passports				6	6	0	0
# genus 5 passports				6	8	12	0
# genus 6 passports				3	0	0	0
# genus 7 passports				3	18	12	0
# genus 8 passports					6	6	0
# genus 9 passports					15	18	24
# genus 11 passports					7	12	0
# genus 12 passports					3	0	0
# genus 13 passports					6	30	12
# genus 14 passports					3	0	0
# genus 15 passports					3	18	12
# genus 16 passports						6	6

Passport counts



degree	2	4	8	16	32	64	128
# genus 17 passports						39	25
# genus 19 passports						18	0
# genus 21 passports						30	48
# genus 23 passports						9	12
# genus 24 passports						3	0
# genus 25 passports						24	78
# genus 27 passports						6	0
# genus 28 passports						3	0
# genus 29 passports						6	30
# genus 30 passports						3	0
# genus 31 passports						3	18
# genus 32 passports							6
# genus 33 passports							117
# genus 37 passports							114
# genus 39 passports							18
# genus 41 passports							93

Passport counts



degree	2	4	8	16	32	64	128
# genus 45 passports							48
# genus 47 passports							9
# genus 48 passports							3
# genus 49 passports							72
# genus 53 passports							26
# genus 55 passports							6
# genus 56 passports							3
# genus 57 passports							24
# genus 59 passports							6
# genus 60 passports							3
# genus 61 passports							6
# genus 62 passports							3
# genus 63 passports							3
total passports	3	7	16	41	96	267	834



Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$.

Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. Given a permutation triple $\tilde{\sigma}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. Given a permutation triple $\tilde{\sigma}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to $\tilde{\sigma}$.

Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. Given a permutation triple $\tilde{\sigma}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to $\tilde{\sigma}$. Let $X \subseteq \mathbb{A}_K^n$ with defining equations $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$.

Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. Given a permutation triple $\tilde{\sigma}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to $\tilde{\sigma}$. Let $X \subseteq \mathbb{A}_K^n$ with defining equations $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$. Our goal is to find $f \in K(X)^\times$ such that

Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. Given a permutation triple $\tilde{\sigma}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to $\tilde{\sigma}$. Let $X \subseteq \mathbb{A}_K^n$ with defining equations $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$. Our goal is to find $f \in K(X)^\times$ such that

$$\begin{array}{ccc} \tilde{X} & & \frac{\overline{K(X)}[y]}{(y^2-f)} \\ \downarrow \tilde{\phi} & \searrow \psi & \downarrow \\ & X & \overline{K(X)} \\ & \swarrow \phi & \uparrow d \\ \mathbb{P}^1 & & \overline{K(\mathbb{P}^1)} \end{array}$$

Computing 2-solvable Belyĭ maps



Let $\phi : X \rightarrow \mathbb{P}^1$ be a Belyĭ map of degree $d = 2^\ell$ corresponding to $\sigma \in S_d^3$. Given a permutation triple $\tilde{\sigma}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to $\tilde{\sigma}$. Let $X \subseteq \mathbb{A}_K^n$ with defining equations $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$. Our goal is to find $f \in K(X)^\times$ such that

$$\begin{array}{ccc} \tilde{X} & & \frac{\overline{K(X)}[y]}{(y^2-f)} \\ \downarrow \tilde{\phi} & \searrow \psi & \downarrow \\ & X & \overline{K(X)} \\ & \swarrow \phi & \uparrow d \\ \mathbb{P}^1 & & \overline{K(\mathbb{P}^1)} \end{array}$$

with ψ (and hence $\tilde{\phi}$) satisfying the ramification conditions imposed by $\tilde{\sigma}$.





The procedure to find $f \in K(X)$ is as follows:



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ .



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ . These are determined by $\tilde{\sigma}$.



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ . These are determined by $\tilde{\sigma}$.
2. Build a degree 0 divisor $D = \sum_P n_P P$ with n_{Q_i} odd for every i .



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ . These are determined by $\tilde{\sigma}$.
2. Build a degree 0 divisor $D = \sum_P n_P P$ with n_{Q_i} odd for every i .
3. Try to find f in the (computable) Riemann-Roch space $L(D)$.



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ . These are determined by $\tilde{\sigma}$.
2. Build a degree 0 divisor $D = \sum_P n_P P$ with n_{Q_i} odd for every i .
3. Try to find f in the (computable) Riemann-Roch space $L(D)$.

There are (at least) two remarks to make about this process:



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ . These are determined by $\tilde{\sigma}$.
2. Build a degree 0 divisor $D = \sum_P n_P P$ with n_{Q_i} odd for every i .
3. Try to find f in the (computable) Riemann-Roch space $L(D)$.

There are (at least) two remarks to make about this process:

- Extending the base field K may be necessary to determine D .



The procedure to find $f \in K(X)$ is as follows:

1. Let $\{Q_i\}$ be the points on X that we want to be ramification values of ψ . These are determined by $\tilde{\sigma}$.
2. Build a degree 0 divisor $D = \sum_P n_P P$ with n_{Q_i} odd for every i .
3. Try to find f in the (computable) Riemann-Roch space $L(D)$.

There are (at least) two remarks to make about this process:

- ▶ Extending the base field K may be necessary to determine D .
- ▶ Class group obstruction.






$$\tilde{\sigma} = ((1432), (13)(24), (1432)), \quad \sigma = ((12), (1)(2), (12))$$



$$\tilde{\sigma} = ((1432), (13)(24), (1432)), \quad \sigma = ((12), (1)(2), (12))$$





$$X_2 : x_2^2 = x_1$$

Diagram illustrating a mapping or transformation. It shows a vertical line connecting $X_2 : x_2^2 = x_1$ to X_1 .



$$\tilde{\sigma} = ((1432), (13)(24), (1432)), \quad \sigma = ((12), (1)(2), (12))$$







Passport: 8T1-8,4,8-g3, size 2

Belyĭ curve: $X : y^2 + (x^4 + 1)y = -2x^4$

Belyĭ map: $(y + 1)^2$



Passport: 8T1-8,4,8-g3, size 2

Belyĭ curve: $X : y^2 + (x^4 + 1)y = -2x^4$

Belyĭ map: $(y + 1)^2$

Passport: 16T1-16,8,16-g7, size 4

Belyĭ curve: $X : y^2 + (x^8 + 1)y = -2x^8$

Belyĭ map: $(y + 1)^2$



Nonhyperelliptic Belyĭ maps



$128S1-128,32,128-g62 \rightarrow 64S1-64,16,64-g30 \rightarrow$
 $32S1-32,8,32-g14 \rightarrow 16T1-16,4,16-g6 \rightarrow 8T1-8,2,8-g2 \rightarrow$
 $4T1-4,1,4-g0 \rightarrow 2T1-2,1,2-g0$



$128S1-128,32,128-g62 \rightarrow 64S1-64,16,64-g30 \rightarrow$
 $32S1-32,8,32-g14 \rightarrow 16T1-16,4,16-g6 \rightarrow 8T1-8,2,8-g2 \rightarrow$
 $4T1-4,1,4-g0 \rightarrow 2T1-2,1,2-g0$

$$\begin{aligned} X \subset \mathbb{A}^6 : & x_1^5 - x_1 - x_2^2 \\ & x_1 - x_1^3 + x_2 x_4^4 \\ & x_1^3 x_3 - x_1 x_3 - x_2 x_4^2 \\ & x_1^2 x_4^2 - x_2 x_3 + x_4^2 \\ & x_2 x_3 - x_1^2 - 1 \\ & x_3 x_4^2 - 1 \\ & x_5^2 - x_4 \\ & x_6^2 - x_5 \\ \phi : & x_3^4 x_2^2 - 2 x_3^2 x_2 + 1 \end{aligned}$$



128S69-8,16,16-g49: size 4

64S7-4,8,8-g17

32S10-4,8,4-g7

16T12-4,8,2-g2

8T4-2,4,2-g0

4T2-2,2,2-g0

2T1-2,2,1-g0

Networks?





<https://math.dartmouth.edu/~mjmusty/32.html>



<https://math.dartmouth.edu/~mjmusty/32.html>

- Is every 2-solvable Belyĭ map defined over an abelian extension of \mathbb{Q} ?



<https://math.dartmouth.edu/~mjmusty/32.html>

- ▶ Is every 2-solvable Belyĭ map defined over an abelian extension of \mathbb{Q} ?
- ▶ What can we say in the hyperelliptic case?



<https://math.dartmouth.edu/~mjmusty/32.html>

- ▶ Is every 2-solvable Belyĭ map defined over an abelian extension of \mathbb{Q} ?
- ▶ What can we say in the hyperelliptic case?
- ▶ What infinite families of 2-groups appear as monodromy groups of Belyĭ maps?



Thanks to the following for helpful discussions:

- ▶ Sam Schiavone
- ▶ Jeroen Sijsling
- ▶ John Voight



Thanks to the following for helpful discussions:

- ▶ Sam Schiavone
- ▶ Jeroen Sijsling
- ▶ John Voight

Thanks for listening!



Thanks to the following for helpful discussions:

- ▶ Sam Schiavone
- ▶ Jeroen Sijsling
- ▶ John Voight

Thanks for listening! (unless there is extra time...)

Applications?



Applications?



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K .

Applications?



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X .



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2.



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J .



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model.



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$.



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$.



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$. (Baker 1893).



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$. (Baker 1893). In general one must compute the adjoint ideal.



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$. (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology.



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$. (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology. Integrating yields a $g \times 2g$ period matrix Π



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$. (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology. Integrating yields a $g \times 2g$ period matrix Π with Λ the \mathbb{Z} -span of the columns of Π .



Let X be a 2-solvable Belyĭ curve of degree d and genus g defined over K . We would like to compute the 2-torsion field of the jacobian J of X . The field $K(J[2])/K$ will be ramified only at 2. To start, we must compute a period matrix for J . We can embed X in \mathbb{P}^2 with a singular model. Now consider an affine patch $f(x, y) = 0$. The space of holomorphic 1-forms on X is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by (i, j) where $(i + 1, j + 1)$ is an interior point of the Newton polygon of $f(x, y)$. (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology. Integrating yields a $g \times 2g$ period matrix Π with Λ the \mathbb{Z} -span of the columns of Π . J is identified with \mathbb{C}^g / Λ .

Applications?





The next piece is the Abel-Jacobi map



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left(\int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left(\int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for $t \in \frac{1}{2}\Lambda/\Lambda$,



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left(\int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for $t \in \frac{1}{2}\Lambda/\Lambda$, our task is to find $\{Q_1, \dots, Q_g\}$ with $Q_j \in X(\overline{K})$ such that



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left(\int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for $t \in \frac{1}{2}\Lambda/\Lambda$, our task is to find $\{Q_1, \dots, Q_g\}$ with $Q_j \in X(\overline{K})$ such that

$$\sum_{j=1}^g \left(\int_{P_0}^{Q_j} \omega_i \right)_{i=1, \dots, g}$$



The next piece is the Abel-Jacobi map

$$\begin{aligned} \text{AJ} : X &\rightarrow \mathbb{C}^g / \Lambda \\ P &\mapsto \left(\int_{P_0}^P \omega_j \right)_{j=1, \dots, g} \end{aligned}$$

Now for $t \in \frac{1}{2}\Lambda / \Lambda$, our task is to find $\{Q_1, \dots, Q_g\}$ with $Q_j \in X(\bar{K})$ such that

$$\sum_{j=1}^g \left(\int_{P_0}^{Q_j} \omega_i \right)_{i=1, \dots, g}$$

The coordinates of the Q_j generate the field $K(J[2])$.