



# 2-solvable Belyĭ maps




Michael Musty  
Algebra and Number Theory Seminar  
Dartmouth College  
May 8, 2018

# 2-solvable Belyĭ maps




Michael Musty  
Algebra and Number Theory Seminar  
Dartmouth College  
May 8, 2018

# 2-solvable Belyĭ maps




Michael Musty  
Algebra and Number Theory Seminar  
Dartmouth College  
May 8, 2018

# 2-solvable Belyĭ maps



Michael Musty  
Algebra and Number Theory Seminar  
Dartmouth College  
May 8, 2018

# 2-solvable Belyĭ maps



Michael Musty  
Algebra and Number Theory Seminar  
Dartmouth College  
May 8, 2018



1. What is a 2-solvable Belyĭ map?
2. Motivation: Beckmann's Theorem
3. An algorithm to compute 2-solvable Belyĭ maps
  - (a) Computing permutation triples
  - (b) Computing equations
4. Examples
5. Application: Number fields obtained from 2-torsion points





## Theorem (G.V. Belyĭ 1979)

*A smooth projective curve  $X$  over  $\mathbb{C}$  can be defined over  $\overline{\mathbb{Q}}$  if and only if there exists a branched covering of compact connected Riemann surfaces  $\phi : X \rightarrow \mathbb{P}^1$  unramified (unbranched) above  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .*





## Theorem (G.V. Belyĭ 1979)

*A smooth projective curve  $X$  over  $\mathbb{C}$  can be defined over  $\overline{\mathbb{Q}}$  if and only if there exists a branched covering of compact connected Riemann surfaces  $\phi : X \rightarrow \mathbb{P}^1$  unramified (unbranched) above  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .*

Such a map is called a **Belyĭ map**.



## Theorem (G.V. Belyĭ 1979)

*A smooth projective curve  $X$  over  $\mathbb{C}$  can be defined over  $\overline{\mathbb{Q}}$  if and only if there exists a branched covering of compact connected Riemann surfaces  $\phi : X \rightarrow \mathbb{P}^1$  unramified (unbranched) above  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ .*

Such a map is called a **Belyĭ map**.

Two Belyĭ maps  $\phi : X \rightarrow \mathbb{P}^1$  and  $\phi' : X' \rightarrow \mathbb{P}^1$  are **isomorphic** if there is an isomorphism  $\iota : X \rightarrow X'$  such that  $\phi' \iota = \phi$ .





A **passport**  $\mathcal{P}$  consists of the data  $(g, G, \lambda)$  where  $g \geq 0$  is an integer,  $G \leq S_d$  is a transitive subgroup, and  $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$  is a triple of partitions of  $d$ .



A **passport**  $\mathcal{P}$  consists of the data  $(g, G, \lambda)$  where  $g \geq 0$  is an integer,  $G \leq S_d$  is a transitive subgroup, and  $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$  is a triple of partitions of  $d$ .

The passport of a Belyĭ map  $\phi : X \rightarrow \mathbb{P}^1$  is  $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$  with  $g(X)$  the genus of  $X$ ,  $\text{Mon}(\phi)$  the monodromy group of  $\phi$ , and the partitions specified by ramification.



A **passport**  $\mathcal{P}$  consists of the data  $(g, G, \lambda)$  where  $g \geq 0$  is an integer,  $G \leq S_d$  is a transitive subgroup, and  $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$  is a triple of partitions of  $d$ .

The passport of a Belyĭ map  $\phi : X \rightarrow \mathbb{P}^1$  is  $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$  with  $g(X)$  the genus of  $X$ ,  $\text{Mon}(\phi)$  the monodromy group of  $\phi$ , and the partitions specified by ramification.

There is an action of  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  on Belyĭ maps.



A **passport**  $\mathcal{P}$  consists of the data  $(g, G, \lambda)$  where  $g \geq 0$  is an integer,  $G \leq S_d$  is a transitive subgroup, and  $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$  is a triple of partitions of  $d$ .

The passport of a Belyĭ map  $\phi : X \rightarrow \mathbb{P}^1$  is  $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$  with  $g(X)$  the genus of  $X$ ,  $\text{Mon}(\phi)$  the monodromy group of  $\phi$ , and the partitions specified by ramification.

There is an action of  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  on Belyĭ maps. This action preserves passports.

# Passports of permutation triples





# Passports of permutation triples



A **transitive permutation triple** is a triple

$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .



A **transitive permutation triple** is a triple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .

Two such triples  $\sigma$  and  $\sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$



A **transitive permutation triple** is a triple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .

Two such triples  $\sigma$  and  $\sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple  $\sigma$  is  $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$

# Passports of permutation triples



A **transitive permutation triple** is a triple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .

Two such triples  $\sigma$  and  $\sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple  $\sigma$  is  $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$  where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$



A **transitive permutation triple** is a triple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .

Two such triples  $\sigma$  and  $\sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple  $\sigma$  is  $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$  where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \#\text{cycles of } \tau,$$



A **transitive permutation triple** is a triple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .

Two such triples  $\sigma$  and  $\sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple  $\sigma$  is  $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$  where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \#\text{cycles of } \tau,$$

and  $\lambda(\sigma)$  is specified by cycle structures.



A **transitive permutation triple** is a triple  $\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$  with  $\langle \sigma \rangle$  a transitive subgroup of  $S_d$  and  $\sigma_\infty \sigma_1 \sigma_0 = 1$ .

Two such triples  $\sigma$  and  $\sigma'$  are **simultaneously conjugate** if there exists  $\tau \in S_d$  with

$$(\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = (\sigma'_0, \sigma'_1, \sigma'_\infty).$$

The passport of a permutation triple  $\sigma$  is  $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$  where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \# \text{cycles of } \tau,$$

and  $\lambda(\sigma)$  is specified by cycle structures.

The **size** of a passport  $\mathcal{P}$  is the number of simultaneous conjugacy classes of transitive permutation triples with passport  $\mathcal{P}$ .

# A group-theoretic description of Belyĭ maps









## Lemma

*The set of transitive permutation triples of degree  $d$  up to simultaneous conjugation is in bijection with the set of Belyĭ maps of degree  $d$  up to isomorphism.*

# A Zoo of Bijections









All up to the appropriate version of equivalence in each category.



# Example 2T1-2, 1, 2-g0







# 2-solvable (Galois) Belyĭ maps









## Theorem (Beckmann-Kazez 1989)

*Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map with monodromy group  $G$ .*



## Theorem (Beckmann-Kazez 1989)

*Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map with monodromy group  $G$ .  
Suppose  $p$  does not divide  $\#G$ .*



## Theorem (Beckmann-Kazez 1989)

*Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map with monodromy group  $G$ .*

*Suppose  $p$  does not divide  $\#G$ . Then there exists a number field  $M$  such that  $p$  is unramified in  $M$  and*



## Theorem (Beckmann-Kazez 1989)

*Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map with monodromy group  $G$ . Suppose  $p$  does not divide  $\#G$ . Then there exists a number field  $M$  such that  $p$  is unramified in  $M$  and  $\phi$  is defined over  $M$  with good reduction at all primes  $\mathfrak{p}$  of  $M$  above  $p$ .*



## Theorem (Beckmann-Kazez 1989)

*Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map with monodromy group  $G$ . Suppose  $p$  does not divide  $\#G$ . Then there exists a number field  $M$  such that  $p$  is unramified in  $M$  and  $\phi$  is defined over  $M$  with good reduction at all primes  $\mathfrak{p}$  of  $M$  above  $p$ .*

**Upshot:**



## Theorem (Beckmann-Kazez 1989)

*Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map with monodromy group  $G$ . Suppose  $p$  does not divide  $\#G$ . Then there exists a number field  $M$  such that  $p$  is unramified in  $M$  and  $\phi$  is defined over  $M$  with good reduction at all primes  $\mathfrak{p}$  of  $M$  above  $p$ .*

**Upshot:** Every 2-solvable Belyĭ curve has a model with good reduction away from  $p = 2$ .

# Computing 2-solvable permutation triples





# Computing 2-solvable permutation triples



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ .



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . We want to find  $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$  corresponding to  $\tilde{\sigma} \in S_{2d}^3$  such that



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . We want to find  $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$  corresponding to  $\tilde{\sigma} \in S_{2d}^3$  such that

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow \tilde{\phi} & \searrow 2 & \\ & X & \\ & \swarrow \phi & \\ & \mathbb{P}^1 & \end{array}$$



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . We want to find  $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$  corresponding to  $\tilde{\sigma} \in S_{2d}^3$  such that

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow \tilde{\phi} & \searrow 2 & \\ & X & \\ & \swarrow \phi & \\ & \mathbb{P}^1 & \end{array}$$

Such a  $\tilde{\sigma}$  sits in the following exact sequence of groups:



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . We want to find  $\tilde{\phi} : \tilde{X} \rightarrow \mathbb{P}^1$  corresponding to  $\tilde{\sigma} \in S_{2d}^3$  such that

$$\begin{array}{ccc} \tilde{X} & & \\ \downarrow \tilde{\phi} & \searrow 2 & \\ & X & \\ & \swarrow \phi & \\ & \mathbb{P}^1 & \end{array}$$

Such a  $\tilde{\sigma}$  sits in the following exact sequence of groups:

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1.$$

# Computing 2-solvable permutation triples





## Theorem

*Given all 2-solvable permutation triples of degree  $2^\ell$ , there is an effective algorithm to compute all 2-solvable permutation triples of degree  $2^{\ell+1}$ .*



## Theorem

*Given all 2-solvable permutation triples of degree  $2^\ell$ , there is an effective algorithm to compute all 2-solvable permutation triples of degree  $2^{\ell+1}$ .*

*Moreover, these computations have been explicitly carried out up to degree 128.*





## Theorem

*Given all 2-solvable permutation triples of degree  $2^\ell$ , there is an effective algorithm to compute all 2-solvable permutation triples of degree  $2^{\ell+1}$ .*

*Moreover, these computations have been explicitly carried out up to degree 128.*

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.



## Theorem

*Given all 2-solvable permutation triples of degree  $2^\ell$ , there is an effective algorithm to compute all 2-solvable permutation triples of degree  $2^{\ell+1}$ .*

*Moreover, these computations have been explicitly carried out up to degree 128.*

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.

This allows us to efficiently compute all equivalence classes of extensions for a given permutation triple.



## Theorem

*Given all 2-solvable permutation triples of degree  $2^\ell$ , there is an effective algorithm to compute all 2-solvable permutation triples of degree  $2^{\ell+1}$ .*

*Moreover, these computations have been explicitly carried out up to degree 128.*

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.

This allows us to efficiently compute all equivalence classes of extensions for a given permutation triple.

For each extension, we get 8 possible  $\tilde{\sigma}$ .



## Theorem

*Given all 2-solvable permutation triples of degree  $2^\ell$ , there is an effective algorithm to compute all 2-solvable permutation triples of degree  $2^{\ell+1}$ .*

*Moreover, these computations have been explicitly carried out up to degree 128.*

The main tool used here is Derek Holt's algorithm (1980s) to compute the second cohomology group of a finite group.

This allows us to efficiently compute all equivalence classes of extensions for a given permutation triple.

For each extension, we get 8 possible  $\tilde{\sigma}$ . We then check the necessary conditions and do some bookkeeping.

# Passport counts



| degree               | 2 | 4 | 8 | 16 | 32 | 64 | 128 |
|----------------------|---|---|---|----|----|----|-----|
| # genus 0 passports  | 3 | 4 | 6 | 6  | 6  | 6  | 6   |
| # genus 1 passports  |   | 3 | 3 | 3  | 3  | 3  | 3   |
| # genus 2 passports  |   |   | 4 | 6  | 0  | 0  | 0   |
| # genus 3 passports  |   |   | 3 | 8  | 12 | 0  | 0   |
| # genus 4 passports  |   |   |   | 6  | 6  | 0  | 0   |
| # genus 5 passports  |   |   |   | 6  | 8  | 12 | 0   |
| # genus 6 passports  |   |   |   | 3  | 0  | 0  | 0   |
| # genus 7 passports  |   |   |   | 3  | 18 | 12 | 0   |
| # genus 8 passports  |   |   |   |    | 6  | 6  | 0   |
| # genus 9 passports  |   |   |   |    | 15 | 18 | 24  |
| # genus 11 passports |   |   |   |    | 7  | 12 | 0   |
| # genus 12 passports |   |   |   |    | 3  | 0  | 0   |
| # genus 13 passports |   |   |   |    | 6  | 30 | 12  |
| # genus 14 passports |   |   |   |    | 3  | 0  | 0   |
| # genus 15 passports |   |   |   |    | 3  | 18 | 12  |
| # genus 16 passports |   |   |   |    |    | 6  | 6   |

# Passport counts



| degree               | 2 | 4 | 8 | 16 | 32 | 64 | 128 |
|----------------------|---|---|---|----|----|----|-----|
| # genus 17 passports |   |   |   |    |    | 39 | 25  |
| # genus 19 passports |   |   |   |    |    | 18 | 0   |
| # genus 21 passports |   |   |   |    |    | 30 | 48  |
| # genus 23 passports |   |   |   |    |    | 9  | 12  |
| # genus 24 passports |   |   |   |    |    | 3  | 0   |
| # genus 25 passports |   |   |   |    |    | 24 | 78  |
| # genus 27 passports |   |   |   |    |    | 6  | 0   |
| # genus 28 passports |   |   |   |    |    | 3  | 0   |
| # genus 29 passports |   |   |   |    |    | 6  | 30  |
| # genus 30 passports |   |   |   |    |    | 3  | 0   |
| # genus 31 passports |   |   |   |    |    | 3  | 18  |
| # genus 32 passports |   |   |   |    |    |    | 6   |
| # genus 33 passports |   |   |   |    |    |    | 117 |
| # genus 37 passports |   |   |   |    |    |    | 114 |
| # genus 39 passports |   |   |   |    |    |    | 18  |
| # genus 41 passports |   |   |   |    |    |    | 93  |

# Passport counts



| degree               | 2 | 4 | 8  | 16 | 32 | 64  | 128 |
|----------------------|---|---|----|----|----|-----|-----|
| # genus 45 passports |   |   |    |    |    |     | 48  |
| # genus 47 passports |   |   |    |    |    |     | 9   |
| # genus 48 passports |   |   |    |    |    |     | 3   |
| # genus 49 passports |   |   |    |    |    |     | 72  |
| # genus 53 passports |   |   |    |    |    |     | 26  |
| # genus 55 passports |   |   |    |    |    |     | 6   |
| # genus 56 passports |   |   |    |    |    |     | 3   |
| # genus 57 passports |   |   |    |    |    |     | 24  |
| # genus 59 passports |   |   |    |    |    |     | 6   |
| # genus 60 passports |   |   |    |    |    |     | 3   |
| # genus 61 passports |   |   |    |    |    |     | 6   |
| # genus 62 passports |   |   |    |    |    |     | 3   |
| # genus 63 passports |   |   |    |    |    |     | 3   |
| total passports      | 3 | 7 | 16 | 41 | 96 | 267 | 834 |





# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ .

# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . Given a permutation triple  $\tilde{\sigma}$  with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . Given a permutation triple  $\tilde{\sigma}$  with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to  $\tilde{\sigma}$ .

# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . Given a permutation triple  $\tilde{\sigma}$  with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to  $\tilde{\sigma}$ . Let  $X \subseteq \mathbb{A}_K^n$  with defining equations  $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$ .

# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . Given a permutation triple  $\tilde{\sigma}$  with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to  $\tilde{\sigma}$ . Let  $X \subseteq \mathbb{A}_K^n$  with defining equations  $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$ . Our goal is to find  $f \in K(X)^\times$  such that

# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . Given a permutation triple  $\tilde{\sigma}$  with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to  $\tilde{\sigma}$ . Let  $X \subseteq \mathbb{A}_K^n$  with defining equations  $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$ . Our goal is to find  $f \in K(X)^\times$  such that

$$\begin{array}{ccc} \tilde{X} & & \frac{\overline{K}(X)[y]}{(y^2-f)} \\ \downarrow \tilde{\phi} & \searrow \psi & \swarrow 2 \\ & X & \searrow d \\ & \swarrow \phi & \\ \mathbb{P}^1 & & \overline{K}(\mathbb{P}^1) \end{array}$$

# Computing 2-solvable Belyĭ maps



Let  $\phi : X \rightarrow \mathbb{P}^1$  be a Belyĭ map of degree  $d = 2^\ell$  corresponding to  $\sigma \in S_d^3$ . Given a permutation triple  $\tilde{\sigma}$  with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \langle \tilde{\sigma} \rangle \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1 ,$$

let us now consider the problem of finding the Belyĭ map corresponding to  $\tilde{\sigma}$ . Let  $X \subseteq \mathbb{A}_K^n$  with defining equations  $\{g_i\}_{i=1}^s \subset K[x_1, \dots, x_n]$ . Our goal is to find  $f \in K(X)^\times$  such that

$$\begin{array}{ccc} \tilde{X} & & \frac{\overline{K}(X)[y]}{(y^2-f)} \\ \downarrow \tilde{\phi} & \searrow \psi & \swarrow 2 \\ & X & \searrow \\ & \swarrow \phi & \overline{K}(X) \\ \mathbb{P}^1 & & \swarrow d \\ & & \overline{K}(\mathbb{P}^1) \end{array}$$

with  $\psi$  (and hence  $\tilde{\phi}$ ) satisfying the ramification conditions imposed by  $\tilde{\sigma}$ .







The procedure to find  $f \in K(X)$  is as follows:



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ .



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ . These are determined by  $\tilde{\sigma}$ .



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ . These are determined by  $\tilde{\sigma}$ .
2. Build a degree 0 divisor  $D = \sum_P n_P P$  with  $n_{Q_i}$  odd for every  $i$ .



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ . These are determined by  $\tilde{\sigma}$ .
2. Build a degree 0 divisor  $D = \sum_P n_P P$  with  $n_{Q_i}$  odd for every  $i$ .
3. Try to find  $f$  in the (computable) Riemann-Roch space  $L(D)$ .



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ . These are determined by  $\tilde{\sigma}$ .
2. Build a degree 0 divisor  $D = \sum_P n_P P$  with  $n_{Q_i}$  odd for every  $i$ .
3. Try to find  $f$  in the (computable) Riemann-Roch space  $L(D)$ .

There are (at least) two remarks to make about this process:



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ . These are determined by  $\tilde{\sigma}$ .
2. Build a degree 0 divisor  $D = \sum_P n_P P$  with  $n_{Q_i}$  odd for every  $i$ .
3. Try to find  $f$  in the (computable) Riemann-Roch space  $L(D)$ .

There are (at least) two remarks to make about this process:

- Extending the base field  $K$  may be necessary to determine  $D$ .



The procedure to find  $f \in K(X)$  is as follows:

1. Let  $\{Q_i\}$  be the points on  $X$  that we want to be ramification values of  $\psi$ . These are determined by  $\tilde{\sigma}$ .
2. Build a degree 0 divisor  $D = \sum_P n_P P$  with  $n_{Q_i}$  odd for every  $i$ .
3. Try to find  $f$  in the (computable) Riemann-Roch space  $L(D)$ .

There are (at least) two remarks to make about this process:

- ▶ Extending the base field  $K$  may be necessary to determine  $D$ .
- ▶ Class group obstruction.








$$\tilde{\sigma} = ((1432), (13)(24), (1432)), \quad \sigma = ((12), (1)(2), (12))$$



$$\tilde{\sigma} = ((1432), (13)(24), (1432)), \quad \sigma = ((12), (1)(2), (12))$$





$$X_2 : x_2^2 = x_1$$

Diagram illustrating a mapping from  $X_2$  to  $X_1$ , defined by the equation  $x_2^2 = x_1$ .



$$\tilde{\sigma} = ((1432), (13)(24), (1432)), \quad \sigma = ((12), (1)(2), (12))$$







Passport: 8T1-8,4,8-g3, size 2

Belyĭ curve:  $X : y^2 + (x^4 + 1)y = -2x^4$

Belyĭ map:  $(y + 1)^2$



Passport: 8T1-8,4,8-g3, size 2

Belyĭ curve:  $X : y^2 + (x^4 + 1)y = -2x^4$

Belyĭ map:  $(y + 1)^2$

Passport: 16T1-16,8,16-g7, size 4

Belyĭ curve:  $X : y^2 + (x^8 + 1)y = -2x^8$

Belyĭ map:  $(y + 1)^2$





# Nonhyperelliptic Belyĭ maps



$128S1-128,32,128-g62 \rightarrow 64S1-64,16,64-g30 \rightarrow$   
 $32S1-32,8,32-g14 \rightarrow 16T1-16,4,16-g6 \rightarrow 8T1-8,2,8-g2 \rightarrow$   
 $4T1-4,1,4-g0 \rightarrow 2T1-2,1,2-g0$



128S1-128,32,128-g62  $\rightarrow$  64S1-64,16,64-g30  $\rightarrow$   
32S1-32,8,32-g14  $\rightarrow$  16T1-16,4,16-g6  $\rightarrow$  8T1-8,2,8-g2  $\rightarrow$   
4T1-4,1,4-g0  $\rightarrow$  2T1-2,1,2-g0

$$\begin{aligned} X \subset \mathbb{A}^6 : & x_1^5 - x_1 - x_2^2 \\ & x_1 - x_1^3 + x_2 x_4^4 \\ & x_1^3 x_3 - x_1 x_3 - x_2 x_4^2 \\ & x_1^2 x_4^2 - x_2 x_3 + x_4^2 \\ & x_2 x_3 - x_1^2 - 1 \\ & x_3 x_4^2 - 1 \\ & x_5^2 - x_4 \\ & x_6^2 - x_5 \\ \phi : & x_3^4 x_2^2 - 2x_3^2 x_2 + 1 \end{aligned}$$



128S69-8,16,16-g49: size 4

64S7-4,8,8-g17

32S10-4,8,4-g7

16T12-4,8,2-g2

8T4-2,4,2-g0

4T2-2,2,2-g0

2T1-2,2,1-g0

# Networks?





<https://math.dartmouth.edu/~mjmusty/32.html>



<https://math.dartmouth.edu/~mjmusty/32.html>

- Is every 2-solvable Belyĭ map defined over an abelian extension of  $\mathbb{Q}$ ?



<https://math.dartmouth.edu/~mjmusty/32.html>

- ▶ Is every 2-solvable Belyĭ map defined over an abelian extension of  $\mathbb{Q}$ ?
- ▶ What can we say in the hyperelliptic case?



<https://math.dartmouth.edu/~mjmusty/32.html>

- ▶ Is every 2-solvable Belyĭ map defined over an abelian extension of  $\mathbb{Q}$ ?
- ▶ What can we say in the hyperelliptic case?
- ▶ What infinite families of 2-groups appear as monodromy groups of Belyĭ maps?





Thanks to the following for helpful discussions:

- ▶ Sam Schiavone
- ▶ Jeroen Sijsling
- ▶ John Voight



Thanks to the following for helpful discussions:

- ▶ Sam Schiavone
- ▶ Jeroen Sijsling
- ▶ John Voight

Thanks for listening!



Thanks to the following for helpful discussions:

- ▶ Sam Schiavone
- ▶ Jeroen Sijsling
- ▶ John Voight

Thanks for listening! (unless there is extra time...)

# Applications?



# Applications?



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ .

# Applications?



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ .

# Applications?



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2.



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ .





Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model.



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ .



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ .



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ . (Baker 1893).



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ . (Baker 1893). In general one must compute the adjoint ideal.



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ . (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology.



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ . (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology. Integrating yields a  $g \times 2g$  period matrix  $\Pi$





Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \text{ and } i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ . (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology. Integrating yields a  $g \times 2g$  period matrix  $\Pi$  with  $\Lambda$  the  $\mathbb{Z}$ -span of the columns of  $\Pi$ .



Let  $X$  be a 2-solvable Belyĭ curve of degree  $d$  and genus  $g$  defined over  $K$ . We would like to compute the 2-torsion field of the jacobian  $J$  of  $X$ . The field  $K(J[2])/K$  will be ramified only at 2. To start, we must compute a period matrix for  $J$ . We can embed  $X$  in  $\mathbb{P}^2$  with a singular model. Now consider an affine patch  $f(x, y) = 0$ . The space of holomorphic 1-forms on  $X$  is a subspace of

$$\left\{ \frac{x^i y^j dx}{\partial_y f(x, y)} : 0 \leq i, j \quad \text{and} \quad i + j \leq d - 3 \right\}$$

In some cases the exact space is given by  $(i, j)$  where  $(i + 1, j + 1)$  is an interior point of the Newton polygon of  $f(x, y)$ . (Baker 1893). In general one must compute the adjoint ideal. The next piece we need is a basis in homology. Integrating yields a  $g \times 2g$  period matrix  $\Pi$  with  $\Lambda$  the  $\mathbb{Z}$ -span of the columns of  $\Pi$ .  $J$  is identified with  $\mathbb{C}^g / \Lambda$ .

# Applications?





The next piece is the Abel-Jacobi map



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left( \int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left( \int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for  $t \in \frac{1}{2}\Lambda/\Lambda$ ,



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left( \int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for  $t \in \frac{1}{2}\Lambda/\Lambda$ , our task is to find  $\{Q_1, \dots, Q_g\}$  with  $Q_j \in X(\overline{K})$  such that



The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left( \int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for  $t \in \frac{1}{2}\Lambda/\Lambda$ , our task is to find  $\{Q_1, \dots, Q_g\}$  with  $Q_j \in X(\overline{K})$  such that

$$\sum_{j=1}^g \left( \int_{P_0}^{Q_j} \omega_i \right)_{i=1, \dots, g}$$





The next piece is the Abel-Jacobi map

$$\text{AJ} : X \rightarrow \mathbb{C}^g / \Lambda$$

$$P \mapsto \left( \int_{P_0}^P \omega_j \right)_{j=1, \dots, g}$$

Now for  $t \in \frac{1}{2}\Lambda/\Lambda$ , our task is to find  $\{Q_1, \dots, Q_g\}$  with  $Q_j \in X(\overline{K})$  such that

$$\sum_{j=1}^g \left( \int_{P_0}^{Q_j} \omega_i \right)_{i=1, \dots, g}$$

The coordinates of the  $Q_j$  generate the field  $K(J[2])$ .