

2-group Belyi maps

Michael Musty

July 9, 2019

Dartmouth College

Acknowledgements

- Dave, Tom, Carl, and John
- Sam, Anna, Jeroen, Edgar, Florian, and Richard
- Mary, Jim, Matt, and Nicole

Outline

Motivation

Background

Computing permutation triples

Computing equations

A refined conjecture

Examples

Motivation



Galois representations

Let X be an irreducible, smooth projective algebraic curve of genus $g \geq 1$ over a number field K . Let $G_K := \text{Gal}(K^{\text{al}} | K)$ be the absolute Galois group of K and let $\ell \in \mathbb{Z}$ be prime.

Galois representations

Let X be an irreducible, smooth projective algebraic curve of genus $g \geq 1$ over a number field K . Let $G_K := \text{Gal}(K^{\text{al}} | K)$ be the absolute Galois group of K and let $\ell \in \mathbb{Z}$ be prime.

Let $J := \text{Jac}(X)$ be the **Jacobian variety** of X . J is an abelian variety of dimension g .

Galois representations

Let X be an irreducible, smooth projective algebraic curve of genus $g \geq 1$ over a number field K . Let $G_K := \text{Gal}(K^{\text{al}} | K)$ be the absolute Galois group of K and let $\ell \in \mathbb{Z}$ be prime.

Let $J := \text{Jac}(X)$ be the **Jacobian variety** of X . J is an abelian variety of dimension g .

G_K acts on the ℓ -torsion points $J[\ell](K^{\text{al}}) \cong (\mathbb{Z}/\ell\mathbb{Z})^{2g}$ of X .

Galois representations

Let X be an irreducible, smooth projective algebraic curve of genus $g \geq 1$ over a number field K . Let $G_K := \text{Gal}(K^{\text{al}} | K)$ be the absolute Galois group of K and let $\ell \in \mathbb{Z}$ be prime.

Let $J := \text{Jac}(X)$ be the **Jacobian variety** of X . J is an abelian variety of dimension g .

G_K acts on the ℓ -torsion points $J[\ell](K^{\text{al}}) \cong (\mathbb{Z}/\ell\mathbb{Z})^{2g}$ of X .

This action determines a **mod- ℓ Galois representation**

$$\rho: G_K \rightarrow \text{Aut}(J[\ell]) \cong \text{GL}_{2g}(\mathbb{Z}/\ell\mathbb{Z}).$$

Galois representations

Let X be an irreducible, smooth projective algebraic curve of genus $g \geq 1$ over a number field K . Let $G_K := \text{Gal}(K^{\text{al}} | K)$ be the absolute Galois group of K and let $\ell \in \mathbb{Z}$ be prime.

Let $J := \text{Jac}(X)$ be the **Jacobian variety** of X . J is an abelian variety of dimension g .

G_K acts on the ℓ -torsion points $J[\ell](K^{\text{al}}) \cong (\mathbb{Z}/\ell\mathbb{Z})^{2g}$ of X .

This action determines a **mod- ℓ Galois representation**

$$\rho: G_K \rightarrow \text{Aut}(J[\ell]) \cong \text{GL}_{2g}(\mathbb{Z}/\ell\mathbb{Z}).$$

The geometry of X and the arithmetic of ρ are intimately related.

Galois representations

Let X be an irreducible, smooth projective algebraic curve of genus $g \geq 1$ over a number field K . Let $G_K := \text{Gal}(K^{\text{al}} | K)$ be the absolute Galois group of K and let $\ell \in \mathbb{Z}$ be prime.

Let $J := \text{Jac}(X)$ be the **Jacobian variety** of X . J is an abelian variety of dimension g .

G_K acts on the ℓ -torsion points $J[\ell](K^{\text{al}}) \cong (\mathbb{Z}/\ell\mathbb{Z})^{2g}$ of X .

This action determines a **mod- ℓ Galois representation**

$$\rho: G_K \rightarrow \text{Aut}(J[\ell]) \cong \text{GL}_{2g}(\mathbb{Z}/\ell\mathbb{Z}).$$

The geometry of X and the arithmetic of ρ are intimately related. For example, if X has good reduction at a prime \mathfrak{p} above $p \neq \ell$, then \mathfrak{p} will be unramified in the **ℓ -torsion field** $K(J[\ell])$.

Belyi's theorem

A **Belyi map** is a morphism $\phi: X \rightarrow \mathbb{P}^1$ of smooth projective algebraic curves over \mathbb{C} that is unramified outside of $\{0, 1, \infty\}$.

A **Belyi map** is a morphism $\phi: X \rightarrow \mathbb{P}^1$ of smooth projective algebraic curves over \mathbb{C} that is unramified outside of $\{0, 1, \infty\}$.

Theorem (Belyi 1979)

An algebraic curve (smooth projective) X over \mathbb{C} can be defined over a number field if and only if X admits a Belyi map.

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

The **genus** of ϕ is the genus of the curve X .

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

The **genus** of ϕ is the genus of the curve X .

The **degree** of ϕ is the degree of the field extension

$$K(\mathbb{P}^1) \hookrightarrow K(X).$$

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

The **genus** of ϕ is the genus of the curve X .

The **degree** of ϕ is the degree of the field extension

$$K(\mathbb{P}^1) \hookrightarrow K(X).$$

ϕ is **geometrically Galois** if $K^{\text{al}}(X)$ is a Galois extension.

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

The **genus** of ϕ is the genus of the curve X .

The **degree** of ϕ is the degree of the field extension

$$K(\mathbb{P}^1) \hookrightarrow K(X).$$

ϕ is **geometrically Galois** if $K^{\text{al}}(X)$ is a Galois extension.

The **monodromy group** of ϕ , $\text{Mon}(\phi)$, is the image of the map

$$\pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \star) \rightarrow S_d$$

obtained by path lifting.

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

The **genus** of ϕ is the genus of the curve X .

The **degree** of ϕ is the degree of the field extension

$$K(\mathbb{P}^1) \hookrightarrow K(X).$$

ϕ is **geometrically Galois** if $K^{\text{al}}(X)$ is a Galois extension.

The **monodromy group of ϕ** , $\text{Mon}(\phi)$, is the image of the map

$$\pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \star) \rightarrow S_d$$

obtained by path lifting.

When ϕ is Galois we can identify $\text{Mon}(\phi)$ with $\text{Gal}(K^{\text{al}}(X) | K^{\text{al}}(\mathbb{P}^1))$.

Some definitions

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map defined over K .

The **genus** of ϕ is the genus of the curve X .

The **degree** of ϕ is the degree of the field extension

$$K(\mathbb{P}^1) \hookrightarrow K(X).$$

ϕ is **geometrically Galois** if $K^{\text{al}}(X)$ is a Galois extension.

The **monodromy group** of ϕ , $\text{Mon}(\phi)$, is the image of the map

$$\pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \star) \rightarrow S_d$$

obtained by path lifting.

When ϕ is Galois we can identify $\text{Mon}(\phi)$ with $\text{Gal}(K^{\text{al}}(X) | K^{\text{al}}(\mathbb{P}^1))$.

We can now state Beckmann's theorem to relate Belyi maps to our original motivation.

Theorem (Beckmann 1989)

*Let $\phi: X \rightarrow \mathbb{P}^1$ be a Galois Belyi map with monodromy group G .
Let p be a prime not dividing $\#G$.*

Theorem (Beckmann 1989)

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Galois Belyi map with monodromy group G .

Let p be a prime not dividing $\#G$.

Then there exists a number field M satisfying the following properties.

Theorem (Beckmann 1989)

*Let $\phi: X \rightarrow \mathbb{P}^1$ be a Galois Belyi map with monodromy group G .
Let p be a prime not dividing $\#G$.*

Then there exists a number field M satisfying the following properties.

- *p is unramified in M*
- *ϕ is defined over M*
- *X is defined over M*
- *X has good reduction at all primes \mathfrak{p} of M above p*

Why $p = 2$?

Conjecture (Gross 1998)

For every prime p , there exists a nonsolvable Galois number field ramified only at p .

Why $p = 2$?

Conjecture (Gross 1998)

For every prime p , there exists a nonsolvable Galois number field ramified only at p .

$p \geq 11$: existence (Serre), explicit (Edixhoven, Mascot)

$p = 7$: existence (Dieulefait)

$p = 5$: existence (Dembélé, Greenberg, Voight), explicit (Roberts)

$p = 3$: existence (Dembélé, Greenberg, Voight)

$p = 2$: existence (Dembélé)

Why $p = 2$?

Conjecture (Gross 1998)

For every prime p , there exists a nonsolvable Galois number field ramified only at p .

$p \geq 11$: existence (Serre), explicit (Edixhoven, Mascot)

$p = 7$: existence (Dieulefait)

$p = 5$: existence (Dembélé, Greenberg, Voight), explicit (Roberts)

$p = 3$: existence (Dembélé, Greenberg, Voight)

$p = 2$: existence (Dembélé)

The hope is that an explicit nonsolvable field ramified only at 2 can be obtained as $K(\text{Jac}(X)[2])$ where X is the domain of a Galois Belyi map with monodromy group a 2-group.

Why $p = 2$?

Conjecture (Gross 1998)

For every prime p , there exists a nonsolvable Galois number field ramified only at p .

$p \geq 11$: existence (Serre), explicit (Edixhoven, Mascot)

$p = 7$: existence (Dieulefait)

$p = 5$: existence (Dembélé, Greenberg, Voight), explicit (Roberts)

$p = 3$: existence (Dembélé, Greenberg, Voight)

$p = 2$: existence (Dembélé)

The hope is that an explicit nonsolvable field ramified only at 2 can be obtained as $K(\text{Jac}(X)[2])$ where X is the domain of a Galois Belyi map with monodromy group a 2-group.

We call these Belyi maps **2-group Belyi maps**.

Main results

Motivated by the applications of 2-group Belyi maps to arithmetic geometry, we now state the main results.

Main results

Motivated by the applications of 2-group Belyi maps to arithmetic geometry, we now state the main results.

- implementation of an algorithm to enumerate isomorphism classes of 2-group Belyi maps

Main results

Motivated by the applications of 2-group Belyi maps to arithmetic geometry, we now state the main results.

- implementation of an algorithm to enumerate isomorphism classes of 2-group Belyi maps
- implementation of an algorithm to compute equations for 2-group Belyi maps over finite fields

Main results

Motivated by the applications of 2-group Belyi maps to arithmetic geometry, we now state the main results.



- implementation of an algorithm to enumerate isomorphism classes of 2-group Belyi maps
- implementation of an algorithm to compute equations for 2-group Belyi maps over finite fields
- implementation of a *method* to compute equations for 2-group Belyi maps over number fields

Main results

Motivated by the applications of 2-group Belyi maps to arithmetic geometry, we now state the main results.

- implementation of an algorithm to enumerate isomorphism classes of 2-group Belyi maps
- implementation of an algorithm to compute equations for 2-group Belyi maps over finite fields
- implementation of a *method* to compute equations for 2-group Belyi maps over number fields
- computational and theoretical evidence supporting a conjecture that every 2-group Belyi map is defined over an abelian extension of the rationals

2-group Belyi maps as iterated quadratic extensions



Background



Isomorphism of Belyi maps

Let $\phi: X \rightarrow \mathbb{P}^1$ and $\phi': X' \rightarrow \mathbb{P}^1$ be Belyi maps of degree d .

Isomorphism of Belyi maps

Let $\phi: X \rightarrow \mathbb{P}^1$ and $\phi': X' \rightarrow \mathbb{P}^1$ be Belyi maps of degree d . ϕ and ϕ' are **isomorphic** (respectively **lax isomorphic**) if the diagrams

$$\begin{array}{ccc} X & \xrightarrow{\sim} & X' \\ \phi \searrow & & \swarrow \phi' \\ & \mathbb{P}^1 & \end{array}, \text{ respectively } \begin{array}{ccc} X & \xrightarrow{\sim} & X' \\ \phi \downarrow & & \downarrow \phi' \\ \mathbb{P}^1 & \xrightarrow[\beta]{\sim} & \mathbb{P}^1 \end{array}$$

commute where $\beta(\{0, 1, \infty\}) = \{0, 1, \infty\}$.

Permutation Triples

A **transitive permutation triple of degree d** is a triple

$$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$$

such that

- $\sigma_\infty \sigma_1 \sigma_0 = 1$
- σ generates a transitive subgroup of S_d

Permutation Triples

A **transitive permutation triple of degree d** is a triple

$$\sigma = (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$$

such that

- $\sigma_\infty \sigma_1 \sigma_0 = 1$
- σ generates a transitive subgroup of S_d

The set of degree d Belyi maps up to isomorphism is in bijection with the set of degree d transitive permutation triples up to **simultaneous conjugation** and the group $\langle \sigma \rangle$ is the monodromy group of ϕ .

Passports

A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

Passports

A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

The **passport of a Belyi map** $\phi : X \rightarrow \mathbb{P}^1$ is $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$ with $g(X)$ the genus of X , $\text{Mon}(\phi)$ the monodromy group of ϕ , and the partitions from ramification.

Passports

A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

The **passport of a Belyi map** $\phi : X \rightarrow \mathbb{P}^1$ is $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$ with $g(X)$ the genus of X , $\text{Mon}(\phi)$ the monodromy group of ϕ , and the partitions from ramification.

The **passport of a permutation triple** σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$ where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \#\text{cycles of } \tau,$$

and $\lambda(\sigma)$ is specified by cycle structures.

Passports

A **passport** \mathcal{P} consists of the data (g, G, λ) where $g \geq 0$ is an integer, $G \leq S_d$ is a transitive subgroup, and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d .

The **passport of a Belyi map** $\phi : X \rightarrow \mathbb{P}^1$ is $(g(X), \text{Mon}(\phi), (\lambda_0, \lambda_1, \lambda_\infty))$ with $g(X)$ the genus of X , $\text{Mon}(\phi)$ the monodromy group of ϕ , and the partitions from ramification.

The **passport of a permutation triple** σ is $(g(\sigma), \langle \sigma \rangle, \lambda(\sigma))$ where

$$g(\sigma) = 1 - d + (e(\sigma_0) - e(\sigma_1) - e(\sigma_\infty))/2$$

with

$$e(\tau) = d - \#\text{cycles of } \tau,$$

and $\lambda(\sigma)$ is specified by cycle structures.

We now discuss the importance of organizing triples by passport. 12/36

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

A number field K is a **field of definition** of X if X can be defined by polynomial equations over K . Similarly for ϕ .

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

A number field K is a **field of definition** of X if X can be defined by polynomial equations over K . Similarly for ϕ .

The **field of moduli** $M(X)$ of X is the fixed field of the field automorphisms $\{\tau \in \text{Aut}(\mathbb{C}) : \tau(X) \simeq X\}$. Similarly for ϕ , $M(\phi)$.

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

A number field K is a **field of definition** of X if X can be defined by polynomial equations over K . Similarly for ϕ .

The **field of moduli** $M(X)$ of X is the fixed field of the field automorphisms $\{\tau \in \text{Aut}(\mathbb{C}) : \tau(X) \simeq X\}$. Similarly for ϕ , $M(\phi)$.

The **size** of a passport \mathcal{P} is the number of simultaneous conjugacy classes of permutation triples σ with passport \mathcal{P} .

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

A number field K is a **field of definition** of X if X can be defined by polynomial equations over K . Similarly for ϕ .

The **field of moduli** $M(X)$ of X is the fixed field of the field automorphisms $\{\tau \in \text{Aut}(\mathbb{C}) : \tau(X) \simeq X\}$. Similarly for ϕ , $M(\phi)$.

The **size** of a passport \mathcal{P} is the number of simultaneous conjugacy classes of permutation triples σ with passport \mathcal{P} .

Theorem

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map with passport \mathcal{P} . Then the degree of $M(\phi)$ is bounded by the size of \mathcal{P} .

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

A number field K is a **field of definition** of X if X can be defined by polynomial equations over K . Similarly for ϕ .

The **field of moduli** $M(X)$ of X is the fixed field of the field automorphisms $\{\tau \in \text{Aut}(\mathbb{C}) : \tau(X) \simeq X\}$. Similarly for ϕ , $M(\phi)$.

The **size** of a passport \mathcal{P} is the number of simultaneous conjugacy classes of permutation triples σ with passport \mathcal{P} .

Theorem

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map with passport \mathcal{P} . Then the degree of $M(\phi)$ is bounded by the size of \mathcal{P} .

A Belyi map need not be defined over its field of moduli \mathbb{Q}^{mod}

Fields of moduli, fields of definition, and passports

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map.

A number field K is a **field of definition** of X if X can be defined by polynomial equations over K . Similarly for ϕ .

The **field of moduli** $M(X)$ of X is the fixed field of the field automorphisms $\{\tau \in \text{Aut}(\mathbb{C}) : \tau(X) \simeq X\}$. Similarly for ϕ , $M(\phi)$.

The **size** of a passport \mathcal{P} is the number of simultaneous conjugacy classes of permutation triples σ with passport \mathcal{P} .

Theorem

Let $\phi: X \rightarrow \mathbb{P}^1$ be a Belyi map with passport \mathcal{P} . Then the degree of $M(\phi)$ is bounded by the size of \mathcal{P} .

A Belyi map need not be defined over its field of moduli \mathbb{Q}^{tr}

The situation improves, however, in the Galois setting...

The Galois setting

Let $\phi: X \rightarrow \mathbb{P}^1$ be a *Galois* Belyi map of degree d with monodromy group G corresponding to the permutation triple σ .

The Galois setting

Let $\phi: X \rightarrow \mathbb{P}^1$ be a *Galois* Belyi map of degree d with monodromy group G corresponding to the permutation triple σ .

Then

- ϕ and X are defined over $M(\phi)$,
- $\#G = d$,
- all cycles of σ_s have the same length for $s \in \{0, 1, \infty\}$,
- and if we let a, b, c be the orders of $\sigma_0, \sigma_1, \sigma_\infty$ respectively, we have

$$g(X) = 1 + \frac{\#G}{2} \left(1 - \frac{1}{a} - \frac{1}{b} - \frac{1}{c} \right).$$

Function fields

Let $\phi: X \rightarrow \mathbb{P}^1$ be a degree d Belyi map defined over K .

Function fields

Let $\phi: X \rightarrow \mathbb{P}^1$ be a degree d Belyi map defined over K .

This corresponds to a degree d extension $K(\mathbb{P}^1) \hookrightarrow K(X)$.

Function fields

Let $\phi: X \rightarrow \mathbb{P}^1$ be a degree d Belyi map defined over K .

This corresponds to a degree d extension $K(\mathbb{P}^1) \hookrightarrow K(X)$.

$K(\mathbb{P}^1) \cong K(x)$ the **rational function field** of K in one variable.

Function fields

Let $\phi: X \rightarrow \mathbb{P}^1$ be a degree d Belyi map defined over K .

This corresponds to a degree d extension $K(\mathbb{P}^1) \hookrightarrow K(X)$.

$K(\mathbb{P}^1) \cong K(x)$ the **rational function field** of K in one variable.

For K perfect, $K(X)$ has a primitive element and can be written as

$$K(x)(\alpha) = \frac{K(x)[y]}{(\text{minpoly}_{\alpha, K(x)}(y))}$$

Function fields

Let $\phi: X \rightarrow \mathbb{P}^1$ be a degree d Belyi map defined over K .

This corresponds to a degree d extension $K(\mathbb{P}^1) \hookrightarrow K(X)$.

$K(\mathbb{P}^1) \cong K(x)$ the **rational function field** of K in one variable.

For K perfect, $K(X)$ has a primitive element and can be written as

$$K(x)(\alpha) = \frac{K(x)[y]}{(\text{minpoly}_{\alpha, K(x)}(y))}$$

Ramification in this setting corresponds to the factorization of ideals (x) , $(x-1)$, $(1/x)$ in maximal orders of $K(X)$.

Function fields

Let $\phi: X \rightarrow \mathbb{P}^1$ be a degree d Belyi map defined over K .

This corresponds to a degree d extension $K(\mathbb{P}^1) \hookrightarrow K(X)$.

$K(\mathbb{P}^1) \cong K(x)$ the **rational function field** of K in one variable.

For K perfect, $K(X)$ has a primitive element and can be written as

$$K(x)(\alpha) = \frac{K(x)[y]}{(\text{minpoly}_{\alpha, K(x)}(y))}$$

Ramification in this setting corresponds to the factorization of ideals (x) , $(x-1)$, $(1/x)$ in maximal orders of $K(X)$.

The monodromy group in this setting corresponds to field automorphisms of the Galois closure of $K(X)$ fixing $K(x)$.

Computing permutation triples



Setup

We first define some terminology for permutation triples corresponding to 2-group Belyi maps.

Setup

We first define some terminology for permutation triples corresponding to 2-group Belyi maps.

A **2-group permutation triple** of degree $d \in \mathbb{Z}_{\geq 1}$ is a triple of permutations $\sigma := (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ satisfying

- $\sigma_\infty \sigma_1 \sigma_0 = \text{id}$;
- $G := \langle \sigma_0, \sigma_1 \rangle$ is a transitive subgroup of S_d ; and
- G is a 2-group of order d embedded in S_d via its left regular representation.

Setup

We first define some terminology for permutation triples corresponding to 2-group Belyi maps.

A **2-group permutation triple** of degree $d \in \mathbb{Z}_{\geq 1}$ is a triple of permutations $\sigma := (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ satisfying

- $\sigma_\infty \sigma_1 \sigma_0 = \text{id}$;
- $G := \langle \sigma_0, \sigma_1 \rangle$ is a transitive subgroup of S_d ; and
- G is a 2-group of order d embedded in S_d via its left regular representation.

G is called the **monodromy group** of σ .

Setup

We first define some terminology for permutation triples corresponding to 2-group Belyi maps.

A **2-group permutation triple** of degree $d \in \mathbb{Z}_{\geq 1}$ is a triple of permutations $\sigma := (\sigma_0, \sigma_1, \sigma_\infty) \in S_d^3$ satisfying

- $\sigma_\infty \sigma_1 \sigma_0 = \text{id}$;
- $G := \langle \sigma_0, \sigma_1 \rangle$ is a transitive subgroup of S_d ; and
- G is a 2-group of order d embedded in S_d via its left regular representation.

G is called the **monodromy group** of σ .

We say two degree d 2-group permutation triples σ, σ' are **simultaneously conjugate** if there exists $\tau \in S_d$ such that

$$\sigma^\tau := (\tau^{-1} \sigma_0 \tau, \tau^{-1} \sigma_1 \tau, \tau^{-1} \sigma_\infty \tau) = \sigma'$$

Lifting permutation triples

Let σ be a 2-group permutation triple.

Lifting permutation triples

Let σ be a 2-group permutation triple.

A **lift** of σ is a 2-group permutation triple $\tilde{\sigma} \in S_{2d}^3$ such that $\langle \tilde{\sigma} \rangle$ is isomorphic to some extension \tilde{G} of $\mathbb{Z}/2\mathbb{Z}$ by G as in the exact sequence below.

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1$$

Lifting permutation triples

Let σ be a 2-group permutation triple.

A **lift** of σ is a 2-group permutation triple $\tilde{\sigma} \in S_{2d}^3$ such that $\langle \tilde{\sigma} \rangle$ is isomorphic to some extension \tilde{G} of $\mathbb{Z}/2\mathbb{Z}$ by G as in the exact sequence below.

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota} \tilde{G} \xrightarrow{\pi} \langle \sigma \rangle \longrightarrow 1$$

For a 2-group permutation triple σ , we denote the set of lifts of σ by $\text{Lifts}(\sigma)$ and $\text{Lifts}(\sigma)/\sim$ denotes the set of lifts up to simultaneous conjugation.

Algorithm to compute $\text{Lifts}(\sigma)/\sim$

Input: σ a 2-group permutation triple of degree d

Output: $\text{Lifts}(\sigma)/\sim$

Algorithm to compute $\text{Lifts}(\sigma)/\sim$

Input: σ a 2-group permutation triple of degree d

Output: $\text{Lifts}(\sigma)/\sim$

1. Let $G = \langle \sigma \rangle$ and compute representatives of $H^2(G, A)$ where $A := \mathbb{Z}/2\mathbb{Z}$ with the trivial G -module structure

Algorithm to compute $\text{Lifts}(\sigma)/\sim$

Input: σ a 2-group permutation triple of degree d

Output: $\text{Lifts}(\sigma)/\sim$

1. Let $G = \langle \sigma \rangle$ and compute representatives of $H^2(G, A)$ where $A := \mathbb{Z}/2\mathbb{Z}$ with the trivial G -module structure
2. For each $f \in H^2(G, A)$ compute the corresponding extension

$$1 \longrightarrow A \xrightarrow{\iota_f} \tilde{G}_f \xrightarrow{\pi_f} G \longrightarrow 1$$

Algorithm to compute $\text{Lifts}(\sigma)/\sim$

Input: σ a 2-group permutation triple of degree d

Output: $\text{Lifts}(\sigma)/\sim$

1. Let $G = \langle \sigma \rangle$ and compute representatives of $H^2(G, A)$ where $A := \mathbb{Z}/2\mathbb{Z}$ with the trivial G -module structure
2. For each $f \in H^2(G, A)$ compute the corresponding extension

$$1 \longrightarrow A \xrightarrow{\iota_f} \tilde{G}_f \xrightarrow{\pi_f} G \longrightarrow 1$$

3. For each extension \tilde{G}_f compute the set $\text{Lifts}(\sigma, f)$ defined by
$$\left\{ \tilde{\sigma} : \tilde{\sigma}_s \in \pi_f^{-1}(\sigma_s) \text{ for } s \in \{0, 1, \infty\}, \tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1, \langle \tilde{\sigma} \rangle = \tilde{G}_f \right\}$$

Algorithm to compute $\text{Lifts}(\sigma)/\sim$

Input: σ a 2-group permutation triple of degree d

Output: $\text{Lifts}(\sigma)/\sim$

1. Let $G = \langle \sigma \rangle$ and compute representatives of $H^2(G, A)$ where $A := \mathbb{Z}/2\mathbb{Z}$ with the trivial G -module structure
2. For each $f \in H^2(G, A)$ compute the corresponding extension

$$1 \longrightarrow A \xrightarrow{\iota_f} \tilde{G}_f \xrightarrow{\pi_f} G \longrightarrow 1$$

3. For each extension \tilde{G}_f compute the set $\text{Lifts}(\sigma, f)$ defined by
$$\left\{ \tilde{\sigma} : \tilde{\sigma}_s \in \pi_f^{-1}(\sigma_s) \text{ for } s \in \{0, 1, \infty\}, \tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1, \langle \tilde{\sigma} \rangle = \tilde{G}_f \right\}$$
- 4.

$$\text{Lifts}(\sigma) := \bigcup_{f \in H^2(G, A)} \text{Lifts}(\sigma, f)$$

Algorithm to compute $\text{Lifts}(\sigma)/\sim$

Input: σ a 2-group permutation triple of degree d

Output: $\text{Lifts}(\sigma)/\sim$

1. Let $G = \langle \sigma \rangle$ and compute representatives of $H^2(G, A)$ where $A := \mathbb{Z}/2\mathbb{Z}$ with the trivial G -module structure
2. For each $f \in H^2(G, A)$ compute the corresponding extension

$$1 \longrightarrow A \xrightarrow{\iota_f} \tilde{G}_f \xrightarrow{\pi_f} G \longrightarrow 1$$

3. For each extension \tilde{G}_f compute the set $\text{Lifts}(\sigma, f)$ defined by
$$\left\{ \tilde{\sigma} : \tilde{\sigma}_s \in \pi_f^{-1}(\sigma_s) \text{ for } s \in \{0, 1, \infty\}, \tilde{\sigma}_\infty \tilde{\sigma}_1 \tilde{\sigma}_0 = 1, \langle \tilde{\sigma} \rangle = \tilde{G}_f \right\}$$
- 4.

$$\text{Lifts}(\sigma) := \bigcup_{f \in H^2(G, A)} \text{Lifts}(\sigma, f)$$

5. Quotient $\text{Lifts}(\sigma)$ by simultaneous conjugation

Example computing $\text{Lifts}(\sigma)/\sim$: setup

Let $\sigma = ((1\ 2), \text{id}, (1\ 2))$. Then $G = \langle \sigma \rangle = \mathbb{Z}/2\mathbb{Z}$.

Example computing $\text{Lifts}(\sigma)/\sim$: setup

Let $\sigma = ((1\ 2), \text{id}, (1\ 2))$. Then $G = \langle \sigma \rangle = \mathbb{Z}/2\mathbb{Z}$.

$\tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_1} \tilde{G}_1 \xrightarrow{\pi_1} G \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_2} \tilde{G}_2 \xrightarrow{\pi_2} G \longrightarrow 1$$

Example computing $\text{Lifts}(\sigma)/\sim$: setup

Let $\sigma = ((1\ 2), \text{id}, (1\ 2))$. Then $G = \langle \sigma \rangle = \mathbb{Z}/2\mathbb{Z}$.

$\tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_1} \tilde{G}_1 \xrightarrow{\pi_1} G \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_2} \tilde{G}_2 \xrightarrow{\pi_2} G \longrightarrow 1$$

Each map π_1, π_2 pulls back to 4 triples that multiply to id:

Example computing $\text{Lifts}(\sigma)/\sim$: setup

Let $\sigma = ((1\ 2), \text{id}, (1\ 2))$. Then $G = \langle \sigma \rangle = \mathbb{Z}/2\mathbb{Z}$.

$\tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_1} \tilde{G}_1 \xrightarrow{\pi_1} G \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_2} \tilde{G}_2 \xrightarrow{\pi_2} G \longrightarrow 1$$

Each map π_1, π_2 pulls back to 4 triples that multiply to id:

$$T_1 = \left\{ ((1\ 2)(3\ 4), \text{id}, (1\ 2)(3\ 4)), ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)), \right. \\ \left. ((1\ 4)(2\ 3), \text{id}, (1\ 4)(2\ 3)), ((1\ 4)(2\ 3), (1\ 3)(2\ 4), (1\ 2)(3\ 4)) \right\}$$

Example computing $\text{Lifts}(\sigma)/\sim$: setup

Let $\sigma = ((1\ 2), \text{id}, (1\ 2))$. Then $G = \langle \sigma \rangle = \mathbb{Z}/2\mathbb{Z}$.

$\tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}$ with

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_1} \tilde{G}_1 \xrightarrow{\pi_1} G \longrightarrow 1$$

$$1 \longrightarrow \mathbb{Z}/2\mathbb{Z} \xrightarrow{\iota_2} \tilde{G}_2 \xrightarrow{\pi_2} G \longrightarrow 1$$

Each map π_1, π_2 pulls back to 4 triples that multiply to id:

$$T_1 = \left\{ ((1\ 2)(3\ 4), \text{id}, (1\ 2)(3\ 4)), ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)), \right. \\ \left. ((1\ 4)(2\ 3), \text{id}, (1\ 4)(2\ 3)), ((1\ 4)(2\ 3), (1\ 3)(2\ 4), (1\ 2)(3\ 4)) \right\}$$

$$T_2 = \left\{ ((1\ 4\ 3\ 2), \text{id}, (1\ 2\ 3\ 4)), ((1\ 2\ 3\ 4), (1\ 3)(2\ 4), (1\ 2\ 3\ 4)), \right. \\ \left. ((1\ 2\ 3\ 4), \text{id}, (1\ 4\ 3\ 2)), ((1\ 4\ 3\ 2), (1\ 3)(2\ 4), (1\ 4\ 3\ 2)) \right\}$$

Example computing $\text{Lifts}(\sigma)/\sim$: action on blocks

Choose $\alpha = (1\ 3)(2\ 4)$ to be the generator of $\iota_1(\mathbb{Z}/2\mathbb{Z})$ in \tilde{G}_1 .

Example computing $\text{Lifts}(\sigma)/\sim$: action on blocks

Choose $\alpha = (1\,3)(2\,4)$ to be the generator of $\iota_1(\mathbb{Z}/2\mathbb{Z})$ in \tilde{G}_1 .

Each triple in T_1 must act on the *blocks* $\{\boxed{1\,3}, \boxed{2\,4}\}$ corresponding to the permutations in σ .

Example computing $\text{Lifts}(\sigma)/\sim$: action on blocks

Choose $\alpha = (1\ 3)(2\ 4)$ to be the generator of $\iota_1(\mathbb{Z}/2\mathbb{Z})$ in \tilde{G}_1 .

Each triple in T_1 must act on the *blocks* $\{\boxed{1\ 3}, \boxed{2\ 4}\}$ corresponding to the permutations in σ .

Let $(\tilde{\sigma}_0, \tilde{\sigma}_1, \tilde{\sigma}_\infty) = ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 4))$.

Example computing $\text{Lifts}(\sigma)/\sim$: action on blocks

Choose $\alpha = (1\ 3)(2\ 4)$ to be the generator of $\iota_1(\mathbb{Z}/2\mathbb{Z})$ in \tilde{G}_1 .

Each triple in T_1 must act on the *blocks* $\{\boxed{1\ 3}, \boxed{2\ 4}\}$ corresponding to the permutations in σ .

Let $(\tilde{\sigma}_0, \tilde{\sigma}_1, \tilde{\sigma}_\infty) = ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 4))$.

Note that $\tilde{\sigma}_0(\boxed{1\ 3}) = \boxed{2\ 4}$ and $\tilde{\sigma}_0(\boxed{2\ 4}) = \boxed{1\ 3}$.

Example computing $\text{Lifts}(\sigma)/\sim$: action on blocks

Choose $\alpha = (13)(24)$ to be the generator of $\iota_1(\mathbb{Z}/2\mathbb{Z})$ in \tilde{G}_1 .

Each triple in T_1 must act on the *blocks* $\{\boxed{13}, \boxed{24}\}$ corresponding to the permutations in σ .

Let $(\tilde{\sigma}_0, \tilde{\sigma}_1, \tilde{\sigma}_\infty) = ((12)(34), (13)(24), (14)(24))$.

Note that $\tilde{\sigma}_0(\boxed{13}) = \boxed{24}$ and $\tilde{\sigma}_0(\boxed{24}) = \boxed{13}$.

The induced permutation of $\tilde{\sigma}_0$ on blocks is $(\boxed{13}, \boxed{24})$ which is the same as the permutation $\sigma_0 = (12)$

Example computing $\text{Lifts}(\sigma)/\sim$: action on blocks

Choose $\alpha = (1\ 3)(2\ 4)$ to be the generator of $\iota_1(\mathbb{Z}/2\mathbb{Z})$ in \tilde{G}_1 .

Each triple in T_1 must act on the *blocks* $\{\boxed{1\ 3}, \boxed{2\ 4}\}$ corresponding to the permutations in σ .

Let $(\tilde{\sigma}_0, \tilde{\sigma}_1, \tilde{\sigma}_\infty) = ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 4))$.

Note that $\tilde{\sigma}_0(\boxed{1\ 3}) = \boxed{2\ 4}$ and $\tilde{\sigma}_0(\boxed{2\ 4}) = \boxed{1\ 3}$.

The induced permutation of $\tilde{\sigma}_0$ on blocks is $(\boxed{1\ 3}, \boxed{2\ 4})$ which is the same as the permutation $\sigma_0 = (1\ 2)$

Choosing

$$\alpha := (1\ d+1)(2\ d+2) \dots (d-1\ 2d-1)(d\ 2d)$$

allows us to label blocks by reducing modulo d .

Example computing $\text{Lifts}(\sigma)/\sim$: conclude

We currently have triples that multiply to id and have the correct action on blocks, but we only want triples that generate the correct group.

Example computing $\text{Lifts}(\sigma)/\sim$: conclude

We currently have triples that multiply to id and have the correct action on blocks, but we only want triples that generate the correct group.

$$\text{Lifts}(\sigma, \tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}) = \left\{ ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)), ((1\ 4)(2\ 3), (1\ 3)(2\ 4), (1\ 2)(3\ 4)) \right\}$$

Example computing $\text{Lifts}(\sigma)/\sim$: conclude

We currently have triples that multiply to id and have the correct action on blocks, but we only want triples that generate the correct group.

$$\text{Lifts}(\sigma, \tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}) = \left\{ ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)), ((1\ 4)(2\ 3), (1\ 3)(2\ 4), (1\ 2)(3\ 4)) \right\}$$

$$\text{Lifts}(\sigma, \tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}) = T_2 = \left\{ ((1\ 4\ 3\ 2), \text{id}, (1\ 2\ 3\ 4)), ((1\ 2\ 3\ 4), (1\ 3)(2\ 4), (1\ 2\ 3\ 4)), \right. \\ \left. ((1\ 2\ 3\ 4), \text{id}, (1\ 4\ 3\ 2)), ((1\ 4\ 3\ 2), (1\ 3)(2\ 4), (1\ 4\ 3\ 2)) \right\}$$

Example computing $\text{Lifts}(\sigma)/\sim$: conclude

We currently have triples that multiply to id and have the correct action on blocks, but we only want triples that generate the correct group.

$$\text{Lifts}(\sigma, \tilde{G}_1 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}) = \left\{ ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)), ((1\ 4)(2\ 3), (1\ 3)(2\ 4), (1\ 2)(3\ 4)) \right\}$$

$$\text{Lifts}(\sigma, \tilde{G}_2 \cong \mathbb{Z}/4\mathbb{Z}) = T_2 = \left\{ ((1\ 4\ 3\ 2), \text{id}, (1\ 2\ 3\ 4)), ((1\ 2\ 3\ 4), (1\ 3)(2\ 4), (1\ 2\ 3\ 4)), \right. \\ \left. ((1\ 2\ 3\ 4), \text{id}, (1\ 4\ 3\ 2)), ((1\ 4\ 3\ 2), (1\ 3)(2\ 4), (1\ 4\ 3\ 2)) \right\}$$

Lastly, we quotient by simultaneous conjugation to obtain

$$\text{Lifts}(\sigma)/\sim = \left\{ ((1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)), \right. \\ \left. ((1\ 4\ 3\ 2), \text{id}, (1\ 2\ 3\ 4)), ((1\ 2\ 3\ 4), (1\ 3)(2\ 4), (1\ 2\ 3\ 4)) \right\}$$

Bipartite graphs of permutation triples

Now that we can lift permutation triples, we now describe some notation for the bipartite graphs that organize these triples.

Bipartite graphs of permutation triples

Now that we can lift permutation triples, we now describe some notation for the bipartite graphs that organize these triples.

For $i \in \mathbb{Z}_{\geq 1}$ we define the bipartite graph denoted \mathcal{G}_{2^i} with the following node sets.

- $\mathcal{G}_{2^i}^{\text{above}}$: the set of isomorphism classes of 2-group Belyi maps of degree 2^i indexed by 2-group permutation triples $\tilde{\sigma}$ up to simultaneous conjugation in S_{2^i}
- $\mathcal{G}_{2^i}^{\text{below}}$: the set of isomorphism classes of 2-group Belyi maps of degree 2^{i-1} indexed by 2-group permutation triples σ up to simultaneous conjugation in $S_{2^{i-1}}$

Bipartite graphs of permutation triples

Now that we can lift permutation triples, we now describe some notation for the bipartite graphs that organize these triples.

For $i \in \mathbb{Z}_{\geq 1}$ we define the bipartite graph denoted \mathcal{G}_{2^i} with the following node sets.

- $\mathcal{G}_{2^i}^{\text{above}}$: the set of isomorphism classes of 2-group Belyi maps of degree 2^i indexed by 2-group permutation triples $\tilde{\sigma}$ up to simultaneous conjugation in S_{2^i}
- $\mathcal{G}_{2^i}^{\text{below}}$: the set of isomorphism classes of 2-group Belyi maps of degree 2^{i-1} indexed by 2-group permutation triples σ up to simultaneous conjugation in $S_{2^{i-1}}$

For every pair of nodes $(\tilde{\sigma}, \sigma) \in \mathcal{G}_{2^i}^{\text{above}} \times \mathcal{G}_{2^i}^{\text{below}}$ there is an edge between σ and $\tilde{\sigma}$ if and only if $\tilde{\sigma}$ is simultaneously conjugate to a lift of σ .

Algorithm to compute \mathcal{G}_{2^i}

Input: The bipartite graph $\mathcal{G}_{2^{i-1}}$

Output: The bipartite graph \mathcal{G}_{2^i}

Algorithm to compute \mathcal{G}_{2^i}

Input: The bipartite graph $\mathcal{G}_{2^{i-1}}$

Output: The bipartite graph \mathcal{G}_{2^i}

1.

$$\text{Lifts}(\mathcal{G}_{2^{i-1}}) := \bigcup_{\sigma \in \mathcal{G}_{2^{i-1}}^{\text{above}}} \text{Lifts}(\sigma)/\sim$$

Algorithm to compute \mathcal{G}_{2^i}

Input: The bipartite graph $\mathcal{G}_{2^{i-1}}$

Output: The bipartite graph \mathcal{G}_{2^i}

1.

$$\text{Lifts}(\mathcal{G}_{2^{i-1}}) := \bigcup_{\sigma \in \mathcal{G}_{2^{i-1}}^{\text{above}}} \text{Lifts}(\sigma)/\sim$$

2. Quotient $\text{Lifts}(\mathcal{G}_{2^{i-1}})$ by simultaneous conjugation in S_{2^i} to obtain $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$

Algorithm to compute \mathcal{G}_{2^i}

Input: The bipartite graph $\mathcal{G}_{2^{i-1}}$

Output: The bipartite graph \mathcal{G}_{2^i}

1.

$$\text{Lifts}(\mathcal{G}_{2^{i-1}}) := \bigcup_{\sigma \in \mathcal{G}_{2^{i-1}}^{\text{above}}} \text{Lifts}(\sigma)/\sim$$

2. Quotient $\text{Lifts}(\mathcal{G}_{2^{i-1}})$ by simultaneous conjugation in S_{2^i} to obtain $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$
3. Define $\mathcal{G}_{2^i}^{\text{below}} := \mathcal{G}_{2^{i-1}}^{\text{above}}$ and define $\mathcal{G}_{2^i}^{\text{above}}$ by representatives of $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$

Algorithm to compute \mathcal{G}_{2^i}

Input: The bipartite graph $\mathcal{G}_{2^{i-1}}$

Output: The bipartite graph \mathcal{G}_{2^i}

1.

$$\text{Lifts}(\mathcal{G}_{2^{i-1}}) := \bigcup_{\sigma \in \mathcal{G}_{2^{i-1}}^{\text{above}}} \text{Lifts}(\sigma)/\sim$$

2. Quotient $\text{Lifts}(\mathcal{G}_{2^{i-1}})$ by simultaneous conjugation in S_{2^i} to obtain $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$
3. Define $\mathcal{G}_{2^i}^{\text{below}} := \mathcal{G}_{2^{i-1}}^{\text{above}}$ and define $\mathcal{G}_{2^i}^{\text{above}}$ by representatives of $\text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$
4. For every pair $(\tilde{\sigma}, \sigma) \in \mathcal{G}_{2^i}^{\text{above}} \times \mathcal{G}_{2^i}^{\text{below}}$ place an edge between $\tilde{\sigma}$ and σ if and only if there is a triple in the equivalence class $[\tilde{\sigma}] \in \text{Lifts}(\mathcal{G}_{2^{i-1}})/\sim$ that is a lift of σ

Results : number of triples and passports

Theorem (M.)


The following tables list the number of isomorphism classes of 2-group Belyi maps, the number of passports, and number of lax passports respectively up to degree 256.

d	1	2	4	8	16	32	64	128	256
# triples	1	3	7	19	55	151	503	1799	7175


d	1	2	4	8	16	32	64	128	256
# passports	1	3	7	16	41	96	267	834	2893

d	1	2	4	8	16	32	64	128	256
# lax passports	1	1	3	6	14	31	85	257	882


Results : distribution of genera



Results : groups by nilpotency class



Results : passport sizes



Computing equations



A motivating example

Algorithm in characteristic $p \geq 3$

Implementation in characteristic zero

A refined conjecture



Passports

Recall that a passport \mathcal{P} consists of the data (g, G, λ) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d corresponding to conjugacy classes (C_0, C_1, C_∞) of S_d .

Passports

Recall that a passport \mathcal{P} consists of the data (g, G, λ) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d corresponding to conjugacy classes (C_0, C_1, C_∞) of S_d .

The size of \mathcal{P} is the cardinality of the set $\Sigma_{\mathcal{P}}$ defined by

$$\left\{ (\sigma_0, \sigma_1, \sigma_\infty) \in C_0 \times C_1 \times C_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1 \rangle = G \right\} / \sim$$

where \sim denotes simultaneous conjugation in S_d .

Passports

Recall that a passport \mathcal{P} consists of the data (g, G, λ) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d corresponding to conjugacy classes (C_0, C_1, C_∞) of S_d .

The size of \mathcal{P} is the cardinality of the set $\Sigma_{\mathcal{P}}$ defined by

$$\left\{ (\sigma_0, \sigma_1, \sigma_\infty) \in C_0 \times C_1 \times C_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1 \rangle = G \right\} / \sim$$

where \sim denotes simultaneous conjugation in S_d .

As a result of the action of $G_{\mathbb{Q}}$ on \mathcal{P} , the size of \mathcal{P} bounds the degree of the field of moduli of any Belyi map with passport \mathcal{P} .

Passports

Recall that a passport \mathcal{P} consists of the data (g, G, λ) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $\lambda = (\lambda_0, \lambda_1, \lambda_\infty)$ is a triple of partitions of d corresponding to conjugacy classes (C_0, C_1, C_∞) of S_d .

The size of \mathcal{P} is the cardinality of the set $\Sigma_{\mathcal{P}}$ defined by

$$\left\{ (\sigma_0, \sigma_1, \sigma_\infty) \in C_0 \times C_1 \times C_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1 \rangle = G \right\} / \sim$$

where \sim denotes simultaneous conjugation in S_d .

As a result of the action of $G_{\mathbb{Q}}$ on \mathcal{P} , the size of \mathcal{P} bounds the degree of the field of moduli of any Belyi map with passport \mathcal{P} .

To instead analyze $\text{Gal}(\mathbb{Q}^{\text{al}} | \mathbb{Q}^{\text{ab}})$ we *refine* the notion of a passport.

Refined passports

A **refined passport** \mathcal{P} consists of the data (g, G, c) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $c = (c_0, c_1, c_\infty)$ is a triple of conjugacy classes of G .

Refined passports

A **refined passport** \mathcal{P} consists of the data (g, G, c) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $c = (c_0, c_1, c_\infty)$ is a triple of conjugacy classes of G .

The size of \mathcal{P} is the cardinality of the set $\Sigma_{\mathcal{P}}$ defined by

$$\left\{ (\sigma_0, \sigma_1, \sigma_\infty) \in c_0 \times c_1 \times c_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1 \rangle = G \right\} / \sim$$

where $(\sigma_0, \sigma_1, \sigma_\infty) \sim (\sigma'_0, \sigma'_1, \sigma'_\infty)$ if there exists $\alpha \in \text{Aut}(G)$ with $\alpha(\sigma_s) = \sigma'_s$ for every $s \in \{0, 1, \infty\}$.

Refined passports

A **refined passport** \mathcal{P} consists of the data (g, G, c) where $g \in \mathbb{Z}_{\geq 0}$, G is a transitive subgroup of S_d and $c = (c_0, c_1, c_\infty)$ is a triple of conjugacy classes of G .

The size of \mathcal{P} is the cardinality of the set $\Sigma_{\mathcal{P}}$ defined by

$$\left\{ (\sigma_0, \sigma_1, \sigma_\infty) \in c_0 \times c_1 \times c_\infty : \sigma_\infty \sigma_1 \sigma_0 = 1 \text{ and } \langle \sigma_0, \sigma_1 \rangle = G \right\} / \sim$$

where $(\sigma_0, \sigma_1, \sigma_\infty) \sim (\sigma'_0, \sigma'_1, \sigma'_\infty)$ if there exists $\alpha \in \text{Aut}(G)$ with $\alpha(\sigma_s) = \sigma'_s$ for every $s \in \{0, 1, \infty\}$.

As was the case with passport, every permutation triple σ determines a refined passport $\mathcal{P}(\sigma)$.

A refined conjecture

Theorem (M.)

The size of $\mathcal{P}(\sigma)$ is equal to 1 for every 2-group permutation triple σ with degree ≤ 256 .

A refined conjecture

Theorem (M.)

The size of $\mathcal{P}(\sigma)$ is equal to 1 for every 2-group permutation triple σ with degree ≤ 256 .

Conjecture (ARC)

The size of $\mathcal{P}(\sigma)$ is equal to 1 for every 2-group permutation triple.

A refined conjecture

Theorem (M.)

The size of $\mathcal{P}(\sigma)$ is equal to 1 for every 2-group permutation triple σ with degree ≤ 256 .

Conjecture (ARC)

The size of $\mathcal{P}(\sigma)$ is equal to 1 for every 2-group permutation triple.

Theorem (M.)

ARC is true for 2-group permutation triples σ with $\langle \sigma \rangle$ dihedral.

Examples



$4T1-4,1,4-g0$

4T1-4,1,4-g0

Backup slides