

Macro PS1

Michael B. Nattinger*

November 6, 2020

1 Question 1

1.1 Part A

$$\begin{aligned} V(A_t, c_{t-1}) &= \max_{c_t, A_{t+1}} u(c_t, c_{t-1}) + \beta V(A_{t+1}, c_t) \\ \text{s.t. } A_{t+1} &= R(A_t - c_t) \end{aligned}$$

We can rewrite this as the following:

$$V(A_t, c_{t-1}) = \max_{c_t} u(c_t, c_{t-1}) + \beta V(R(A_t - c_t), c_t) \quad (1)$$

$$V(A_t, c_{t-1}) = \max_{A_t} u\left(A_t - \frac{A_{t+1}}{R}, c_{t-1}\right) + \beta V\left(A_{t+1}, A_t - \frac{A_{t+1}}{R}\right) \quad (2)$$

To ensure the solution is unique, with a strictly increasing, strictly concave value function that is differentiable on the interior of the feasible set, we require the following assumptions:

- $u(\cdot)$ is continuously differentiable, strictly concave, and strictly increasing in its arguments.
- $0 < \beta < 1$
- $\lim_{k \rightarrow 0} u'(k, u) = \lim_{k \rightarrow 0} u'(u, k) = \infty$
- $\lim_{k \rightarrow \infty} u'(k, u) = \lim_{k \rightarrow \infty} u'(u, k) = 0$
- The utility function is bounded?
- Do we need anything else?

*I worked on this assignment with my study group: Alex von Hafften, Andrew Smith, and Ryan Mather. I have also discussed problem(s) with Emily Case, Sarah Bass, and Danny Edgel.

Assuming the above conditions hold, we can solve the value function as it is written in (1) by taking a first order condition with respect to c_t , and then applying the envelope theorem twice:

$$\begin{aligned}
0 &= \frac{\partial u}{\partial c_t}(c_t, c_{t-1}) - R\beta \frac{\partial V}{\partial R(A_t - c_t)}(R(A_t - c_t), c_t) + \beta \frac{\partial V}{\partial c_t}(R(A_t - c_t), c_t) \\
\frac{\partial V}{\partial R(A_t - c_t)}(R(A_t - c_t), c_t) &= 999 \\
\frac{\partial V}{\partial c_t}(R(A_t - c_t), c_t) &= \frac{\partial u}{\partial c_t}(c_{t+1}, c_t) \\
\Rightarrow \frac{\partial u}{\partial c_t}(c_t, c_{t-1}) + \beta() &= R\beta()
\end{aligned}$$

Instead we will use the value function as it is written in (2) by taking a first order condition with respect to A_{t+1} , and then applying the envelope theorem twice:

$$\begin{aligned}
0 &= \frac{-1}{R} \frac{\partial u}{\partial A_t - \frac{A_{t+1}}{R}} \left(A_t - \frac{A_{t+1}}{R}, c_{t-1} \right) + \beta \frac{\partial V}{\partial A_{t+1}} \left(A_{t+1}, A_t - \frac{A_{t+1}}{R} \right) - \frac{\beta}{R} \frac{\partial V}{\partial A_t - \frac{A_{t+1}}{R}} \left(A_{t+1}, A_t - \frac{A_{t+1}}{R} \right) \\
\frac{\partial V}{\partial R(A_t - c_t)}(R(A_t - c_t), c_t) &= 999 \\
\frac{\partial V}{\partial c_t}(R(A_t - c_t), c_t) &= \frac{\partial u}{\partial c_t}(c_{t+1}, c_t) \\
\Rightarrow \frac{\partial u}{\partial c_t}(c_t, c_{t-1}) + \beta() &= R\beta()
\end{aligned}$$

2 Question 2

2.1 Part A

The sequence problem is to maximize profits:

$$\max_{\{x_t\}_{i=1}^{\infty}} \left(\frac{1}{1+r} \right)^t \left(ax_t - \frac{b}{2}x_t^2 - \frac{c}{2}(x_{t+1} - x_t)^2 \right)$$

We can rewrite this as a bellman equation in the following way:

$$V(x) = \max_y ax - \frac{b}{2}x^2 - \frac{c}{2}(y - x)^2 + \delta V(y) \quad (3)$$

We can rewrite this as follows:

$$T(v)(x) = \max_y ax - \frac{b}{2}x^2 - \frac{c}{2}(y - x)^2 + \delta v(y) \quad (4)$$

where the fixed point of our T operator in (4) is the solution to the Bellman equation in (3).

2.2 Part B

Let $L < 0$ be arbitrary. If we set $y = 0, x < \frac{L}{a}$ then $F(x, y) = ax - \frac{b}{2}x^2 - \frac{c}{2}(y - x)^2 \leq ax < L$ so F is unbounded below.

This F function is of the form of a polynomial with limits of negative infinity in all directions, so we can find the global maximum by taking first order conditions:

$$\begin{aligned} 0 &= a - bx + c(y - x) \\ 0 &= -c(y - x) \Rightarrow y - x = 0 \Rightarrow y = x \\ \Rightarrow y = x &= \frac{a}{b} \\ F\left(\frac{a}{b}, \frac{a}{b}\right) &= a\left(\frac{a}{b}\right) - \frac{b}{2}\left(\frac{a}{b}\right)^2 - 0 \\ &= \frac{a^2}{2b} \end{aligned}$$

Therefore, the maximum value F can take is $\frac{a^2}{2b}$
We can find bounds on \hat{v} in the following way:

$$\begin{aligned} \hat{v} &= \frac{a^2}{2b} + \delta \hat{v} \\ \Rightarrow \hat{v} &= \frac{a^2}{2b(1 - \delta)} \end{aligned}$$

2.3 Part C

$$\begin{aligned} T\hat{v}(x) &= \max_y ax - \frac{b}{2}x^2 - \frac{c}{2}(y - x)^2 + \delta \hat{v} \\ 0 &= -c(y - x) \Rightarrow y = x, \\ \Rightarrow T\hat{v}(x) &= ax - \frac{b}{2}x^2 + \delta \hat{v} \\ &\leq \frac{a^2}{2b} + \delta \frac{a^2}{2b(1 - \delta)} = \frac{a^2}{2b(1 - \delta)} \\ &= \hat{v}. \end{aligned}$$

2.4 Part D

We showed the base case of the induction in Part C. Now we will show the induction step.

Assume that $T^n \hat{v}(x)$ takes the form $T^n \hat{v}(x) = \alpha_n x - \frac{1}{2} \beta_n x^2 + \gamma_n$. Then,

$$\begin{aligned} T^{n+1} \hat{v}(x) &= \max_y ax - \frac{b}{2} x^2 - \frac{c}{2} (y - x)^2 + \delta(\alpha_n x - \frac{1}{2} \beta_n x^2 + \gamma_n) \\ y = x \Rightarrow T^{n+1} \hat{v}(x) &= ax - \frac{b}{2} x^2 + \delta \alpha_n x - \delta \frac{1}{2} \beta_n x^2 + \delta \gamma_n \\ &= (a + \delta \alpha_n) x - \frac{b + \delta \beta_n}{2} x^2 + \delta \gamma_n \\ &= \alpha_{n+1} x - \frac{1}{2} \beta_{n+1} x^2 + \gamma_{n+1} \end{aligned}$$

where $\alpha_{n+1} = (a + \delta \alpha_n)$, $\beta_{n+1} = b + \delta \beta_n$, $\gamma_{n+1} = \delta \gamma_n$.

2.5 Part E

Note that $\alpha_n = a + \delta a + \delta^2 a + \dots$, $\beta_n = b + \delta b + \delta^2 b + \dots$, $\gamma_n = \delta^n \hat{v}$. Thus, we can take the limit of α, β as geometric sums, and the limit of γ_n is 0. Therefore,

$$\begin{aligned} \tilde{V} &= \lim_{n \rightarrow \infty} T^n \hat{v} = \frac{a}{1 - \delta} x - \frac{1}{2} \frac{b}{1 - \delta} x^2. \\ T\tilde{V} &= \max_y ax - \frac{b}{2} x^2 - \frac{c}{2} (y - x)^2 + \delta \frac{a}{1 - \delta} x - \frac{1}{2} \frac{b}{1 - \delta} x^2, \\ y = x \Rightarrow T\tilde{V} &= ax - \frac{b}{2} x^2 + \delta \frac{a}{1 - \delta} x - \frac{1}{2} \frac{b}{1 - \delta} x^2 \\ &= \frac{a}{1 - \delta} x - \frac{1}{2} \frac{b}{1 - \delta} x^2 \\ &= \tilde{T}. \end{aligned}$$

Therefore, the limit function \tilde{V} satisfies the Bellman equation.

3 Question 3

$$V(k) = \max_{k'} \pi(k) - \gamma(k' - (1 - \delta)k) + \frac{1}{R} V(k')$$

$$\begin{aligned} 0 &= -\gamma'(k' - (1 - \delta)k) + \frac{1}{R} (\pi'(k') - (1 - \delta)\gamma'(k'' - (1 - \delta)k')) \\ \Rightarrow \pi'(k') &= R\gamma'(k' - (1 - \delta)k) + (1 - \delta)\gamma'(k'' - (1 - \delta)k') \end{aligned}$$

4 Question 4