

Ex0909A3: The generalized canonical transformation

Submitted by: Michael Pukshanski, Lior Weitzhandler

The problem: A system is described by the Hamiltonian $H(q,p,t)$. The coordinate q is transformed $q \rightarrow Q = \psi(q,t)$

1. Find the most general transformation $p \rightarrow P(q,p,t)$ so that $(q,p) \rightarrow (Q,P)$ will be canonical.
2. Calculate the appropriate generating function (explain why $F_1(q,Q,t)$ isn't relevant).
3. Given the new Hamiltonian $H' = 0$. Find the original Hamiltonian.
4. Prove that for $\psi(q+\omega t)$, the original Hamiltonian is a function of p alone. What does it mean?

The solution:

1. answer 1
2. answer 2
3. answer 3
4. answer 4
5. This is how you write an equation

$$\mathcal{I} = \int_0^z d^4x \sqrt{-g} \left[\varphi^2 (\mathcal{R} - 6sg^{\mu\nu} \kappa_\mu \kappa_\nu) + 4\omega g^{\mu\nu} D_\mu \varphi D_\nu \varphi + \lambda \varphi^4 + \frac{1}{4} g^{\mu\nu} g^{\lambda\sigma} X_{\mu\lambda} X_{\nu\sigma} \right]. \quad (1)$$

6. If you wish to add a set of equations (add the ampersands to align the two equations):

$$s = \frac{3+2\omega}{2\omega} \neq 1, \quad (2)$$

$$D_\mu \varphi = \varphi_{;\mu} + s \kappa_\mu \varphi, \quad (3)$$

If you wish to have equations with no numbers add an asterisk

$$g^{\mu\nu} (\varphi_{;\mu}^2 + 2\kappa_\mu \varphi^2)_{;\nu} = \frac{\partial W_{eff}(\varphi^2)}{\partial \varphi^2},$$
$$\frac{\partial W_{eff}(\varphi^2)}{\partial \varphi^2} = \frac{1}{3+2} \omega \left(\frac{1}{2} \varphi V'(\varphi) - 2V(\varphi) \right).$$

To write a vector

$$\vec{r} = x\hat{x} + y\hat{y} + z\hat{z} \quad ; \quad \vec{r}_i = |\vec{r}| \hat{r} \quad (4)$$

To write a matrix

$$\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \text{"\quad\quad" makes space in equations and this is how you add text.} \quad (5)$$

You can also use “align” to tightly align the equal sign, i.e “=” is aligned: see Eq. (??) and Eq. (??), also note that every end of an equation needs to be punctuated, i.e : “,” or “.”, according to the sentence.

$$\lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad (6)$$

$$\lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -2 & 0 \end{pmatrix}, \quad (7)$$

and centered

$$\left[\frac{\lambda_i}{2}, \frac{\lambda_j}{2} \right] = i \sum_{k=1}^n f^{ijk} \frac{\lambda_k}{2},$$

$$f^{147} = f^{165} = f^{246} = f^{257} = f^{345} = f^{376} = \frac{1}{2}.$$

7. Known arguments are **not** *written in italic mode*, some have special syntax in LaTeX otherwise just use “\mathrm”:

$$\cos(\omega t); \quad \sin(\omega t); \quad \mathrm{Tr} \left[\hat{A} \right]; \quad \det \left[\hat{\mathcal{A}} \right]; \quad \tan(\omega t); \quad \mathrm{e}^{\frac{t}{\tau}}; \quad \exp \left(\frac{t}{\tau} \right); \quad \log \left(\frac{t}{\tau} \right). \quad (8)$$

For all Latex related knowledge go to <https://en.wikibooks.org/wiki/LaTeX>

To include a picture use, note that the picture file has to be in the same folder as the *.tex file.