CS 107, Probability, Spring 2019 Lecture 05

Michael Poghosyan

AUA

25 January 2019

Content

- Classical Probability Models: Countably Infinite Sample Spaces
- Classical Probability Models: Finite Sample Spaces with Equally Likely Outcomes

Surprize

Surprize

Quiz Time!

Let us do a scientific experiment:

Let us do a scientific experiment: Take a deep breath. Keep it for few seconds.

Let us do a scientific experiment: Take a deep breath. Keep it for few seconds. Do you remember Julius Caesar and his last words?

Let us do a scientific experiment: Take a deep breath. Keep it for few seconds. Do you remember Julius Caesar and his last words?



Figure: Gaius Julius Caesar

Let us do a scientific experiment: Take a deep breath. Keep it for at least for few seconds. Do you remember Julius Caesar and his last words?




Figure: Gaius Julius Caesar

Let us do a scientific experiment: Take a deep breath. Keep it for at least for few seconds. Do you remember Julius Caesar and his last words?

Let us do a scientific experiment: Take a deep breath. Keep it for at least for few seconds. Do you remember Julius Caesar and his last words?

You too, Brutus.

Let us do a scientific experiment: Take a deep breath. Keep it for at least for few seconds. Do you remember Julius Caesar and his last words?

You too, Brutus.

Vnimanie, the question:

Let us do a scientific experiment: Take a deep breath. Keep it for at least for few seconds. Do you remember Julius Caesar and his last words?

You too, Brutus.

Vnimanie, the question: what are the chances that you just inhaled a molecule which the great Caesar exhaled when saying his last words?

Let us do a scientific experiment: Take a deep breath. Keep it for at least for few seconds. Do you remember Julius Caesar and his last words?

You too, Brutus.

Vnimanie, the question: what are the chances that you just inhaled a molecule which the great Caesar exhaled when saying his last words?

The answer is: the probability is larger than 99%! $\stackrel{..}{\sim}$

We continue considering Discrete Models. We assume that we have a countably infinite Sample Space:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n, \omega_{n+1}...\},\$$

We continue considering Discrete Models. We assume that we have a countably infinite Sample Space:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n, \omega_{n+1}...\},\$$

We take $\mathcal{F}=2^{\Omega}$.

We continue considering Discrete Models. We assume that we have a countably infinite Sample Space:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n, \omega_{n+1}...\},\$$

We take $\mathcal{F}=2^{\Omega}$. Then we take arbitrary real numbers $p_k, k \in \mathbb{N}$ with

$$p_k \ge 0, \quad k \in \mathbb{N} \quad \text{and} \quad \sum_{k=1}^{\infty} p_k = 1,$$

and define



We continue considering Discrete Models. We assume that we have a countably infinite Sample Space:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n, \omega_{n+1}...\},\$$

We take $\mathcal{F}=2^{\Omega}$. Then we take arbitrary real numbers $p_k, k \in \mathbb{N}$ with

$$p_k \ge 0, \quad k \in \mathbb{N} \quad \text{and} \quad \sum_{k=1}^{\infty} p_k = 1,$$

and define

$$\mathbb{P}(\{\omega_k\}) = p_k, \qquad k \in \mathbb{N}.$$



In the table form, this looks like

Outcome	ω_1	ω_2	ω_3	 ω_n	
Probability	p_1	p_2	<i>p</i> ₃	 p_n	

In the table form, this looks like

Now, we define for any nonempty event $A \in \mathcal{F}$ (i.e., for any nonempty subset $A \subset \Omega$),

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k,$$

and also $\mathbb{P}(\varnothing) = 0$.



In the table form, this looks like

Now, we define for any nonempty event $A \in \mathcal{F}$ (i.e., for any nonempty subset $A \subset \Omega$),

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k,$$

and also $\mathbb{P}(\varnothing) = 0$.

Now we have the complete Model.



The Moral is:

The Moral is:

For the Discrete case (i.e., when the Sample Space is either finite or countably infinite), it is enough to define the Probability Measure only for the outcomes: $\mathbb{P}(\{\omega_k\})$.

The Moral is:

For the Discrete case (i.e., when the Sample Space is either finite or countably infinite), it is enough to define the Probability Measure only for the outcomes: $\mathbb{P}(\{\omega_k\})$.

The Probability of any other Event can be obtained from these outcome probabilities.

The Moral is:

For the Discrete case (i.e., when the Sample Space is either finite or countably infinite), it is enough to define the Probability Measure only for the outcomes: $\mathbb{P}(\{\omega_k\})$.

The Probability of any other Event can be obtained from these outcome probabilities.

Example:

Now assume we have a Discrete Model with finitely many outcomes:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n\}$$

Now assume we have a Discrete Model with finitely many outcomes:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n\}$$

One of the important Probability Models is the Equally Likely Outcomes (or the uniform discrete distribution) case:

$$\mathbb{P}(\{\omega_1\}) = \mathbb{P}(\{\omega_2\}) = \dots = \mathbb{P}(\{\omega_n\}) =$$

Now assume we have a Discrete Model with finitely many outcomes:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n\}$$

One of the important Probability Models is the Equally Likely Outcomes (or the uniform discrete distribution) case:

$$\mathbb{P}(\{\omega_1\}) = \mathbb{P}(\{\omega_2\}) = \dots = \mathbb{P}(\{\omega_n\}) = \frac{1}{n}.$$

Now assume we have a Discrete Model with finitely many outcomes:

$$\Omega = \{\omega_1, \omega_2, ..., \omega_n\}$$

One of the important Probability Models is the Equally Likely Outcomes (or the uniform discrete distribution) case:

$$\mathbb{P}(\{\omega_1\}) = \mathbb{P}(\{\omega_2\}) = \dots = \mathbb{P}(\{\omega_n\}) = \frac{1}{n}.$$

In the table form:



$$\mathbb{P}(A) =$$

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k =$$

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k = \sum_{\omega_k \in A} \frac{1}{n} =$$

Now, consider an event $A \subset \Omega$. Then

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k = \sum_{\omega_k \in A} \frac{1}{n} =$$

 $=\frac{1}{n}$ (number of elements in A) =

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k = \sum_{\omega_k \in A} \frac{1}{n} =$$

$$= \frac{1}{n} \cdot (\text{number of elements in } A) = \frac{\text{number of elements in } A}{\text{total number of elements in } \Omega}.$$

Now, consider an event $A \subset \Omega$. Then

$$\mathbb{P}(A) = \sum_{\omega_k \in A} p_k = \sum_{\omega_k \in A} \frac{1}{n} =$$

 $=\frac{1}{n}\cdot(\text{number of elements in }A)=\frac{\text{number of elements in }A}{\text{total number of elements in }\Omega}.$ Or, in other words,

$$\mathbb{P}(A) = \frac{\text{number of elements favorable for the event } A}{\text{total number of possible outcomes}} = \frac{\#A}{\#\Omega}.$$

• What is the Equally Probable Outcomes (or the Discrete Uniform) Model?

- What is the Equally Probable Outcomes (or the Discrete Uniform) Model?
- Give an example of an Equally Probable Outcomes model.

- What is the Equally Probable Outcomes (or the Discrete Uniform) Model?
- Give an example of an Equally Probable Outcomes model.
- Is it possible to define the Equally Probable Outcomes Model for the Countably Infinite Sample Space case?

- What is the Equally Probable Outcomes (or the Discrete Uniform) Model?
- Give an example of an Equally Probable Outcomes model.
- Is it possible to define the Equally Probable Outcomes Model for the Countably Infinite Sample Space case?
- How to calculate the probability of an event in the Equally Probable Outcomes Case?

Example: Assume we have a box containing 12 red balls, 21 white balls and 14 green balls. We pick a ball at random from the box.

Example: Assume we have a box containing 12 red balls, 21 white balls and 14 green balls. We pick a ball at random from the box.

• What is the probability that we will choose a green ball?

Example: Assume we have a box containing 12 red balls, 21 white balls and 14 green balls. We pick a ball at random from the box.

- What is the probability that we will choose a green ball?
- Why this kind of problems are interesting/important in Probability Theory? Can you give a real -life example?

Example: Assume we have a box containing 12 red balls, 21 white balls and 14 green balls. We pick a ball at random from the box.

- What is the probability that we will choose a green ball?
- Why this kind of problems are interesting/important in Probability Theory? Can you give a real -life example?

Example: In our Parliament we have 132 seats. 88 Seats are for "My Step" Alliance, 26 seats are for "Prosperous Armenia" party and 18 are for the "Bright Armenia". What is the probability that a randomly chosen parliamentarian will be from the "Bright Armenia"?