

# AUA CS 108, Statistics, Fall 2019

## Lecture 43 (-1)

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

04 Dec 2019

# Contents

- ▶ Linear Regression

# Last Lecture ReCap

- ▶ Maybe we can skip this this time?

# Intro to Linear Regression

Recall that our Mission (for these last few lectures 😊) was to give a Model for the Relationship between two and more Variables.

---

# Intro to Linear Regression

Recall that our Mission (for these last few lectures 😊) was to give a Model for the Relationship between two and more Variables.

Usually, we assume that the Joint Distribution of  $X$  and  $Y$  is given in the following form: for the fixed value of  $X$ , say,  $X = x$ ,  $Y$  has some Distribution, depending (or, maybe independent of)  $x$ .

---

<sup>1</sup>And, in fact, we can recover the Joint Distribution, having the Distribution of  $X$  and the Conditional Distribution  $Y|X = x$ .

# Intro to Linear Regression

Recall that our Mission (for these last few lectures 😊) was to give a Model for the Relationship between two and more Variables.

Usually, we assume that the Joint Distribution of  $X$  and  $Y$  is given in the following form: for the fixed value of  $X$ , say,  $X = x$ ,  $Y$  has some Distribution, depending (or, maybe independent of)  $x$ . We will denote that Distribution by

$$Y|X = x$$

and call it the<sup>1</sup> **Conditional Distribution of  $Y$  given  $X = x$** .

---

<sup>1</sup>And, in fact, we can recover the Joint Distribution, having the Distribution of  $X$  and the Conditional Distribution  $Y|X = x$ .

# Intro to Linear Regression

Recall that our Mission (for these last few lectures 😊) was to give a Model for the Relationship between two and more Variables.

Usually, we assume that the Joint Distribution of  $X$  and  $Y$  is given in the following form: for the fixed value of  $X$ , say,  $X = x$ ,  $Y$  has some Distribution, depending (or, maybe independent of)  $x$ . We will denote that Distribution by

$$Y|X = x$$

and call it the<sup>1</sup> **Conditional Distribution of  $Y$  given  $X = x$** .

**Example:** Say,  $(X, Y) \sim \text{Unif}([0, 1] \times [0, 2])$ . What is  $Y|X = 1$ , or, in general,  $Y|X = x$ ?

---

<sup>1</sup>And, in fact, we can recover the Joint Distribution, having the Distribution of  $X$  and the Conditional Distribution  $Y|X = x$ .

# Intro to Linear Regression

Recall that our Mission (for these last few lectures 😊) was to give a Model for the Relationship between two and more Variables.

Usually, we assume that the Joint Distribution of  $X$  and  $Y$  is given in the following form: for the fixed value of  $X$ , say,  $X = x$ ,  $Y$  has some Distribution, depending (or, maybe independent of)  $x$ . We will denote that Distribution by

$$Y|X = x$$

and call it the<sup>1</sup> **Conditional Distribution of  $Y$  given  $X = x$** .

**Example:** Say,  $(X, Y) \sim \text{Unif}([0, 1] \times [0, 2])$ . What is  $Y|X = 1$ , or, in general,  $Y|X = x$ ?

**Example:** Now, assume  $(X, Y) \sim \text{Unif}(D)$ , where  $D$  is the triangle with vertices at  $(-1, 0)$ ,  $(0, 1)$  and  $(1, 0)$ . What is  $Y|X = x$ ?

---


<sup>1</sup>And, in fact, we can recover the Joint Distribution, having the Distribution of  $X$  and the Conditional Distribution  $Y|X = x$ .



## Example

**Example:** Assume  $(X, Y) \sim \mathcal{N}(\mu, \Sigma)$ , where

$$\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 4 & 0.5 \\ 0.5 & 1 \end{bmatrix}$$



What is  $Y|X = x$ ?

## Intro to Linear Regression

We will simplify our model and consider the case when  $\mathbf{X}$  is not Stochastic (is not Random), but  $Y|\mathbf{X} = \mathbf{x}$  is Random.

## Intro to Linear Regression

We will simplify our model and consider the case when  $\mathbf{X}$  is not Stochastic (is not Random), but  $Y|\mathbf{X} = \mathbf{x}$  is Random.

**Example:** Assume  $x \in [0, 2\pi]$  and


$$Y|X = x \sim \mathcal{N}(\sin x, 0.3^2)$$

# Intro to Linear Regression

We will simplify our model and consider the case when  $\mathbf{X}$  is not Stochastic (is not Random), but  $Y|\mathbf{X} = \mathbf{x}$  is Random.

**Example:** Assume  $x \in [0, 2\pi]$  and

$$Y|X = x \sim \mathcal{N}(\sin x, 0.3^2)$$



## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ .

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number.

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$  ?

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$  ? Well, of course, the first and most important numerical characteristics of a r.v. is its



## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$  ? Well, of course, the first and most important numerical characteristics of a r.v. is its **Mean**:  $\mathbb{E}(Y|\mathbf{X} = \mathbf{x})$ .

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$ ? Well, of course, the first and most important numerical characteristics of a r.v. is its **Mean**:  $\mathbb{E}(Y|\mathbf{X} = \mathbf{x})$ . Since for each  $\mathbf{x}$  this will be a number, we will obtain a (deterministic) function of  $\mathbf{x}$ , called the **Regression Function**:

$$RegFun(\mathbf{x}) = \mathbb{E}(Y|\mathbf{X} = \mathbf{x}).$$

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$ ? Well, of course, the first and most important numerical characteristics of a r.v. is its **Mean**:  $\mathbb{E}(Y|\mathbf{X} = \mathbf{x})$ . Since for each  $\mathbf{x}$  this will be a number, we will obtain a (deterministic) function of  $\mathbf{x}$ , called the **Regression Function**:

$$RegFun(\mathbf{x}) = \mathbb{E}(Y|\mathbf{X} = \mathbf{x}).$$

So our r.v.  $Y|\mathbf{X} = \mathbf{x}$  is a r.v. around  $RegFun(\mathbf{x})$ , or simply,

$$(Y|\mathbf{X} = \mathbf{x}) = RegFun(\mathbf{x}) + \varepsilon$$

where  $\varepsilon$  is a r.v. (just take  $\varepsilon = (Y|\mathbf{X} = \mathbf{x}) - RegFun(\mathbf{x})$ ),

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$ ? Well, of course, the first and most important numerical characteristics of a r.v. is its **Mean**:  $\mathbb{E}(Y|\mathbf{X} = \mathbf{x})$ . Since for each  $\mathbf{x}$  this will be a number, we will obtain a (deterministic) function of  $\mathbf{x}$ , called the **Regression Function**:

$$RegFun(\mathbf{x}) = \mathbb{E}(Y|\mathbf{X} = \mathbf{x}).$$

So our r.v.  $Y|\mathbf{X} = \mathbf{x}$  is a r.v. around  $RegFun(\mathbf{x})$ , or simply,

$$(Y|\mathbf{X} = \mathbf{x}) = RegFun(\mathbf{x}) + \varepsilon$$

where  $\varepsilon$  is a r.v. (just take  $\varepsilon = (Y|\mathbf{X} = \mathbf{x}) - RegFun(\mathbf{x})$ ), or, which is the same,

$$Y = RegFun(\mathbf{X}) + \varepsilon.$$

## Intro to Linear Regression

Now, about the Regression. As we talked above,  $Y|\mathbf{X} = \mathbf{x}$  is a r.v., for each  $\mathbf{x}$ . Now, we want to give a deterministic characteristic for each r.v.  $Y|\mathbf{X} = \mathbf{x}$ , i.e., for each  $\mathbf{x}$ , we want to describe the r.v.  $Y|\mathbf{X} = \mathbf{x}$  by just giving one number. Can you give just one number that will describe  $Y|\mathbf{X} = \mathbf{x}$ ? Well, of course, the first and most important numerical characteristics of a r.v. is its **Mean**:  $\mathbb{E}(Y|\mathbf{X} = \mathbf{x})$ . Since for each  $\mathbf{x}$  this will be a number, we will obtain a (deterministic) function of  $\mathbf{x}$ , called the **Regression Function**:

$$RegFun(\mathbf{x}) = \mathbb{E}(Y|\mathbf{X} = \mathbf{x}).$$

So our r.v.  $Y|\mathbf{X} = \mathbf{x}$  is a r.v. around  $RegFun(\mathbf{x})$ , or simply,

$$(Y|\mathbf{X} = \mathbf{x}) = RegFun(\mathbf{x}) + \varepsilon$$

where  $\varepsilon$  is a r.v. (just take  $\varepsilon = (Y|\mathbf{X} = \mathbf{x}) - RegFun(\mathbf{x})$ ), or, which is the same,

$$Y = RegFun(\mathbf{X}) + \varepsilon.$$

Here, importantly,  $\mathbb{E}(\varepsilon) = 0$ , for any  $\mathbf{x}$ .

# Intro to Linear Regression

**Example:** Say, given the height of a person, the weight will be some fixed number  $\pm$  something random, i.e.,

$$Weight = g(Height) + random\ term.$$

# Intro to Linear Regression

**Example:** Say, given the height of a person, the weight will be some fixed number  $\pm$  something random, i.e.,

$$Weight = g(Height) + random\ term.$$

Say, see Wiki page for [BMI](#).