

# CS 107, Probability, Spring 2019

## Lecture 35

Michael Poghosyan

AUA

15 April 2019

- Multivariate Normal (Gaussian) Distribution
- Transformation of Random Vectors
- Independent Random Variables

## R Code for Bivariate Normal

```
mu <- c(0,0) # The Mean
Sigma <- matrix(c(1, .5, .5, 1), nrow = 2) #Cov Matrix
#Version 1
library(MASS)
data <- mvrnorm(5000, mu = mu, Sigma = Sigma )
plot(data, pch = 20, asp = 1, cex = 0.6)
#Version 2
#install.packages("mvtnorm")
library(mvtnorm)
data <- rmvnorm(1000, mean = mu, sigma = Sigma)
plot(data, pch = 20, asp = 1, xlim = c(-3,3))
```

# Multivariate Normal (Gaussian) Distribution

Now, about the MultiVariate Normal Distribution. The definition goes in line with the 2D (BiVariate) case:

# Multivariate Normal (Gaussian) Distribution

Now, about the MultiVariate Normal Distribution. The definition goes in line with the 2D (BiVariate) case:

We assume we are given a vector in  $\mathbb{R}^n$ :

$$\mu = \begin{bmatrix} \mu_1 \\ \vdots \\ \mu_n \end{bmatrix},$$

# Multivariate Normal (Gaussian) Distribution

Now, about the MultiVariate Normal Distribution. The definition goes in line with the 2D (BiVariate) case:

We assume we are given a vector in  $\mathbb{R}^n$ :

$$\mu = \begin{bmatrix} \mu_1 \\ \vdots \\ \mu_n \end{bmatrix},$$

and a Symmetric Positive Definite Matrix

$$\Sigma = \begin{bmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1n} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{n1} & \sigma_{n2} & \cdots & \sigma_{nn} \end{bmatrix}$$

# Multivariate Normal (Gaussian) Distribution

## Multivariate Normal (Gaussian) Distribution

We say that the r. vector  $\mathbf{X} = (X_1, X_2, \dots, X_n)$  has a Multivariate Normal (or Gaussian) Distribution with the **mean**  $\mu$  and the **covariance matrix**  $\Sigma$ , and we will write

$$\mathbf{X} \sim \mathcal{N}(\mu, \Sigma),$$

# Multivariate Normal (Gaussian) Distribution

## Multivariate Normal (Gaussian) Distribution

We say that the r. vector  $\mathbf{X} = (X_1, X_2, \dots, X_n)$  has a Multivariate Normal (or Gaussian) Distribution with the **mean**  $\mu$  and the **covariance matrix**  $\Sigma$ , and we will write

$$\mathbf{X} \sim \mathcal{N}(\mu, \Sigma),$$

if the Joint PDF of  $\mathbf{X}$  is given by

$$f(\mathbf{x}) = \frac{1}{\sqrt{(2\pi)^n \det(\Sigma)}} \cdot \exp \left\{ -\frac{1}{2} (\mathbf{x} - \mu)^T \cdot \Sigma^{-1} \cdot (\mathbf{x} - \mu) \right\},$$

for any  $\mathbf{x} \in \mathbb{R}^n$ .



# Multivariate Normal (Gaussian) Distribution

Another, equivalent definition of the Multivariate Normal Distribution is the following:

## Multivariate Normal (Gaussian) Distribution

We say that the r. vector  $\mathbf{X} = (X_1, X_2, \dots, X_n)$  has a Multivariate Normal (or Gaussian) Distribution, if for any numbers<sup>a</sup>  $a_1, a_2, \dots, a_n \in \mathbb{R}$ , the r.v.

$$Y = a_1 \cdot X_1 + a_2 \cdot X_2 + \dots + a_n \cdot X_n$$

is Normally Distributed.

---

<sup>a</sup>We need to take care of the case  $a_k = 0$  for all  $k$ . We can exclude this case, say, in the definition.

# Marginals of Multivariate Normal Distribution

It is remarkable that the Marginal Distributions of Multivariate Normal Distribution are again Normal. In particular, if

$$(X, Y) \sim \mathcal{N} \left( \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}, \begin{bmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{bmatrix} \right)$$

# Marginals of Multivariate Normal Distribution

It is remarkable that the Marginal Distributions of Multivariate Normal Distribution are again Normal. In particular, if

$$(X, Y) \sim \mathcal{N} \left( \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}, \begin{bmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{bmatrix} \right)$$

then

$$X \sim \mathcal{N}(\mu_1, \sigma_{11}) \quad \text{and} \quad Y \sim \mathcal{N}(\mu_2, \sigma_{22}).$$

# Transformation of Random Vectors

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment.

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ .

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ . Say,  $X + Y$ ,  $\sin(X + Y)$ ,...

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .



# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ . Say,  $X + Y$ ,  $\sin(X + Y)$ ,...
- We can use 2 Real-Valued functions  $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$  to transform  $(X, Y)$  onto new r.vector  $(U, V)$  by

$$U = g_1(X, Y), \quad V = g_2(X, Y);$$

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ . Say,  $X + Y$ ,  $\sin(X + Y)$ ,...
- We can use 2 Real-Valued functions  $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$  to transform  $(X, Y)$  onto new r.vector  $(U, V)$  by

$$U = g_1(X, Y), \quad V = g_2(X, Y);$$

$$\text{Say, } U = X + Y, \quad V = X - Y;$$

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ . Say,  $X + Y$ ,  $\sin(X + Y)$ ,...
- We can use 2 Real-Valued functions  $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$  to transform  $(X, Y)$  onto new r.vector  $(U, V)$  by

$$U = g_1(X, Y), \quad V = g_2(X, Y);$$

Say,  $U = X + Y$ ,  $V = X - Y$ ; or  $U = X + 2Y$ ,  $V = X \cdot Y$ ,...

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ . Say,  $X + Y$ ,  $\sin(X + Y)$ ,...
- We can use 2 Real-Valued functions  $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$  to transform  $(X, Y)$  onto new r.vector  $(U, V)$  by

$$U = g_1(X, Y), \quad V = g_2(X, Y);$$

Say,  $U = X + Y$ ,  $V = X - Y$ ; or  $U = X + 2Y$ ,  $V = X \cdot Y$ ,...

- We can obtain 3 r.v.s from our  $X, Y$ , ...

---

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Assume we have 2 r.v.s  $X$  and  $Y$ , defined on the same Experiment. Using  $X$  and  $Y$ , we can obtain new r.v.s.

- Say, we can use a Real-Valued function<sup>1</sup>  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$  to obtain a 1D r.v.  $Z = g(X, Y)$ . Say,  $X + Y$ ,  $\sin(X + Y)$ ,...
- We can use 2 Real-Valued functions  $g_1, g_2: \mathbb{R}^2 \rightarrow \mathbb{R}$  to transform  $(X, Y)$  onto new r.vector  $(U, V)$  by

$$U = g_1(X, Y), \quad V = g_2(X, Y);$$

Say,  $U = X + Y$ ,  $V = X - Y$ ; or  $U = X + 2Y$ ,  $V = X \cdot Y$ ,...

- We can obtain 3 r.v.s from our  $X, Y$ , ...

Here we will consider the 1D case, when we transform our pair  $X, Y$  onto 1 r.v.  $Z$ .

<sup>1</sup>It is enough to have our  $g$  defined not on the whole  $\mathbb{R}^2$  but on all possible pairs  $(X, Y)$ , in some subregion of  $\mathbb{R}^2$ .

# Transformation of Random Vectors

Now assume we have a r.vector  $(X, Y)$ , and we form a new r.variable (1D r.v.!) from  $X$  and  $Y$ . Say, we take a function  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $g = g(x, y)$ , and consider the r.v.

$$Z = g(X, Y).$$

# Transformation of Random Vectors

Now assume we have a r.vector  $(X, Y)$ , and we form a new r.variable (1D r.v.!) from  $X$  and  $Y$ . Say, we take a function  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $g = g(x, y)$ , and consider the r.v.

$$Z = g(X, Y).$$

**Example:** Say, we are interested in the r.v.  $Z = 2X - 3Y$ ,  
 $T = X^2 + Y^2$ ,  $K = \sin(X^2 + Y^2)$ , ...

# Transformation of Random Vectors

Now assume we have a r.vector  $(X, Y)$ , and we form a new r.variable (1D r.v.!) from  $X$  and  $Y$ . Say, we take a function  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $g = g(x, y)$ , and consider the r.v.

$$Z = g(X, Y).$$

**Example:** Say, we are interested in the r.v.  $Z = 2X - 3Y$ ,  
 $T = X^2 + Y^2$ ,  $K = \sin(X^2 + Y^2)$ , ...

The general problem here is:

Knowing the Joint Distribution of  $X$  and  $Y$ , find the Distribution of  $Z = g(X, Y)$ .



# Transformation of Random Vectors

Now assume we have a r.vector  $(X, Y)$ , and we form a new r.variable (1D r.v.!) from  $X$  and  $Y$ . Say, we take a function  $g: \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $g = g(x, y)$ , and consider the r.v.

$$Z = g(X, Y).$$

**Example:** Say, we are interested in the r.v.  $Z = 2X - 3Y$ ,  $T = X^2 + Y^2$ ,  $K = \sin(X^2 + Y^2)$ , ...

The general problem here is:

Knowing the Joint Distribution of  $X$  and  $Y$ , find the Distribution of  $Z = g(X, Y)$ .

In particular, very important is the question of distribution of the sum  $Z = X + Y$ , and, in general, of the sum of  $n$  r.v.s

$$Y = X_1 + X_2 + \dots + X_n.$$

# Examples

Some Examples:

# Examples

Some Examples:

- Say, my portfolio consists of 100 shares of Google (Alphabet Inc.) and 200 shares of Amazon.

# Examples

Some Examples:

- Say, my portfolio consists of 100 shares of Google (Alphabet Inc.) and 200 shares of Amazon. So I am a Rich Man!! 😊

# Examples

Some Examples:

- Say, my portfolio consists of 100 shares of Google (Alphabet Inc.) and 200 shares of Amazon. So I am a Rich Man!! 😊 Let  $Z$  be the price of my portfolio at the end of this year. Then

$$Z = 100 \cdot GOOG + 200 \cdot AMZN,$$

where  $GOOG$  and  $AMZN$  are r.v. showing the prices of Google and Amazon Stocks at the end of the year.

# Examples

Some Examples:

- Say, my portfolio consists of 100 shares of Google (Alphabet Inc.) and 200 shares of Amazon. So I am a Rich Man!! 😊 Let  $Z$  be the price of my portfolio at the end of this year. Then

$$Z = 100 \cdot GOOG + 200 \cdot AMZN,$$

where  $GOOG$  and  $AMZN$  are r.v. showing the prices of Google and Amazon Stocks at the end of the year.

- My expenses at AUA cafeteria next week can be written as

$$Z = X_1 + X_3 + X_5,$$

where  $X_i$  is

# Examples

Some Examples:

- Say, my portfolio consists of 100 shares of Google (Alphabet Inc.) and 200 shares of Amazon. So I am a Rich Man!! 😊 Let  $Z$  be the price of my portfolio at the end of this year. Then

$$Z = 100 \cdot GOOG + 200 \cdot AMZN,$$

where  $GOOG$  and  $AMZN$  are r.v. showing the prices of Google and Amazon Stocks at the end of the year.

- My expenses at AUA cafeteria next week can be written as

$$Z = X_1 + X_3 + X_5,$$

where  $X_i$  is the amount I will spend on the day  $i$  of the week.

# Examples

Some Examples:

- Say, my portfolio consists of 100 shares of Google (Alphabet Inc.) and 200 shares of Amazon. So I am a Rich Man!! 😊 Let  $Z$  be the price of my portfolio at the end of this year. Then

$$Z = 100 \cdot GOOG + 200 \cdot AMZN,$$

where  $GOOG$  and  $AMZN$  are r.v. showing the prices of Google and Amazon Stocks at the end of the year.

- My expenses at AUA cafeteria next week can be written as

$$Z = X_1 + X_3 + X_5,$$

where  $X_i$  is the amount I will spend on the day  $i$  of the week. Of course,  $X_i$ -s are r.v.s depending on how much I will be hungry.



# Examples

Another Example:

# Examples

Another Example:

- Usually, insurance companies are interested in the amount they will pay for some future time interval, say, for the upcoming week.

# Examples

Another Example:

- Usually, insurance companies are interested in the amount they will pay for some future time interval, say, for the upcoming week. This amount can be written as:

$$Y = X_1 + X_2 + \dots + X_7,$$

where  $X_i$  is the claim size for the day  $i$  of the week.  $X_i$ -s are r.v., so  $Y$  is a r.v.

# Examples

Another Example:

- Usually, insurance companies are interested in the amount they will pay for some future time interval, say, for the upcoming week. This amount can be written as:

$$Y = X_1 + X_2 + \dots + X_7,$$

where  $X_i$  is the claim size for the day  $i$  of the week.  $X_i$ -s are r.v., so  $Y$  is a r.v.

- Going further, the same companies can also model their daily claims: If they will have  $M$  claims in a day, with sizes  $X_1, X_2, \dots, X_M$ , then their daily claim size will be

$$Z = X_1 + X_2 + \dots + X_M.$$

# Examples

Another Example:

- Usually, insurance companies are interested in the amount they will pay for some future time interval, say, for the upcoming week. This amount can be written as:

$$Y = X_1 + X_2 + \dots + X_7,$$

where  $X_i$  is the claim size for the day  $i$  of the week.  $X_i$ -s are r.v., so  $Y$  is a r.v.

- Going further, the same companies can also model their daily claims: If they will have  $M$  claims in a day, with sizes  $X_1, X_2, \dots, X_M$ , then their daily claim size will be

$$Z = X_1 + X_2 + \dots + X_M.$$

Here important is that  $M$  **will be a r.v. itself (!)**, and the distribution of  $Z$  is an 18+ topic.

# Examples

Another Example:

- Usually, insurance companies are interested in the amount they will pay for some future time interval, say, for the upcoming week. This amount can be written as:

$$Y = X_1 + X_2 + \dots + X_7,$$

where  $X_i$  is the claim size for the day  $i$  of the week.  $X_i$ -s are r.v., so  $Y$  is a r.v.

- Going further, the same companies can also model their daily claims: If they will have  $M$  claims in a day, with sizes  $X_1, X_2, \dots, X_M$ , then their daily claim size will be

$$Z = X_1 + X_2 + \dots + X_M.$$

Here important is that  $M$  **will be a r.v. itself (!)**, and the distribution of  $Z$  is an 18+ topic. , Btw, can you give a Model for  $M$ ?

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ .



# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ . That is,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y), \quad \forall x, y.$$

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ . That is,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y), \quad \forall x, y.$$

We want to find the CDF of  $Z$ ,  $F_Z(x)$ .

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ . That is,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y), \quad \forall x, y.$$

We want to find the CDF of  $Z$ ,  $F_Z(x)$ . We know that

$$F_Z(x) = \mathbb{P}(Z \leq x) =$$

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ . That is,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y), \quad \forall x, y.$$

We want to find the CDF of  $Z$ ,  $F_Z(x)$ . We know that

$$F_Z(x) = \mathbb{P}(Z \leq x) = \mathbb{P}(g(X, Y) \leq x).$$

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ . That is,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y), \quad \forall x, y.$$

We want to find the CDF of  $Z$ ,  $F_Z(x)$ . We know that

$$F_Z(x) = \mathbb{P}(Z \leq x) = \mathbb{P}(g(X, Y) \leq x).$$

Unfortunately, it is not possible to express this in terms of  $F_{X,Y}(x, y)$  in the general case.

# CDF of $g(X, Y)$ through the Joint CDF of $X, Y$

First, let us consider the problem of expressing the CDF of  $Z = g(X, Y)$  in terms of the Joint CDF of  $X$  and  $Y$ .

Assume that  $F_{X,Y}(x, y)$  is the CDF of  $(X, Y)$ . That is,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y), \quad \forall x, y.$$

We want to find the CDF of  $Z$ ,  $F_Z(x)$ . We know that

$$F_Z(x) = \mathbb{P}(Z \leq x) = \mathbb{P}(g(X, Y) \leq x).$$

Unfortunately, it is not possible to express this in terms of  $F_{X,Y}(x, y)$  in the general case.

Btw, we have calculated this kind of thing some lecture ago: we were calculating  $\mathbb{P}(Y - X \leq x)$ , i.e., the CDF of  $Z = Y - X$ .

# Transform of Joint Discrete R.V.s

Assume now  $X$  and  $Y$  are Discrete with values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , and with the Joint PMF

$Y \setminus X$	$x_1$	$x_2$	$\dots$
$y_1$	$p_{1,1}$	$p_{2,1}$	$\dots$
$y_2$	$p_{1,2}$	$p_{2,2}$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

# Transform of Joint Discrete R.V.s

Assume now  $X$  and  $Y$  are Discrete with values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , and with the Joint PMF

$Y \setminus X$	$x_1$	$x_2$	$\dots$
$y_1$	$p_{1,1}$	$p_{2,1}$	$\dots$
$y_2$	$p_{1,2}$	$p_{2,2}$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Assume also that  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function, and  $Z = g(X, Y)$ .



# Transform of Joint Discrete R.V.s

Assume now  $X$  and  $Y$  are Discrete with values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , and with the Joint PMF

$Y \setminus X$	$x_1$	$x_2$	$\dots$
$y_1$	$p_{1,1}$	$p_{2,1}$	$\dots$
$y_2$	$p_{1,2}$	$p_{2,2}$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Assume also that  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function, and  $Z = g(X, Y)$ . It is easy to see that  $Z$  is a Discrete r.v., and our aim is to find the PMF of  $Z$ .

# Transform of Joint Discrete R.V.s

Assume now  $X$  and  $Y$  are Discrete with values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , and with the Joint PMF

$Y \setminus X$	$x_1$	$x_2$	$\dots$
$y_1$	$p_{1,1}$	$p_{2,1}$	$\dots$
$y_2$	$p_{1,2}$	$p_{2,2}$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Assume also that  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function, and  $Z = g(X, Y)$ . It is easy to see that  $Z$  is a Discrete r.v., and our aim is to find the PMF of  $Z$ .

Now, the values of  $Z$  will be

# Transform of Joint Discrete R.V.s

Assume now  $X$  and  $Y$  are Discrete with values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , and with the Joint PMF

$Y \setminus X$	$x_1$	$x_2$	$\dots$
$y_1$	$p_{1,1}$	$p_{2,1}$	$\dots$
$y_2$	$p_{1,2}$	$p_{2,2}$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Assume also that  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function, and  $Z = g(X, Y)$ . It is easy to see that  $Z$  is a Discrete r.v., and our aim is to find the PMF of  $Z$ .

Now, the values of  $Z$  will be  $z_{ij} = g(x_i, y_j)$ , and the Probability of  $Z$  being  $z_{ij}$  will be

# Transform of Joint Discrete R.V.s

Assume now  $X$  and  $Y$  are Discrete with values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , and with the Joint PMF

$Y \setminus X$	$x_1$	$x_2$	$\dots$
$y_1$	$p_{1,1}$	$p_{2,1}$	$\dots$
$y_2$	$p_{1,2}$	$p_{2,2}$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Assume also that  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function, and  $Z = g(X, Y)$ . It is easy to see that  $Z$  is a Discrete r.v., and our aim is to find the PMF of  $Z$ .

Now, the values of  $Z$  will be  $z_{ij} = g(x_i, y_j)$ , and the Probability of  $Z$  being  $z_{ij}$  will be

$$\mathbb{P}(Z = z_{ij}) = \sum_{g(x_k, y_m) = z_{ij}} \mathbb{P}(X = x_k, Y = y_m).$$

# Transform of Joint Discrete R.V.s, Example

And also in this case, we can calculate the CDF of  $Z$ :

$$F_Z(x) = \mathbb{P}(Z \leq x) = \mathbb{P}(g(X, Y) \leq x) =$$

# Transform of Joint Discrete R.V.s, Example

And also in this case, we can calculate the CDF of  $Z$ :

$$F_Z(x) = \mathbb{P}(Z \leq x) = \mathbb{P}(g(X, Y) \leq x) = \sum_{g(x_i, y_j) \leq x} \mathbb{P}(X = x_i, Y = y_j).$$

# Transform of Joint Discrete R.V.s, Example

And also in this case, we can calculate the CDF of  $Z$ :

$$F_Z(x) = \mathbb{P}(Z \leq x) = \mathbb{P}(g(X, Y) \leq x) = \sum_{g(x_i, y_j) \leq x} \mathbb{P}(X = x_i, Y = y_j).$$

**Example:** Assume  $X$  and  $Y$  are given by

$Y \setminus X$	$-1$	$1$
$3$	$0.1$	$0.2$
$2$	$0.3$	$0.1$
$4$	$0.1$	$0.2$

- Find the PMF of  $Z = X + Y$ ;
- Find the CDF of  $Z = 2X - Y$ .

# Transform of Joint Continuous R.V.s

Now assume  $X$  and  $Y$  are Continuous r.v.s with Joint PDF  $f_{X,Y}(x,y)$ . Also,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function. Let  $Z = g(X, Y)$ . Again we are interested in the Distribution of  $Z$ .



# Transform of Joint Continuous R.V.s

Now assume  $X$  and  $Y$  are Continuous r.v.s with Joint PDF  $f_{X,Y}(x,y)$ . Also,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function. Let  $Z = g(X, Y)$ . Again we are interested in the Distribution of  $Z$ .

**Note:** First, note that although  $(X, Y)$  is continuous,  $Z$  is not always continuous: it can be both Discrete or Continuous.

# Transform of Joint Continuous R.V.s

Now assume  $X$  and  $Y$  are Continuous r.v.s with Joint PDF  $f_{X,Y}(x,y)$ . Also,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function. Let  $Z = g(X, Y)$ . Again we are interested in the Distribution of  $Z$ .

**Note:** First, note that although  $(X, Y)$  is continuous,  $Z$  is not always continuous: it can be both Discrete or Continuous.

**Example:**  $Z_1 = X^2 + 3X \cdot Y$  is continuous, and

# Transform of Joint Continuous R.V.s

Now assume  $X$  and  $Y$  are Continuous r.v.s with Joint PDF  $f_{X,Y}(x,y)$ . Also,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function. Let  $Z = g(X, Y)$ . Again we are interested in the Distribution of  $Z$ .

**Note:** First, note that although  $(X, Y)$  is continuous,  $Z$  is not always continuous: it can be both Discrete or Continuous.

**Example:**  $Z_1 = X^2 + 3X \cdot Y$  is continuous, and  $Z_2 = [X + Y]$  or

# Transform of Joint Continuous R.V.s

Now assume  $X$  and  $Y$  are Continuous r.v.s with Joint PDF  $f_{X,Y}(x,y)$ . Also,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function. Let  $Z = g(X, Y)$ . Again we are interested in the Distribution of  $Z$ .

**Note:** First, note that although  $(X, Y)$  is continuous,  $Z$  is not always continuous: it can be both Discrete or Continuous.

**Example:**  $Z_1 = X^2 + 3X \cdot Y$  is continuous, and  $Z_2 = [X + Y]$  or

$$Z_3 = \begin{cases} -2, & X^2 - Y \leq 2 \\ 3, & X^2 - Y > 2 \end{cases}$$

are Discrete.

# Transform of Joint Continuous R.V.s

Now assume  $X$  and  $Y$  are Continuous r.v.s with Joint PDF  $f_{X,Y}(x,y)$ . Also,  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a given function. Let  $Z = g(X, Y)$ . Again we are interested in the Distribution of  $Z$ .

**Note:** First, note that although  $(X, Y)$  is continuous,  $Z$  is not always continuous: it can be both Discrete or Continuous.

**Example:**  $Z_1 = X^2 + 3X \cdot Y$  is continuous, and  $Z_2 = \lfloor X + Y \rfloor$  or

$$Z_3 = \begin{cases} -2, & X^2 - Y \leq 2 \\ 3, & X^2 - Y > 2 \end{cases}$$

are Discrete.

**Note:** Second, we can find the CDF of  $Z$  by:

$$F_Z(x) = \mathbb{P}(g(X, Y) \leq x) = \iint_{g(u,v) \leq x} f_{X,Y}(u, v) \, du dv.$$

# The Sum of Joint Continuous R.V.s

Now, finding the PMF or the PDF of  $Z = g(X, Y)$  in the Continuous case is not so easy.

# The Sum of Joint Continuous R.V.s

Now, finding the PMF or the PDF of  $Z = g(X, Y)$  in the Continuous case is not so easy. In principle, we can calculate first the CDF by the formula above, and then differentiate it (if  $Z$  is continuous) to obtain the PDF.

# The Sum of Joint Continuous R.V.s

Now, finding the PMF or the PDF of  $Z = g(X, Y)$  in the Continuous case is not so easy. In principle, we can calculate first the CDF by the formula above, and then differentiate it (if  $Z$  is continuous) to obtain the PDF. Try to solve some problems for yourself!



# The Sum of Joint Continuous R.V.s

Now, finding the PMF or the PDF of  $Z = g(X, Y)$  in the Continuous case is not so easy. In principle, we can calculate first the CDF by the formula above, and then differentiate it (if  $Z$  is continuous) to obtain the PDF. Try to solve some problems for yourself!

Here we will consider one very important case:

# The Sum of Joint Continuous R.V.s

Now, finding the PMF or the PDF of  $Z = g(X, Y)$  in the Continuous case is not so easy. In principle, we can calculate first the CDF by the formula above, and then differentiate it (if  $Z$  is continuous) to obtain the PDF. Try to solve some problems for yourself!

Here we will consider one very important case: we will restrict our attention to the sum of  $X$  and  $Y$ ,

$$Z = X + Y.$$

# The Sum of Joint Continuous R.V.s

Now, assume

$$Z = X + Y.$$

# The Sum of Joint Continuous R.V.s

Now, assume

$$Z = X + Y.$$

- If  $X$  and  $Y$  are **Discrete**, then  $Z = X + Y$  will be Discrete too, with a PMF

$$\mathbb{P}(Z = x) = \mathbb{P}(X + Y = x) = \sum_{x_i + y_j = x} \mathbb{P}(X = x_i, Y = y_j).$$

# The Sum of Joint Continuous R.V.s

Now, assume

$$Z = X + Y.$$

- If  $X$  and  $Y$  are **Discrete**, then  $Z = X + Y$  will be Discrete too, with a PMF

$$\mathbb{P}(Z = x) = \mathbb{P}(X + Y = x) = \sum_{x_i + y_j = x} \mathbb{P}(X = x_i, Y = y_j).$$

- If  $X$  and  $Y$  are **Jointly Continuous** with the Joint PDF  $f_{X,Y}(x, y)$ , then  $Z = X + Y$  will be a Continuous r.v. with the PDF

$$f_{X+Y}(x) = \int_{-\infty}^{\infty} f_{X,Y}(t, x - t) dt \quad \forall x \in \mathbb{R}.$$