

# AUA CS 108, Statistics, Fall 2019

## Lecture 30

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

1 Nov 2019

# Contents

- ▶ The Properties of Maximum Likelihood Estimator
- ▶ Confidence Intervals (CI)

# Last Lecture ReCap

- ▶ Nothing new 😊

## Some Notes about MLE

**Note:** The MLE does not always exist;

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Note:** The MLE is not always unique;

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Note:** The MLE is not always unique;

**Note:** Sometimes it is necessary to find it numerically, say, using the NR Method

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Note:** The MLE is not always unique;

**Note:** Sometimes it is necessary to find it numerically, say, using the NR Method

**Note:** It is remarkable, that ML Estimators, in general (if they exist, of course 😊), possess some nice properties. These properties make MLE one of the widely used methods of Estimation.

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,



## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

# Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Asymptotically Normal and Efficient:

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\theta)}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Asymptotically Normal and Efficient:

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\theta)}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, as people usually use:

$$\sqrt{n} \cdot (\hat{\theta}_n^{MLE} - \theta) \xrightarrow{D} \mathcal{N}\left(0, \frac{1}{\mathcal{I}(\theta)}\right),$$

# Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Asymptotically Normal and Efficient:

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\theta)}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, as people usually use:

$$\sqrt{n} \cdot (\hat{\theta}_n^{MLE} - \theta) \xrightarrow{D} \mathcal{N}\left(0, \frac{1}{\mathcal{I}(\theta)}\right),$$

or, put in other way,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators.



## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

So, MLE is **Consistent** and **Asymptotically Efficient**.

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

So, MLE is **Consistent** and **Asymptotically Efficient**. And this is why, for large Sample Size  $n$ , MLE is the Top 1 Choice, is (almost) unbeatable.

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, in more non-rigorous terms,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, in more non-rigorous terms,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

**Note:** This is almost the above Property,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right),$$

but, instead of  $\mathcal{I}(\theta)$  we have  $\mathcal{I}(\hat{\theta}_n^{MLE})$ .

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, in more non-rigorous terms,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

**Note:** This is almost the above Property,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right),$$

but, instead of  $\mathcal{I}(\theta)$  we have  $\mathcal{I}(\hat{\theta}_n^{MLE})$ .

**Note:** We will use this later, to construct an (approximate) Confidence Interval for  $\theta$  and for testing Hypotheses about  $\theta$ .

## Properties of the MLE, Cont'd

- ▶ If  $\hat{\theta}$  is the MLE for  $\theta$ , then for any function  $g$ , the MLE of  $g(\theta)$  is  $g(\hat{\theta})$ , i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$



## Properties of the MLE, Cont'd

- ▶ If  $\hat{\theta}$  is the MLE for  $\theta$ , then for any function  $g$ , the MLE of  $g(\theta)$  is  $g(\hat{\theta})$ , i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

This is sometimes called the **invariance Principle of MLE**.

## Properties of the MLE, Cont'd

- ▶ If  $\hat{\theta}$  is the MLE for  $\theta$ , then for any function  $g$ , the MLE of  $g(\theta)$  is  $g(\hat{\theta})$ , i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

This is sometimes called the **invariance Principle of MLE**.

**Example** Find the MLE for  $\sigma$  in  $\mathcal{N}(\mu, \sigma^2)$  Model.

**Solution:** OTB

## Other Methods to construct Point Estimators/Estimates

There are other important methods to construct Estimators: e.g.

- ▶ Bayesian Estimation: Maximum APosteriori (MAP) Estimators
- ▶ Bayesian Estimation: Bayes Estimators;
- ▶ OLS
- ▶ etc

# Confidence Intervals

## Prelude

So far we have studied the Parameteric Point Estimation Problem: we were assuming that our data comes from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

## Prelude

So far we have studied the Parameteric Point Estimation Problem: we were assuming that our data comes from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ .

## Prelude

So far we have studied the Parameteric Point Estimation Problem: we were assuming that our data comes from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ . Assume  $\hat{\theta}$  is a Continuous r.v.

## Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data comes from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ . Assume  $\hat{\theta}$  is a Continuous r.v. If the True value of our Parameter is  $\theta^*$ , then

$$\mathbb{P}(\hat{\theta} = \theta^*) = 0,$$

i.e., we will (almost) **never** be correct in our guess.



## Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data comes from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ . Assume  $\hat{\theta}$  is a Continuous r.v. If the True value of our Parameter is  $\theta^*$ , then

$$\mathbb{P}(\hat{\theta} = \theta^*) = 0,$$

i.e., we will (almost) **never** be correct in our guess. Sad news!

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>,

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

And you were asking nice questions about how small is the *error* or how much sure are we in our Estimate (for the Unknown Parameter), and how large  $n$  needs to be to have a good estimate.

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

And you were asking nice questions about how small is the *error* or how much sure are we in our Estimate (for the Unknown Parameter), and how large  $n$  needs to be to have a good estimate.

Here we want to develop the theory of Confidence Intervals, which will contain answers to these questions.

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.



## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;
- ▶ which has the possible smallest length.

# CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;
- ▶ which has the possible smallest length.

Let us state this in Mathematical terms. We will consider here only 1D case, i.e., we will assume  $\theta \in \Theta \subset \mathbb{R}$ .

# Random Intervals and CI

We start by **Random Intervals**:

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ .

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ . We will call intervals of the form  $[X, Y]$  or  $(X, Y)$  or  $(X, Y]$  or  $[X, Y)$  or  $(X, +\infty), \dots$ , **random intervals**.

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ . We will call intervals of the form  $[X, Y]$  or  $(X, Y)$  or  $(X, Y]$  or  $[X, Y)$  or  $(X, +\infty), \dots$ , **random intervals**.

**Example:** Assume  $X \sim \text{Pois}(2.3)$ . Then

$$[X, X + 2)$$

is a Random Interval.

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ . We will call intervals of the form  $[X, Y]$  or  $(X, Y)$  or  $(X, Y]$  or  $[X, Y)$  or  $(X, +\infty), \dots$ , **random intervals**.

**Example:** Assume  $X \sim \text{Pois}(2.3)$ . Then

$$[X, X + 2)$$

is a Random Interval.

**Example:** Let  $X_1, X_2, \dots, X_n$  are IID r.v.s. Then

$$(\bar{X} - 0.1, \bar{X} + 0.1)$$

is a Random Interval.