

AUA CS 108, Statistics, Fall 2019

Lecture 26

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

23 Oct 2019

Contents

- ▶ Cramer-Rao Lower Bound
- ▶ Methods to Obtain/Construct Estimators: The Method of Moments

Last Lecture ReCap

- ▶ Give the definition of the Fisher Information.

Last Lecture ReCap

- ▶ Give the definition of the Fisher Information.
- ▶ Give the Cramer-Rao Lower Bound

Cramer-Rao Inequality, C-R Lower Bound, Refresher

Recall that under some regularity conditions on the family of Distributions \mathcal{F}_θ , the following holds:

Theorem (Cramer-Rao): Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$$

and the Fisher Information for the family \mathcal{F}_θ is $I(\theta)$. Assume also that $\hat{\theta}$ is an unbiased estimator for θ obtained from our Random Sample. Then, under the above mentioned regularity conditions,

$$\text{Var}(\hat{\theta}) \geq \frac{1}{n \cdot I(\theta)}.$$

Consequences of the C-R LB

We will use now the following consequence from the C-R LB:

Consequences of the C-R LB

We will use now the following consequence from the C-R LB:

If there exists an **Unbiased Estimator** $\hat{\theta}$ with

$$\text{Var}(\hat{\theta}) = \frac{1}{n \cdot I(\theta)},$$

then that Estimator is a MVUE for θ .

Consequences of the C-R LB

We will use now the following consequence from the C-R LB:

If there exists an **Unbiased Estimator** $\hat{\theta}$ with

$$\text{Var}(\hat{\theta}) = \frac{1}{n \cdot I(\theta)},$$

then that Estimator is a MVUE for θ . And recall that Unbiased Estimators satisfying this equality are called **Efficient Estimators**.

Example

Example: Show that in the Bernoulli Model, with a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p), \quad p \in [0, 1],$$

the Estimator

$$\hat{p} = \bar{X}$$

is the MVUE of p .

Example

Example: Show that in the Bernoulli Model, with a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p), \quad p \in [0, 1],$$

the Estimator

$$\hat{p} = \bar{X}$$

is the MVUE of p .

Example: Show that in the Poisson Model, with a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda), \quad \lambda > 0,$$

the Estimator

$$\hat{\lambda} = \bar{X}$$

is the MVUE of λ .

Methods to find (good) Estimators

The Problem

Here, in this Section, our Problem will be the following:

The Problem

Here, in this Section, our Problem will be the following:

We are given a Random Sample from a Parametric Family of Distributions,

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

The Problem

Here, in this Section, our Problem will be the following:

We are given a Random Sample from a Parametric Family of Distributions,

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Problem: The Problem is to find/construct a good Estimator for θ , using our Random Sample.

The Method of Moments

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Example: Say, the first Moment of \mathcal{F}_θ is just the Mean of X , $\mathbb{E}(X)$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Example: Say, the first Moment of \mathcal{F}_θ is just the Mean of X , $\mathbb{E}(X)$. The second Moment of \mathcal{F}_θ is $\mathbb{E}(X^2)$ etc.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Example: Say, the first Moment of \mathcal{F}_θ is just the Mean of X , $\mathbb{E}(X)$. The second Moment of \mathcal{F}_θ is $\mathbb{E}(X^2)$ etc.

Note: Note that, in general, the Theoretical Moments of \mathcal{F}_θ are functions of θ .

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Definition: The **Empirical (Sample) Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ is

$$\frac{(X_1)^k + (X_2)^k + \dots + (X_n)^k}{n}.$$

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Definition: The **Empirical (Sample) Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ is

$$\frac{(X_1)^k + (X_2)^k + \dots + (X_n)^k}{n}.$$

Example: Say, the second order Empirical Moment is

$$\frac{(X_1)^2 + (X_2)^2 + \dots + (X_n)^2}{n}.$$

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Definition: The **Empirical (Sample) Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ is

$$\frac{(X_1)^k + (X_2)^k + \dots + (X_n)^k}{n}.$$

Example: Say, the second order Empirical Moment is

$$\frac{(X_1)^2 + (X_2)^2 + \dots + (X_n)^2}{n}.$$

Note: The Empirical Moment is independent of the Parameter θ , it is just a Statistics, it is a function of X_1, X_2, \dots, X_n .

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ .

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**, and denoted by $\hat{\theta}^{MoM}$.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**, and denoted by $\hat{\theta}^{MoM}$.
And if this will not give a value for θ , solve instead

2-nd order Theoretical Moment = 2-nd order Empirical Moment.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**, and denoted by $\hat{\theta}^{MoM}$.
And if this will not give a value for θ , solve instead

2-nd order Theoretical Moment = 2-nd order Empirical Moment.

If this is giving the value for θ , then you have the MoM Estimator.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**, and denoted by $\hat{\theta}^{MoM}$.
And if this will not give a value for θ , solve instead

2-nd order Theoretical Moment = 2-nd order Empirical Moment.

If this is giving the value for θ , then you have the MoM Estimator. Otherwise, continue with the 3-rd Moments, then with 4-th Moments, etc.

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$.

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$. Find the MoM Estimator and MoM Estimate for θ .

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$. Find the MoM Estimator and MoM Estimate for θ .

Solution: OTB

Example: Find the MoM Estimator for λ in the *Exp*(λ) Model.

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for θ in the $Unif[0, \theta]$ Model.

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for θ in the $Unif[0, \theta]$ Model.

Solution: OTB

Example: Find the MoM Estimator for θ in the $Unif[-\theta, \theta]$ Model.

Solution: OTB