

CS 107, Probability, Spring 2019

Lecture 39

Michael Poghosyan

AUA

29 April 2019

Content

- The Expected Value of a R.V.
- The Variance of a R.V.
- Markov and Chebyshev Inequalities

Example:

Question: Is it true that

$$\mathbb{E}(X^2) = \left(\mathbb{E}(X)\right)^2?$$

Expectation and Independence

Assume X and Y are Independent r.v. Then

$$\mathbb{E}(X \cdot Y) = \mathbb{E}(X) \cdot \mathbb{E}(Y).$$

Expectation and Independence

Assume X and Y are Independent r.v. Then

$$\mathbb{E}(X \cdot Y) = \mathbb{E}(X) \cdot \mathbb{E}(Y).$$

Moreover, if g and h are any (nice!) functions, then

$$\mathbb{E}(g(X) \cdot h(Y)) = \mathbb{E}(g(X)) \cdot \mathbb{E}(h(Y)).$$

Expectation and Independence

Assume X and Y are Independent r.v. Then

$$\mathbb{E}(X \cdot Y) = \mathbb{E}(X) \cdot \mathbb{E}(Y).$$

Moreover, if g and h are any (nice!) functions, then

$$\mathbb{E}(g(X) \cdot h(Y)) = \mathbb{E}(g(X)) \cdot \mathbb{E}(h(Y)).$$

Example: Say, if $X \perp\!\!\!\perp Y$, then

$$\mathbb{E}((X + Y)^2) =$$

Expectation and Independence

Assume X and Y are Independent r.v. Then

$$\mathbb{E}(X \cdot Y) = \mathbb{E}(X) \cdot \mathbb{E}(Y).$$

Moreover, if g and h are any (nice!) functions, then

$$\mathbb{E}(g(X) \cdot h(Y)) = \mathbb{E}(g(X)) \cdot \mathbb{E}(h(Y)).$$

Example: Say, if $X \perp\!\!\!\perp Y$, then

$$\mathbb{E}((X + Y)^2) = \mathbb{E}(X^2 + 2XY + Y^2) =$$

Jensen's Inequality

Recall the Jensen's Inequality for convex functions: if $g : [a, b] \rightarrow \mathbb{R}$ is a convex function, $x_1, \dots, x_n \in [a, b]$ and $\alpha_1, \dots, \alpha_n \in [0, 1]$ with $\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$, then

$$g\left(\sum_{k=1}^n \alpha_k x_k\right) \leq \sum_{k=1}^n \alpha_k \cdot g(x_k).$$

Jensen's Inequality

Recall the Jensen's Inequality for convex functions: if $g : [a, b] \rightarrow \mathbb{R}$ is a convex function, $x_1, \dots, x_n \in [a, b]$ and $\alpha_1, \dots, \alpha_n \in [0, 1]$ with $\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$, then

$$g\left(\sum_{k=1}^n \alpha_k x_k\right) \leq \sum_{k=1}^n \alpha_k \cdot g(x_k).$$

Now, the generalization, and Probabilistic Interpretation is:

Jensen's Inequality

Recall the Jensen's Inequality for convex functions: if $g : [a, b] \rightarrow \mathbb{R}$ is a convex function, $x_1, \dots, x_n \in [a, b]$ and $\alpha_1, \dots, \alpha_n \in [0, 1]$ with $\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$, then

$$g\left(\sum_{k=1}^n \alpha_k x_k\right) \leq \sum_{k=1}^n \alpha_k \cdot g(x_k).$$

Now, the generalization, and Probabilistic Interpretation is:

Jensen's Inequality

If X is a r.v. and $g : \mathbb{R} \rightarrow \mathbb{R}$ is a **convex function**, then

$$g(\mathbb{E}(X)) \leq \mathbb{E}(g(X)).$$

Jensen's Inequality

Recall the Jensen's Inequality for convex functions: if $g : [a, b] \rightarrow \mathbb{R}$ is a convex function, $x_1, \dots, x_n \in [a, b]$ and $\alpha_1, \dots, \alpha_n \in [0, 1]$ with $\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$, then

$$g\left(\sum_{k=1}^n \alpha_k x_k\right) \leq \sum_{k=1}^n \alpha_k \cdot g(x_k).$$

Now, the generalization, and Probabilistic Interpretation is:

Jensen's Inequality

If X is a r.v. and $g : \mathbb{R} \rightarrow \mathbb{R}$ is a **convex function**, then

$$g(\mathbb{E}(X)) \leq \mathbb{E}(g(X)).$$

Example: In Economics/Finance, one is defining Risk Aversion using this Inequality.

Partial Numerical Characteristics of R.V.s: Variance and Standard Deviation of a R.V

The Variance of a R.V.

Variance is the second most important numerical (partial) characteristics of the r.v.

The Variance of a R.V.

Variance is the second most important numerical (partial) characteristics of the r.v. Knowing the average, Expected Value of the r.v. is an important thing, but this will not give a sufficient information about the r.v.

The Variance of a R.V.

Variance is the second most important numerical (partial) characteristics of the r.v. Knowing the average, Expected Value of the r.v. is an important thing, but this will not give a sufficient information about the r.v.

Say, if we know that the AMZN stock Expected price at the end of the next week will be \$2K, then this is an important information. And we imagine that the price at the end of the next week will be around \$2K. Is this giving us a nice picture what the actual price will be?

The Variance of a R.V.

Variance is the second most important numerical (partial) characteristics of the r.v. Knowing the average, Expected Value of the r.v. is an important thing, but this will not give a sufficient information about the r.v.

Say, if we know that the AMZN stock Expected price at the end of the next week will be \$2K, then this is an important information. And we imagine that the price at the end of the next week will be around \$2K. Is this giving us a nice picture what the actual price will be? No, of course.

The Variance of a R.V.

Variance is the second most important numerical (partial) characteristics of the r.v. Knowing the average, Expected Value of the r.v. is an important thing, but this will not give a sufficient information about the r.v.

Say, if we know that the AMZN stock Expected price at the end of the next week will be \$2K, then this is an important information. And we imagine that the price at the end of the next week will be around \$2K. Is this giving us a nice picture what the actual price will be? No, of course.

Consider 2 cases:

The Variance of a R.V.

The Variance of a R.V.

- The AMZN stock price at the end of the next week can be anything from the interval $[1000, 3000]$, uniformly.

The Variance of a R.V.

- The AMZN stock price at the end of the next week can be anything from the interval $[1000, 3000]$, uniformly. In this case, the Expected price will be 2000. But we have no much information about the concrete price we will have at the end of the next week, because of the wide range of possibilities.

The Variance of a R.V.

- The AMZN stock price at the end of the next week can be anything from the interval $[1000, 3000]$, uniformly. In this case, the Expected price will be 2000. But we have no much information about the concrete price we will have at the end of the next week, because of the wide range of possibilities.
- The AMZN stock price at the end of the next week can be anything from the interval $[1950, 2050]$, uniformly.

The Variance of a R.V.

- The AMZN stock price at the end of the next week can be anything from the interval $[1000, 3000]$, uniformly. In this case, the Expected price will be 2000. But we have no much information about the concrete price we will have at the end of the next week, because of the wide range of possibilities.
- The AMZN stock price at the end of the next week can be anything from the interval $[1950, 2050]$, uniformly. In this case too, the Expected price will be 2000. But here the information is much more concrete about the price.

So not only the average is important, but also the spread around that average matters!

The Variance of a R.V.

Another example: if X is the wage of a (randomly chosen working) person in Armenia, then $\mathbb{E}(X) = 172,056\text{AMD}$ (by Feb 2019, see <https://www.armstat.am/en/>).

Is this number solely giving a nice picture about the wages?

The Variance of a R.V.

Another example: if X is the wage of a (randomly chosen working) person in Armenia, then $\mathbb{E}(X) = 172,056\text{AMD}$ (by Feb 2019, see <https://www.armstat.am/en/>).

Is this number solely giving a nice picture about the wages? Well, it gives some information, but not a nice one.

The Variance of a R.V.

Another example: if X is the wage of a (randomly chosen working) person in Armenia, then $\mathbb{E}(X) = 172,056\text{AMD}$ (by Feb 2019, see <https://www.armstat.am/en/>).

Is this number solely giving a nice picture about the wages?

Well, it gives some information, but not a nice one.

Say, is it true that all persons are getting around, close to 170,000AMD?

The Variance of a R.V.

Another example: if X is the wage of a (randomly chosen working) person in Armenia, then $\mathbb{E}(X) = 172,056\text{AMD}$ (by Feb 2019, see <https://www.armstat.am/en/>).

Is this number solely giving a nice picture about the wages?

Well, it gives some information, but not a nice one.

Say, is it true that all persons are getting around, close to 170,000AMD? Well, you know the answer 😊.

The Variance of a R.V.

Another example: if X is the wage of a (randomly chosen working) person in Armenia, then $\mathbb{E}(X) = 172,056\text{AMD}$ (by Feb 2019, see <https://www.armstat.am/en/>).

Is this number solely giving a nice picture about the wages? Well, it gives some information, but not a nice one.

Say, is it true that all persons are getting around, close to 170,000AMD? Well, you know the answer 😊. Well, you know that a lot of people are getting less than 100K. And the average is 172,056 because some politicians (and also programmers) are getting more than 500K 😊

The Variance of a R.V.

Another example: if X is the wage of a (randomly chosen working) person in Armenia, then $\mathbb{E}(X) = 172,056\text{AMD}$ (by Feb 2019, see <https://www.armstat.am/en/>).

Is this number solely giving a nice picture about the wages? Well, it gives some information, but not a nice one.

Say, is it true that all persons are getting around, close to 170,000AMD? Well, you know the answer 😊. Well, you know that a lot of people are getting less than 100K. And the average is 172,056 because some politicians (and also programmers) are getting more than 500K 😊

So the spread around the Expected Value is an important characteristic!

The Variance of a R.V.

The spread around the Expected Value can be calculated in different ways. The important method is the Expected Square Deviation from the Mean:

The Variance of a R.V.

The spread around the Expected Value can be calculated in different ways. The important method is the Expected Square Deviation from the Mean:

The Variance

The **Variance** of a r.v. X is the following number:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The Variance of a R.V.

The spread around the Expected Value can be calculated in different ways. The important method is the Expected Square Deviation from the Mean:

The Variance

The **Variance** of a r.v. X is the following number:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The idea is to calculate Deviations from the Mean:

$$X - \mathbb{E}(X)$$

which is a r.v..

The Variance of a R.V.

The spread around the Expected Value can be calculated in different ways. The important method is the Expected Square Deviation from the Mean:

The Variance

The **Variance** of a r.v. X is the following number:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The idea is to calculate Deviations from the Mean:

$$X - \mathbb{E}(X)$$

which is a r.v.. Then we calculate the Square of that: $(X - \mathbb{E}(X))^2$, which is again a r.v..

The Variance of a R.V.

The spread around the Expected Value can be calculated in different ways. The important method is the Expected Square Deviation from the Mean:

The Variance

The **Variance** of a r.v. X is the following number:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The idea is to calculate Deviations from the Mean:

$$X - \mathbb{E}(X)$$

which is a r.v.. Then we calculate the Square of that: $(X - \mathbb{E}(X))^2$, which is again a r.v.. Finally, we calculate the Expected Squared Deviations, $\mathbb{E}\left((X - \mathbb{E}(X))^2\right)$, which we call the Variance of X or the Dispersion of X .

The Variance of a R.V.

So Variance shows the following: in the Mean, how far are the values of a r.v. from its Mean, in the squared distance sense?

The Variance of a R.V.

So Variance shows the following: in the Mean, how far are the values of a r.v. from its Mean, in the squared distance sense?

The drawback in calculation of the Variance is that it is in squared units. Say, if X is a price, in AMDs, then $\mathbb{E}(X)$ is again in AMD units, but $\text{Var}(X)$ will be in AMD^2 units, which is not a convenient measure to report.

The Variance of a R.V.

So Variance shows the following: in the Mean, how far are the values of a r.v. from its Mean, in the squared distance sense?

The drawback in calculation of the Variance is that it is in squared units. Say, if X is a price, in AMDs, then $\mathbb{E}(X)$ is again in AMD units, but $\text{Var}(X)$ will be in AMD^2 units, which is not a convenient measure to report.

So people use the next important characteristic of the Spread, for reporting, the Standard Deviation:

The Standard Deviation of a R.V.

The Standard Deviation

The **Standard Deviation** of a r.v. X is the square root of the Variance:

$$SD(X) = \sigma_X = \sqrt{Var(X)} = \sqrt{\mathbb{E}\left((X - \mathbb{E}(X))^2\right)}.$$

The Standard Deviation of a R.V.

The Standard Deviation

The **Standard Deviation** of a r.v. X is the square root of the Variance:

$$SD(X) = \sigma_X = \sqrt{Var(X)} = \sqrt{\mathbb{E}\left((X - \mathbb{E}(X))^2\right)}.$$

Note: Variance is easy to handle with in terms of calculations, and **Standard Deviation** is the measure that is mostly used in reporting.

The Standard Deviation of a R.V.

The Standard Deviation

The **Standard Deviation** of a r.v. X is the square root of the Variance:

$$SD(X) = \sigma_X = \sqrt{Var(X)} = \sqrt{\mathbb{E}\left((X - \mathbb{E}(X))^2\right)}.$$

Note: Variance is easy to handle with in terms of calculations, and **Standard Deviation** is the measure that is mostly used in reporting.

Note: Another important numerical characteristic of the spread around the mean is the Mean Absolute Deviation (MAD) or the Mean Absolute Error:

$$MAD(X) = \mathbb{E}(|X - \mathbb{E}(X)|).$$

Calculation Formula for the Variance

Recall the definition of the Variance:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The following formula is widely used in Prob and Stat:

Calculation Formula for the Variance

Recall the definition of the Variance:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The following formula is widely used in Prob and Stat:

Variance Calculation Formula

$$\text{Var}(X) = \mathbb{E}(X^2) - \left(\mathbb{E}(X)\right)^2.$$

Calculation Formula for the Variance

Recall the definition of the Variance:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The following formula is widely used in Prob and Stat:

Variance Calculation Formula

$$\text{Var}(X) = \mathbb{E}(X^2) - \left(\mathbb{E}(X)\right)^2.$$

Proof: OTB

Calculation Formula for the Variance

Recall the definition of the Variance:

$$\text{Var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right).$$

The following formula is widely used in Prob and Stat:

Variance Calculation Formula

$$\text{Var}(X) = \mathbb{E}(X^2) - \left(\mathbb{E}(X)\right)^2.$$

Proof: OTB

Important Consequence:

$$\mathbb{E}(X^2) = \text{Var}(X) + \left(\mathbb{E}(X)\right)^2.$$

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) =$$

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) = \sum_i (x_i - \mathbb{E}(X))^2 \cdot p_i =$$

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) = \sum_i (x_i - \mathbb{E}(X))^2 \cdot p_i = \sum_i x_i^2 p_i - \left(\sum_i x_i p_i \right)^2$$

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) = \sum_i (x_i - \mathbb{E}(X))^2 \cdot p_i = \sum_i x_i^2 p_i - \left(\sum_i x_i p_i \right)^2$$

- If X is Continuous with the PDF $f(x)$:

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) = \sum_i (x_i - \mathbb{E}(X))^2 \cdot p_i = \sum_i x_i^2 p_i - \left(\sum_i x_i p_i \right)^2$$

- If X is Continuous with the PDF $f(x)$:

$$\text{Var}(X) =$$

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) = \sum_i (x_i - \mathbb{E}(X))^2 \cdot p_i = \sum_i x_i^2 p_i - \left(\sum_i x_i p_i \right)^2$$

- If X is Continuous with the PDF $f(x)$:

$$\text{Var}(X) = \int_{-\infty}^{\infty} (x - \mathbb{E}(X))^2 f(x) dx =$$

Calculation of the Variance

Now, let us give explicitly calculation methods for the Variance of a r.v. X :

- If X is Discrete with possible values x_i , and PMF $\mathbb{P}(X = x_i) = p_i$:

$$\text{Var}(X) = \sum_i (x_i - \mathbb{E}(X))^2 \cdot p_i = \sum_i x_i^2 p_i - \left(\sum_i x_i p_i \right)^2$$

- If X is Continuous with the PDF $f(x)$:

$$\text{Var}(X) = \int_{-\infty}^{\infty} (x - \mathbb{E}(X))^2 f(x) dx = \int_{-\infty}^{\infty} x^2 f(x) dx - \left(\int_{-\infty}^{\infty} x f(x) dx \right)^2$$

Example:

Example: Calculate the Variance of a r.v. X , if

Example:

Example: Calculate the Variance of a r.v. X , if

- X is taking the values 1, 2, 5 with probabilities 0.4, 0.1, 0.5, respectively;

Example:

Example: Calculate the Variance of a r.v. X , if

- X is taking the values 1, 2, 5 with probabilities 0.4, 0.1, 0.5, respectively;
- $X \sim \text{Unif}[1, 4]$.

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $Var(X) \geq 0$ and $SD(X) \geq 0$;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $\text{Var}(X) \geq 0$ and $\text{SD}(X) \geq 0$;
- $\text{Var}(X) = 0$ iff $X = \text{Const}$ almost surely;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $\text{Var}(X) \geq 0$ and $\text{SD}(X) \geq 0$;
- $\text{Var}(X) = 0$ iff $X = \text{Const}$ almost surely;
- $\text{Var}(X + \alpha) = \text{Var}(X)$ for any real number α ;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $\text{Var}(X) \geq 0$ and $\text{SD}(X) \geq 0$;
- $\text{Var}(X) = 0$ iff $X = \text{Const}$ almost surely;
- $\text{Var}(X + \alpha) = \text{Var}(X)$ for any real number α ;
- $\text{SD}(X + \alpha) = \text{SD}(X)$ for any real number α ;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $\text{Var}(X) \geq 0$ and $\text{SD}(X) \geq 0$;
- $\text{Var}(X) = 0$ iff $X = \text{Const}$ almost surely;
- $\text{Var}(X + \alpha) = \text{Var}(X)$ for any real number α ;
- $\text{SD}(X + \alpha) = \text{SD}(X)$ for any real number α ;
- $\text{Var}(\alpha \cdot X) = \alpha^2 \cdot \text{Var}(X)$;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $\text{Var}(X) \geq 0$ and $\text{SD}(X) \geq 0$;
- $\text{Var}(X) = 0$ iff $X = \text{Const}$ almost surely;
- $\text{Var}(X + \alpha) = \text{Var}(X)$ for any real number α ;
- $\text{SD}(X + \alpha) = \text{SD}(X)$ for any real number α ;
- $\text{Var}(\alpha \cdot X) = \alpha^2 \cdot \text{Var}(X)$;
- $\text{SD}(\alpha \cdot X) = |\alpha| \cdot \text{SD}(X)$;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $\text{Var}(X) \geq 0$ and $\text{SD}(X) \geq 0$;
- $\text{Var}(X) = 0$ iff $X = \text{Const}$ almost surely;
- $\text{Var}(X + \alpha) = \text{Var}(X)$ for any real number α ;
- $\text{SD}(X + \alpha) = \text{SD}(X)$ for any real number α ;
- $\text{Var}(\alpha \cdot X) = \alpha^2 \cdot \text{Var}(X)$;
- $\text{SD}(\alpha \cdot X) = |\alpha| \cdot \text{SD}(X)$;

Note: Interpretation of the 3rd Property: If we shift X by α , then we get $X + \alpha$. Then

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $Var(X) \geq 0$ and $SD(X) \geq 0$;
- $Var(X) = 0$ iff $X = Const$ almost surely;
- $Var(X + \alpha) = Var(X)$ for any real number α ;
- $SD(X + \alpha) = SD(X)$ for any real number α ;
- $Var(\alpha \cdot X) = \alpha^2 \cdot Var(X)$;
- $SD(\alpha \cdot X) = |\alpha| \cdot SD(X)$;

Note: Interpretation of the 3rd Property: If we shift X by α , then we get $X + \alpha$. Then

- The Mean is changed by α : $\mathbb{E}(X + \alpha) = \mathbb{E}(X) + \alpha$;

Properties of a Variance and a SD

Properties of a Variance and a SD

Assume X is a r.v.. Then

- $Var(X) \geq 0$ and $SD(X) \geq 0$;
- $Var(X) = 0$ iff $X = Const$ almost surely;
- $Var(X + \alpha) = Var(X)$ for any real number α ;
- $SD(X + \alpha) = SD(X)$ for any real number α ;
- $Var(\alpha \cdot X) = \alpha^2 \cdot Var(X)$;
- $SD(\alpha \cdot X) = |\alpha| \cdot SD(X)$;

Note: Interpretation of the 3rd Property: If we shift X by α , then we get $X + \alpha$. Then

- The Mean is changed by α : $\mathbb{E}(X + \alpha) = \mathbb{E}(X) + \alpha$;
- The Deviation from the Mean is NOT Changed:
 $Var(X + \alpha) = Var(X)$.

Properties of a Variance and a SD

Another important property is the calculation of the Variance of a sum of several r.v.s. We will give the general formula later, but one important case is the following:

Properties of a Variance and a SD

Another important property is the calculation of the Variance of a sum of several r.v.s. We will give the general formula later, but one important case is the following:

Variance of a Sum of Independent RVs

Assume X and Y are Independent r.v.s, $X \perp\!\!\!\perp Y$. Then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Properties of a Variance and a SD

Another important property is the calculation of the Variance of a sum of several r.v.s. We will give the general formula later, but one important case is the following:

Variance of a Sum of Independent RVs

Assume X and Y are Independent r.v.s, $X \perp\!\!\!\perp Y$. Then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Note: Important is to remember that

Properties of a Variance and a SD

Another important property is the calculation of the Variance of a sum of several r.v.s. We will give the general formula later, but one important case is the following:

Variance of a Sum of Independent RVs

Assume X and Y are Independent r.v.s, $X \perp\!\!\!\perp Y$. Then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Note: Important is to remember that

- The property $\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$ works for ANY r.v. X and Y ;

Properties of a Variance and a SD

Another important property is the calculation of the Variance of a sum of several r.v.s. We will give the general formula later, but one important case is the following:

Variance of a Sum of Independent RVs

Assume X and Y are Independent r.v.s, $X \perp\!\!\!\perp Y$. Then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Note: Important is to remember that

- The property $\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$ works for ANY r.v. X and Y ;
- The property $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$ is NOT TRUE for any r.v.s. It is true, in particular, when $X \perp\!\!\!\perp Y$.

Properties of a Variance and a SD

Another important property is the calculation of the Variance of a sum of several r.v.s. We will give the general formula later, but one important case is the following:

Variance of a Sum of Independent RVs

Assume X and Y are Independent r.v.s, $X \perp\!\!\!\perp Y$. Then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Note: Important is to remember that

- The property $\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$ works for ANY r.v. X and Y ;
- The property $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$ is NOT TRUE for any r.v.s. It is true, in particular, when $X \perp\!\!\!\perp Y$. Later we will see that this property holds only for *uncorrelated* r.v.s.

Example

Example: Assume $X, Y \sim \text{Bernoulli}(0.5)$ and $X \perp\!\!\!\perp Y$. Calculate

$$\text{Var}(2X - 3Y + 5).$$