

AUA CS 108, Statistics, Fall 2019

Lecture 19

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

07 Oct 2019

Contents

- ▶ Parametric Models
- ▶ Statistics v3, Estimators
- ▶ Properties of Estimators: MSE

Last Lecture ReCap

- ▶ What is a Random Sample?

Last Lecture ReCap

- ▶ What is a Random Sample?
- ▶ Why we consider Random Samples?

Last Lecture ReCap

- ▶ What is a Random Sample?
- ▶ Why we consider Random Samples?
- ▶ What is the problem we consider in the Parametric Statistics?

Example

Example: We consider a Sample of heights of some persons:
 x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim$$

Example

Example: We consider a Sample of heights of some persons:
 x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 \geq 0.$$

Example

Example: We consider a Sample of heights of some persons: x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 \geq 0.$$

Here

- ▶ Our Parameter is $(\mu, \sigma^2) = \theta$ (or $(\mu, \sigma) = \theta$), which is 2D;

Example

Example: We consider a Sample of heights of some persons: x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 \geq 0.$$

Here

- ▶ Our Parameter is $(\mu, \sigma^2) = \theta$ (or $(\mu, \sigma) = \theta$), which is 2D;
- ▶ The Parameter Set is $\mathbb{R} \times [0, +\infty) = \Theta \subset \mathbb{R}^2$;

Example

Example: We consider a Sample of heights of some persons: x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 \geq 0.$$

Here

- ▶ Our Parameter is $(\mu, \sigma^2) = \theta$ (or $(\mu, \sigma) = \theta$), which is 2D;
- ▶ The Parameter Set is $\mathbb{R} \times [0, +\infty) = \Theta \subset \mathbb{R}^2$;
- ▶ The Parametric Family of Distributions is $\mathcal{N}(\mu, \sigma^2)$.

Example

Example: We consider a Sample of heights of some persons: x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 \geq 0.$$

Here

- ▶ Our Parameter is $(\mu, \sigma^2) = \theta$ (or $(\mu, \sigma) = \theta$), which is 2D;
- ▶ The Parameter Set is $\mathbb{R} \times [0, +\infty) = \Theta \subset \mathbb{R}^2$;
- ▶ The Parametric Family of Distributions is $\mathcal{N}(\mu, \sigma^2)$.

We can think about $X \sim \mathcal{N}(\mu, \sigma^2)$ as the height of a generic person.

Example

Example: We consider a Sample of heights of some persons: x_1, x_2, \dots, x_n . We assume this is a realization of a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 \geq 0.$$

Here

- ▶ Our Parameter is $(\mu, \sigma^2) = \theta$ (or $(\mu, \sigma) = \theta$), which is 2D;
- ▶ The Parameter Set is $\mathbb{R} \times [0, +\infty) = \Theta \subset \mathbb{R}^2$;
- ▶ The Parametric Family of Distributions is $\mathcal{N}(\mu, \sigma^2)$.

We can think about $X \sim \mathcal{N}(\mu, \sigma^2)$ as the height of a generic person.

Our Problem here is, using the observation x_1, x_2, \dots, x_n , to estimate μ and σ^2 .

Point Estimates

Motivating Example ☺

Motivating Example 😊

Example: I have generated the following Data from a Normal Distribution:

##		[,1]	[,2]	[,3]	[,4]	[,5]	[,6]
##	[1,]	-6.19	-7.70	-1.23	-3.77	-2.726	-4.93
##	[2,]	-7.81	-1.63	4.17	-8.51	-3.654	1.54
##	[3,]	-6.87	-4.55	-2.84	-9.44	-7.891	-9.50
##	[4,]	4.31	-7.72	-3.49	-6.95	0.447	-5.28
##	[5,]	4.27	-5.97	-4.57	-8.91	-1.306	-6.25

Motivating Example 😊

Example: I have generated the following Data from a Normal Distribution:

##		[,1]	[,2]	[,3]	[,4]	[,5]	[,6]
##	[1,]	-6.19	-7.70	-1.23	-3.77	-2.726	-4.93
##	[2,]	-7.81	-1.63	4.17	-8.51	-3.654	1.54
##	[3,]	-6.87	-4.55	-2.84	-9.44	-7.891	-9.50
##	[4,]	4.31	-7.72	-3.49	-6.95	0.447	-5.28
##	[5,]	4.27	-5.97	-4.57	-8.91	-1.306	-6.25

Question: Find/Estimate the Parameter values I was using.

Motivating Example 😊

Example: I have generated the following Data from a Normal Distribution:

```
##          [,1]  [,2]  [,3]  [,4]  [,5]  [,6]  
## [1,] -6.19 -7.70 -1.23 -3.77 -2.726 -4.93  
## [2,] -7.81 -1.63  4.17 -8.51 -3.654  1.54  
## [3,] -6.87 -4.55 -2.84 -9.44 -7.891 -9.50  
## [4,]  4.31 -7.72 -3.49 -6.95  0.447 -5.28  
## [5,]  4.27 -5.97 -4.57 -8.91 -1.306 -6.25
```

Question: Find/Estimate the Parameter values I was using.

Moral: Statistics is like a Detective Story: you need to find the Unknown (murderer?) using some small amount of Observations, Data you have 😊

Statistics, Estimator and Estimate

Let us recall what is our Problem: assume we have a Dataset x_1, \dots, x_n . We assume that this is a realization of a Random Sample X_1, \dots, X_n , coming from one of the Distributions from some Parametric Family:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

And our problem is to estimate the value of our unknown Parameter θ .

Statistics, Estimator and Estimate

Let us recall what is our Problem: assume we have a Dataset x_1, \dots, x_n . We assume that this is a realization of a Random Sample X_1, \dots, X_n , coming from one of the Distributions from some Parametric Family:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

And our problem is to estimate the value of our unknown Parameter θ .

To estimate θ , we will use only X_1, \dots, X_n (or, x_1, x_2, \dots, x_n), since we do not have any other thing.

Statistics, Estimator and Estimate

Let us recall what is our Problem: assume we have a Dataset x_1, \dots, x_n . We assume that this is a realization of a Random Sample X_1, \dots, X_n , coming from one of the Distributions from some Parametric Family:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

And our problem is to estimate the value of our unknown Parameter θ .

To estimate θ , we will use only X_1, \dots, X_n (or, x_1, x_2, \dots, x_n), since we do not have any other thing. Now,

Definition: Any (measurable) function of the Random Sample X_1, X_2, \dots, X_n is called a **Statistics**. So Statistics is a r.v. of the form

$$g(X_1, X_2, \dots, X_n).$$

Statistics, Estimator and Estimate

Let us recall what is our Problem: assume we have a Dataset x_1, \dots, x_n . We assume that this is a realization of a Random Sample X_1, \dots, X_n , coming from one of the Distributions from some Parametric Family:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

And our problem is to estimate the value of our unknown Parameter θ .

To estimate θ , we will use only X_1, \dots, X_n (or, x_1, x_2, \dots, x_n), since we do not have any other thing. Now,

Definition: Any (measurable) function of the Random Sample X_1, X_2, \dots, X_n is called a **Statistics**. So Statistics is a r.v. of the form

$$g(X_1, X_2, \dots, X_n).$$

This is our third meaning of the term *Statistics*.

Example

Example: For example, the followings are Statistics:

$$X_1, \quad \frac{X_1 + X_n}{3}, \quad \sin(X_1 \cdot X_2 + X_3 + \dots + X_n).$$

Example

Example: For example, the followings are Statistics:

$$X_1, \quad \frac{X_1 + X_n}{3}, \quad \sin(X_1 \cdot X_2 + X_3 + \dots + X_n).$$

Definition: The Distribution of the Statistics $g(X_1, X_2, \dots, X_n)$ is called a **Sampling Distribution**.

Example

Example: For example, the followings are Statistics:

$$X_1, \quad \frac{X_1 + X_n}{3}, \quad \sin(X_1 \cdot X_2 + X_3 + \dots + X_n).$$

Definition: The Distribution of the Statistics $g(X_1, X_2, \dots, X_n)$ is called a **Sampling Distribution**.

Example: The Sampling Distribution of the Statistics

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

is

Example

Example: For example, the followings are Statistics:

$$X_1, \quad \frac{X_1 + X_n}{3}, \quad \sin(X_1 \cdot X_2 + X_3 + \dots + X_n).$$

Definition: The Distribution of the Statistics $g(X_1, X_2, \dots, X_n)$ is called a **Sampling Distribution**.

Example: The Sampling Distribution of the Statistics

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

is almost Normal, for large n , by the CLT.

Statistics, Estimator and Estimate

Assume we want to estimate the value of the parameter $\theta \in \Theta$, and we will use the Statistics $g(X_1, \dots, X_n)$ for that.

Statistics, Estimator and Estimate

Assume we want to estimate the value of the parameter $\theta \in \Theta$, and we will use the Statistics $g(X_1, \dots, X_n)$ for that.

Definition: If

- ▶ $g : \mathbb{R}^n \rightarrow \Theta$;
- ▶ g doesn't depend on the unknown θ ;

then the Statistics $g(X_1, X_2, \dots, X_n)$ is called an **Estimator** for θ , and it is usually denoted by

$$\hat{\theta} = \hat{\theta}_n = g(X_1, X_2, \dots, X_n).$$

Statistics, Estimator and Estimate

Assume we want to estimate the value of the parameter $\theta \in \Theta$, and we will use the Statistics $g(X_1, \dots, X_n)$ for that.

Definition: If

- ▶ $g : \mathbb{R}^n \rightarrow \Theta$;
- ▶ g doesn't depend on the unknown θ ;

then the Statistics $g(X_1, X_2, \dots, X_n)$ is called an **Estimator** for θ , and it is usually denoted by

$$\hat{\theta} = \hat{\theta}_n = g(X_1, X_2, \dots, X_n).$$

The value of the Estimator at our observations, $g(x_1, x_2, \dots, x_n)$, is called an **Estimate** for θ , and it is again (unfortunately) denoted by $\hat{\theta} = \hat{\theta}_n$.

Example

Example: Say, we want to estimate the parameter λ in the

$$\{\text{Exp}(\lambda) : \lambda > 0\}$$

model, using the Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Exp}(\lambda).$$

Example

Example: Say, we want to estimate the parameter λ in the

$$\{Exp(\lambda) : \lambda > 0\}$$

model, using the Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} Exp(\lambda).$$

Then the followings are **Esimators**:

► $\hat{\lambda} = \frac{X_1}{3};$

Example

Example: Say, we want to estimate the parameter λ in the

$$\{Exp(\lambda) : \lambda > 0\}$$

model, using the Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} Exp(\lambda).$$

Then the followings are **Esimators**:

- ▶ $\hat{\lambda} = \frac{X_1}{3};$
- ▶ $\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_n}{n};$

Example

Example: Say, we want to estimate the parameter λ in the

$$\{Exp(\lambda) : \lambda > 0\}$$

model, using the Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} Exp(\lambda).$$

Then the followings are **Esimators**:

- ▶ $\hat{\lambda} = \frac{X_1}{3};$
- ▶ $\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_n}{n};$
- ▶ $\hat{\lambda} = (X_1 + X_2 + \dots + X_n)^2;$

Example

Example: Say, we want to estimate the parameter λ in the

$$\{Exp(\lambda) : \lambda > 0\}$$

model, using the Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} Exp(\lambda).$$

Then the followings are **Esimators**:

- ▶ $\hat{\lambda} = \frac{X_1}{3};$
- ▶ $\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_n}{n};$
- ▶ $\hat{\lambda} = (X_1 + X_2 + \dots + X_n)^2;$

And the following is not an estimator:

- ▶ $\hat{\lambda} = \frac{\lambda}{X_1 + X_n},$ since it depends on λ - the unkown parameter value.

Estimators and Estimates

Note: We require our Estimator to be independent of the Parameter θ , since we want to be able to calculate the value of the Estimator on the observation, i.e., we want to be able to calculate the Estimate, we want to make our Estimator *observable*. Since θ is unknown, then we cannot use it in the Estimator - we will not be able to calculate the Estimate.

Estimators and Estimates

Note: We require our Estimator to be independent of the Parameter θ , since we want to be able to calculate the value of the Estimator on the observation, i.e., we want to be able to calculate the Estimate, we want to make our Estimator *observable*. Since θ is unknown, then we cannot use it in the Estimator - we will not be able to calculate the Estimate.

Note: Again,

- ▶ **Estimator** is a function of our Random Sample, it is a r.v.. It doesn't depend on the unknown Parameter. Plugging the values of our observation into the Estimator we will get the Estimate

Estimators and Estimates

Note: We require our Estimator to be independent of the Parameter θ , since we want to be able to calculate the value of the Estimator on the observation, i.e., we want to be able to calculate the Estimate, we want to make our Estimator *observable*. Since θ is unknown, then we cannot use it in the Estimator - we will not be able to calculate the Estimate.

Note: Again,

- ▶ **Estimator** is a function of our Random Sample, it is a r.v.. It doesn't depend on the unknown Parameter. Plugging the values of our observation into the Estimator we will get the Estimate
- ▶ **Estimate** is a number, it is the result of plugging the observation into the Estimator.

Example

Example: Assume we want to estimate the probability of a birth (birthrate) of a girl child in Armenia. To that end, say, we ask a maternity hospital to provide the birth data for a day. And say, our data is:

b, g, g, b, b, g, b

Example

Example: Assume we want to estimate the probability of a birth (birthrate) of a girl child in Armenia. To that end, say, we ask a maternity hospital to provide the birth data for a day. And say, our data is:

$$b, g, g, b, b, g, b$$

First we need to build a Model. To that end, we encode our data by

$$x : 0, 1, 1, 0, 0, 1, 0,$$

where $b = 0$ and $g = 1$:

Example

Example: Assume we want to estimate the probability of a birth (birthrate) of a girl child in Armenia. To that end, say, we ask a maternity hospital to provide the birth data for a day. And say, our data is:

$$b, g, g, b, b, g, b$$

First we need to build a Model. To that end, we encode our data by

$$x : 0, 1, 1, 0, 0, 1, 0,$$

where $b = 0$ and $g = 1$: this is to be able to use one of our standard Distributions.

Example

Example: Assume we want to estimate the probability of a birth (birthrate) of a girl child in Armenia. To that end, say, we ask a maternity hospital to provide the birth data for a day. And say, our data is:

$$b, g, g, b, b, g, b$$

First we need to build a Model. To that end, we encode our data by

$$x : 0, 1, 1, 0, 0, 1, 0,$$

where $b = 0$ and $g = 1$: this is to be able to use one of our standard Distributions. Next, from a Dataset we pass, for a generalization, to a Random Sample

$$X_1, X_2, X_3, X_4, X_5, X_6, X_7,$$

where X_k is the gender of the k -th child *before the observation was made* ($X_k = 1$ if the child will be a girl, and 0 otherwise).

Example, cont'd

Then we will have

$$X_1, X_2, X_3, X_4, X_5, X_6, X_7 \sim \textit{Bernoulli}(p),$$

where p is the Probability of having a girl child.

Example, cont'd

Then we will have

$$X_1, X_2, X_3, X_4, X_5, X_6, X_7 \sim \text{Bernoulli}(p),$$

where p is the Probability of having a girl child. So our Model will be

$$\{\text{Bernoulli}(p) : p \in [0, 1]\},$$

and our task will be to estimate p based on our Sample x .

Example, cont'd

Then we will have

$$X_1, X_2, X_3, X_4, X_5, X_6, X_7 \sim \text{Bernoulli}(p),$$

where p is the Probability of having a girl child. So our Model will be

$$\{\text{Bernoulli}(p) : p \in [0, 1]\},$$

and our task will be to estimate p based on our Sample x .

To estimate p , let us take the following **Estimator**:

$$\hat{p} = \frac{X_1 + X_2 + X_3 + X_4 + X_5 + X_6 + X_7}{7}.$$

Example, cont'd

Then we will have

$$X_1, X_2, X_3, X_4, X_5, X_6, X_7 \sim \text{Bernoulli}(p),$$

where p is the Probability of having a girl child. So our Model will be

$$\{\text{Bernoulli}(p) : p \in [0, 1]\},$$

and our task will be to estimate p based on our Sample x .

To estimate p , let us take the following **Estimator**:

$$\hat{p} = \frac{X_1 + X_2 + X_3 + X_4 + X_5 + X_6 + X_7}{7}.$$

This is a r.v. . The **Estimate** for p , using our Observation, will be

$$\hat{p} = \frac{0 + 1 + 1 + 0 + 0 + 1 + 0}{7} = \frac{3}{7}.$$

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

- ▶ First we specify the Model and unknown Parameter(s);

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

- ▶ First we specify the Model and unknown Parameter(s);
- ▶ Then we find Estimators for our Parameter(s);

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

- ▶ First we specify the Model and unknown Parameter(s);
- ▶ Then we find Estimators for our Parameter(s);
- ▶ Prove that the Estimators found are good ones;

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

- ▶ First we specify the Model and unknown Parameter(s);
- ▶ Then we find Estimators for our Parameter(s);
- ▶ Prove that the Estimators found are good ones;
- ▶ Using Estimators and Data, calculate the Estimate for the Parameter(s).

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

- ▶ First we specify the Model and unknown Parameter(s);
- ▶ Then we find Estimators for our Parameter(s);
- ▶ Prove that the Estimators found are good ones;
- ▶ Using Estimators and Data, calculate the Estimate for the Parameter(s).

In the next few lectures, we will consider what it means that an Estimator is a good one.

Point Estimation Setup

In the Inferential Statistics part, for the Point Estimation of the Parameter(s), we usually do the following steps, after describing the problem (and, maybe after obtaining Data):

- ▶ First we specify the Model and unknown Parameter(s);
- ▶ Then we find Estimators for our Parameter(s);
- ▶ Prove that the Estimators found are good ones;
- ▶ Using Estimators and Data, calculate the Estimate for the Parameter(s).

In the next few lectures, we will consider what it means that an Estimator is a good one. Later, we will consider some general methods to find good Estimators.

Example

Example: Assume we work with the Bernoulli Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p),$$

and we want to estimate the Parameter p .

Question: Which Estimator to use?

Example

Example: Assume we work with the Bernoulli Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p),$$

and we want to estimate the Parameter p .

Question: Which Estimator to use? Say, is

$$\hat{p} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown p ?

Example

Example: Assume we work with the Bernoulli Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p),$$

and we want to estimate the Parameter p .

Question: Which Estimator to use? Say, is

$$\hat{p} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown p ?

Or, maybe,

$$\hat{p} = \frac{X_1 + X_n}{2}$$

is better?

Example

Example: Assume we work with the Bernoulli Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p),$$

and we want to estimate the Parameter p .

Question: Which Estimator to use? Say, is

$$\hat{p} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown p ?

Or, maybe,

$$\hat{p} = \frac{X_1 + X_n}{2}$$

is better?

Or, maybe

$$\hat{p} = \frac{X_{(1)} + X_{(n)}}{2} \quad \text{or} \quad \hat{p} = \text{Median}(X_1, \dots, X_n)?$$

Example

Example: Assume we work with the Gaussian Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2),$$

and we want to estimate the Parameters μ and σ^2 .

Question: Which Estimator to use?

Example

Example: Assume we work with the Gaussian Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2),$$

and we want to estimate the Parameters μ and σ^2 .

Question: Which Estimator to use? Say, is

$$\hat{\mu} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown μ ?

Example

Example: Assume we work with the Gaussian Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2),$$

and we want to estimate the Parameters μ and σ^2 .

Question: Which Estimator to use? Say, is

$$\hat{\mu} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown μ ?

And what about estimating σ^2 ? Can you suggest Estimators?

Example

Example: Assume we work with the Gaussian Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2),$$

and we want to estimate the Parameters μ and σ^2 .

Question: Which Estimator to use? Say, is

$$\hat{\mu} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown μ ?

And what about estimating σ^2 ? Can you suggest Estimators? Say, which one to choose:

$$\hat{\sigma}^2 = \left(\frac{\sum_{k=1}^n |X_k - \bar{X}_n|}{n} \right)^2 \quad \text{or} \quad \hat{\sigma}^2 = \frac{\sum_{k=1}^n (X_k - \bar{X}_n)^2}{n} \quad \text{or}$$

$$\hat{\sigma}^2 = \frac{\sum_{k=1}^n (X_k - \bar{X}_n)^2}{n - 1} \quad \text{or} \quad \hat{\sigma}^2 = \text{other Estimator?}$$

Example

Example: Assume we work with the Exponential Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Exp}(\lambda),$$

and we want to estimate the Parameter λ .

Question: Which Estimator to use?

Example

Example: Assume we work with the Exponential Model: we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Exp}(\lambda),$$

and we want to estimate the Parameter λ .

Question: Which Estimator to use? Say, is

$$\hat{\lambda} = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

good enough to estimate the unknown λ ?

Notation

Before starting, few Notations.

Notation

Before starting, few Notations. Let us consider the Model

$$\{\mathcal{F}_\theta : \theta \in \Theta\}.$$

Notation

Before starting, few Notations. Let us consider the Model

$$\{\mathcal{F}_\theta : \theta \in \Theta\}.$$

Assume $F_\theta(x) = F(x|\theta)$ is the CDF of the Distribution \mathcal{F}_θ and $f_\theta(x) = f(x|\theta)$ is the PD(M)F of \mathcal{F}_θ .

Notation

Before starting, few Notations. Let us consider the Model

$$\{\mathcal{F}_\theta : \theta \in \Theta\}.$$

Assume $F_\theta(x) = F(x|\theta)$ is the CDF of the Distribution \mathcal{F}_θ and $f_\theta(x) = f(x|\theta)$ is the PD(M)F of \mathcal{F}_θ .

If $X \sim \mathcal{F}_\theta$, then we will write

$$\mathbb{E}_\theta(X)$$

the Expected value of X : since $X \sim \mathcal{F}_\theta$, then the Expected value depends on θ .

Notation

Before starting, few Notations. Let us consider the Model

$$\{\mathcal{F}_\theta : \theta \in \Theta\}.$$

Assume $F_\theta(x) = F(x|\theta)$ is the CDF of the Distribution \mathcal{F}_θ and $f_\theta(x) = f(x|\theta)$ is the PD(M)F of \mathcal{F}_θ .

If $X \sim \mathcal{F}_\theta$, then we will write

$$\mathbb{E}_\theta(X)$$

the Expected value of X : since $X \sim \mathcal{F}_\theta$, then the Expected value depends on θ . Say, if $X \sim \mathcal{F}_\theta$ is continuous, then

$$\mathbb{E}_\theta(X) = \int_{-\infty}^{+\infty} x \cdot f(x|\theta) dx.$$

Notation

Before starting, few Notations. Let us consider the Model

$$\{\mathcal{F}_\theta : \theta \in \Theta\}.$$

Assume $F_\theta(x) = F(x|\theta)$ is the CDF of the Distribution \mathcal{F}_θ and $f_\theta(x) = f(x|\theta)$ is the PD(M)F of \mathcal{F}_θ .

If $X \sim \mathcal{F}_\theta$, then we will write

$$\mathbb{E}_\theta(X)$$

the Expected value of X : since $X \sim \mathcal{F}_\theta$, then the Expected value depends on θ . Say, if $X \sim \mathcal{F}_\theta$ is continuous, then

$$\mathbb{E}_\theta(X) = \int_{-\infty}^{+\infty} x \cdot f(x|\theta) dx.$$

And we will use $\text{Var}_\theta(X)$ for the Variance of X .

Properties of Estimators

Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator $\hat{\theta}$ to estimate θ .

Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator $\hat{\theta}$ to estimate θ .

Definition: The **Mean Squared Error** or the **Quadratic Risk** of the estimator $\hat{\theta}_n$ of θ is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator $\hat{\theta}$ to estimate θ .

Definition: The **Mean Squared Error** or the **Quadratic Risk** of the estimator $\hat{\theta}_n$ of θ is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

Note: *MSE* calculates how close are, in the Quadratic Mean sense, possible values of the Estimator $\hat{\theta}$ to the actual (unknown) value of θ .

Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator $\hat{\theta}$ to estimate θ .

Definition: The **Mean Squared Error** or the **Quadratic Risk** of the estimator $\hat{\theta}_n$ of θ is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

Note: *MSE* calculates how close are, in the Quadratic Mean sense, possible values of the Estimator $\hat{\theta}$ to the actual (unknown) value of θ . The smaller the value of *MSE*, the better, of course.

Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator $\hat{\theta}$ to estimate θ .

Definition: The **Mean Squared Error** or the **Quadratic Risk** of the estimator $\hat{\theta}_n$ of θ is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

Note: *MSE* calculates how close are, in the Quadratic Mean sense, possible values of the Estimator $\hat{\theta}$ to the actual (unknown) value of θ . The smaller the value of *MSE*, the better, of course.

Definition: We say that the estimator $\hat{\theta}_n^1$ of θ is **preferable** to $\hat{\theta}_n^2$, another estimator of θ , if

$$MSE(\hat{\theta}_n^1, \theta) \leq MSE(\hat{\theta}_n^2, \theta), \quad \forall \theta \in \Theta,$$

and there exists a θ s.t. $MSE(\hat{\theta}_n^1, \theta) < MSE(\hat{\theta}_n^2, \theta)$.