

CS 107, Probability, Spring 2019

Lecture 03

Michael Poghosyan

AUA

21 January 2019

- Probability Measure
- Properties of the Probability Measure

Linda is thirty-one years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

Give your estimates for the probabilities that Linda is

- (1) an elementary school teacher,
- (2) active in the feminist movement,
- (3) a bank teller,
- (4) an insurance salesperson,
- (5) a bank teller also active in the feminist movement.

Linda is thirty-one years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

Give your estimates for the probabilities that Linda is

- (1) an elementary school teacher,
- (2) active in the feminist movement,
- (3) a bank teller,
- (4) an insurance salesperson,
- (5) a bank teller also active in the feminist movement.

This question is by Daniel Kahneman, psychologist, 2002 Nobel Prize receiver

Probability (Measure) Definition

Assume we have an Experiment with the Sample Space Ω and let \mathcal{F} be the set of all Events in our Experiment.

So $A \in \mathcal{F}$ means that A is an Event in our Experiment. For example, $\Omega \in \mathcal{F}$ and $\emptyset \in \mathcal{F}$

Probability (Measure) Definition

Assume we have an Experiment with the Sample Space Ω and let \mathcal{F} be the set of all Events in our Experiment.

So $A \in \mathcal{F}$ means that A is an Event in our Experiment. For example, $\Omega \in \mathcal{F}$ and $\emptyset \in \mathcal{F}$

We want to be able to measure the Probability of *any Event*, so the Probability Measure is a function defined on all events:

Probability (Measure) Definition

Probability Measure Definition

A function $\mathbb{P} : \mathcal{F} \rightarrow \mathbb{R}$ is called a **Probability Measure** on (Ω, \mathcal{F}) , if it satisfies the following axioms:

P1. For any $A \in \mathcal{F}$,

$$\mathbb{P}(A) \geq 0;$$

Probability (Measure) Definition

Probability Measure Definition

A function $\mathbb{P} : \mathcal{F} \rightarrow \mathbb{R}$ is called a **Probability Measure** on (Ω, \mathcal{F}) , if it satisfies the following axioms:

P1. For any $A \in \mathcal{F}$,

$$\mathbb{P}(A) \geq 0;$$

P2. $\mathbb{P}(\Omega) = 1;$

Probability (Measure) Definition

Probability Measure Definition

A function $\mathbb{P} : \mathcal{F} \rightarrow \mathbb{R}$ is called a **Probability Measure** on (Ω, \mathcal{F}) , if it satisfies the following axioms:

P1. For any $A \in \mathcal{F}$,

$$\mathbb{P}(A) \geq 0;$$

P2. $\mathbb{P}(\Omega) = 1$;

P3. For any sequence of pairwise mutually exclusive (disjoint) events $A_n \in \mathcal{F}$, i.e., for any sequence $A_n \in \mathcal{F}$ with $A_i \cap A_j = \emptyset$ for $i \neq j$, we have

$$\mathbb{P}\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mathbb{P}(A_n).$$

Some Remarks and Facts:

- This is a formal definition of **what is a Probability Measure**, what properties need to share the Probability measure, but we are not giving **how to define the Probability Measure**;

Some Remarks and Facts:

- This is a formal definition of **what is a Probability Measure**, what properties need to share the Probability measure, but we are not giving **how to define the Probability Measure**;
- One can define different Probability Measures on the Same Experiment;

Some Remarks and Facts:

- This is a formal definition of **what is a Probability Measure**, what properties need to share the Probability measure, but we are not giving **how to define the Probability Measure**;
- One can define different Probability Measures on the Same Experiment;
- Probability Measure is very similar (and shares the properties of) any other Measure - Cardinality (no. of elements), Length (in 1D), Area (in 2D), Volume (in 3D and moreD).

Some Remarks and Facts:

- This is a formal definition of **what is a Probability Measure**, what properties need to share the Probability measure, but we are not giving **how to define the Probability Measure**;
- One can define different Probability Measures on the Same Experiment;
- Probability Measure is very similar (and shares the properties of) any other Measure - Cardinality (no. of elements), Length (in 1D), Area (in 2D), Volume (in 3D and moreD).
- The difference is only that the Probability of the Sample Space is 1, $\mathbb{P}(\Omega) = 1$.

Examples

- Experiment: we are rolling a die.

Examples

- Experiment: we are rolling a die.
- The Sample Space in this Example is

$$\Omega = \{\blacksquare_1, \blacksquare_2, \blacksquare_3, \blacksquare_4, \blacksquare_5, \blacksquare_6\}$$

Examples

- Experiment: we are rolling a die.
- The Sample Space in this Example is

$$\Omega = \{\blacksquare, \blacksquare, \blacksquare, \blacksquare, \blacksquare, \blacksquare\}$$

- We can define (this is a definition!) the Probability Measure on this Experiment as:

$$\mathbb{P}(\Omega) = 1,$$

$$\begin{aligned}\mathbb{P}(\{\blacksquare\}) &= \mathbb{P}(\{\blacksquare\}) = \mathbb{P}(\{\blacksquare\}) = \mathbb{P}(\{\blacksquare\}) = \\ &= \mathbb{P}(\{\blacksquare\}) = \mathbb{P}(\{\blacksquare\}) = \frac{1}{6}\end{aligned}$$

Examples, cont'd

- Note that I am not writing $\mathbb{P}(\blacksquare)$, but $\mathbb{P}(\{\blacksquare\})$, since the Probability is defined on **Events**, but not Outcomes. But usually people use the notation $\mathbb{P}(\blacksquare)$ for simplicity, and we will do that.

Examples, cont'd

- Note that I am not writing $\mathbb{P}(\blacksquare)$, but $\mathbb{P}(\{\blacksquare\})$, since the Probability is defined on **Events**, but not Outcomes. But usually people use the notation $\mathbb{P}(\blacksquare)$ for simplicity, and we will do that.
- Another note: the definition above is not complete: we need to define the Probability of **any Event**, so we need to continue our definition by:

$$\mathbb{P}(\{\blacksquare, \blacksquare\}) = \mathbb{P}(\{\blacksquare, \blacksquare\}) = \dots = \mathbb{P}(\{\blacksquare, \blacksquare, \blacksquare\}) = \frac{2}{6},$$

also

$$\mathbb{P}(\{\blacksquare, \blacksquare, \blacksquare\}) = \frac{3}{6}$$

etc., and also $\mathbb{P}(\emptyset) = 0$.

Another Example

- Experiment: we are rolling a die, again.

Another Example

- Experiment: we are rolling a die, again.
- You know the Sample Space 😊

Another Example

- Experiment: we are rolling a die, again.
- You know the Sample Space 😊
- Now we take any non-negative numbers p_1, p_2, \dots, p_6 with

$$p_1 + p_2 + \dots + p_6 = 1,$$

and define

$$\mathbb{P}\left(\left\{\begin{array}{|c|c|c|}\hline \blacksquare & & \\ \hline \bullet & & \\ \hline\end{array}\right\}\right) = p_1, \quad \mathbb{P}\left(\left\{\begin{array}{|c|c|c|}\hline \blacksquare & & \\ \hline \bullet & \bullet & \\ \hline\end{array}\right\}\right) = p_2, \quad \mathbb{P}\left(\left\{\begin{array}{|c|c|c|}\hline \blacksquare & & \\ \hline \bullet & \bullet & \bullet \\ \hline\end{array}\right\}\right) = p_3,$$

$$\mathbb{P}\left(\left\{\begin{array}{|c|c|c|}\hline \blacksquare & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline\end{array}\right\}\right) = p_4, \quad \mathbb{P}\left(\left\{\begin{array}{|c|c|c|}\hline \blacksquare & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline\end{array}\right\}\right) = p_5, \quad \mathbb{P}\left(\left\{\begin{array}{|c|c|c|}\hline \blacksquare & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline\end{array}\right\}\right) = p_6.$$

Another Example

- Experiment: we are rolling a die, again.
- You know the Sample Space ☺
- Now we take any non-negative numbers p_1, p_2, \dots, p_6 with

$$p_1 + p_2 + \dots + p_6 = 1,$$

and define

$$\mathbb{P}(\{\blacksquare\bullet\}) = p_1, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\}) = p_2, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\bullet\}) = p_3,$$

$$\mathbb{P}(\{\blacksquare\bullet\bullet\bullet\bullet\}) = p_4, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\bullet\bullet\bullet\}) = p_5, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\bullet\bullet\bullet\bullet\}) = p_6.$$

and also:

$$\mathbb{P}(\{\blacksquare\bullet, \blacksquare\bullet\bullet\}) = p_1 + p_2, \quad \mathbb{P}(\{\blacksquare\bullet\bullet, \blacksquare\bullet\bullet\bullet, \blacksquare\bullet\bullet\bullet\bullet\}) = p_3 + p_4 + p_6$$

etc, and $\mathbb{P}(\emptyset) = 0$.

Another Example

- Experiment: we are rolling a die, again.
- You know the Sample Space ☺
- Now we take any non-negative numbers p_1, p_2, \dots, p_6 with

$$p_1 + p_2 + \dots + p_6 = 1,$$

and define

$$\mathbb{P}(\{\blacksquare\bullet\}) = p_1, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\}) = p_2, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\bullet\}) = p_3,$$

$$\mathbb{P}(\{\blacksquare\bullet\bullet\bullet\bullet\}) = p_4, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\bullet\bullet\bullet\}) = p_5, \quad \mathbb{P}(\{\blacksquare\bullet\bullet\bullet\bullet\bullet\bullet\}) = p_6.$$

and also:

$$\mathbb{P}(\{\blacksquare\bullet, \blacksquare\bullet\bullet\}) = p_1 + p_2, \quad \mathbb{P}(\{\blacksquare\bullet\bullet, \blacksquare\bullet\bullet\bullet, \blacksquare\bullet\bullet\bullet\bullet\}) = p_3 + p_4 + p_6$$

etc, and $\mathbb{P}(\emptyset) = 0$. In this way we will obtain a Probability Measure.

Remarks and Facts:

As we have seen in the above examples, we need to give how to calculate the Probability of *each Event*.

¹More precisely and correctly, the Probability of each Events containing one Outcome.

Remarks and Facts:

As we have seen in the above examples, we need to give how to calculate the Probability of *each Event*. Even in the above examples, where the Sample Spaces are finite, listing of all values of Probability Measure is very long and boring.

¹More precisely and correctly, the Probability of each Events containing one Outcome.

Remarks and Facts:

As we have seen in the above examples, we need to give how to calculate the Probability of *each Event*. Even in the above examples, where the Sample Spaces are finite, listing of all values of Probability Measure is very long and boring.

Fortunately, in the case of Experiments with Discrete Sample Spaces (i.e., Sample Spaces, that are either Finite or Countably Infinite), one can just define the Probability of each Outcome¹. Then, using the Properties of the Probability Measure, we will be able to calculate the Probability of each Event in that Experiment!

¹More precisely and correctly, the Probability of each Events containing one Outcome.

Remarks and Facts: Cont'd

- Say, in the Coin Tossing Experiment, with the

$$\text{Sample Space} = \{H, T\},$$

we can just give

$$\mathbb{P}(\{H\}) \quad \text{and} \quad \mathbb{P}(\{T\}),$$

Remarks and Facts: Cont'd

- Say, in the Coin Tossing Experiment, with the

$$\text{Sample Space} = \{H, T\},$$

we can just give

$$\mathbb{P}(\{H\}) \quad \text{and} \quad \mathbb{P}(\{T\}),$$

and then we can calculate $\mathbb{P}(\{H, T\})$ by using the property of a Probability Measure that we will give soon:

$$\mathbb{P}(\{H, T\}) = \mathbb{P}(\{H\}) + \mathbb{P}(\{T\}).$$

It remains only to add $\mathbb{P}(\emptyset) = 0$.

Remarks and Facts: Cont'd

- Unfortunately, the same is not the case for Experiments with Continuous Sample Spaces.

²in our example - uniformly

Remarks and Facts: Cont'd

- Unfortunately, the same is not the case for Experiments with Continuous Sample Spaces.
- But still, it will be possible to define the Probability of some Basic Events, and then to extend Probability calculations to any Event.

²in our example - uniformly

Remarks and Facts: Cont'd

- Unfortunately, the same is not the case for Experiments with Continuous Sample Spaces.
- But still, it will be possible to define the Probability of some Basic Events, and then to extend Probability calculations to any Event.
- This is an 18+ topic covered in advanced Probability courses with Measure Theory, and we will skip this part.

²in our example - uniformly

Remarks and Facts: Cont'd

- Unfortunately, the same is not the case for Experiments with Continuous Sample Spaces.
- But still, it will be possible to define the Probability of some Basic Events, and then to extend Probability calculations to any Event.
- This is an 18+ topic covered in advanced Probability courses with Measure Theory, and we will skip this part.
- As an example, if we consider an Experiment of choosing a point (a real number) from $[0, 1] = \Omega$ at random ², then we can choose as Basic Events all intervals $[a, b] \subset [0, 1]$, and define

$$\mathbb{P}([a, b]) = \text{Length}([a, b]) = b - a.$$

²in our example - uniformly

Remarks and Facts: Cont'd

- Unfortunately, the same is not the case for Experiments with Continuous Sample Spaces.
- But still, it will be possible to define the Probability of some Basic Events, and then to extend Probability calculations to any Event.
- This is an 18+ topic covered in advanced Probability courses with Measure Theory, and we will skip this part.
- As an example, if we consider an Experiment of choosing a point (a real number) from $[0, 1] = \Omega$ at random ², then we can choose as Basic Events all intervals $[a, b] \subset [0, 1]$, and define

$$\mathbb{P}([a, b]) = \text{Length}([a, b]) = b - a.$$

Using this, we will be able to calculate the Probability of
any Event (there is another 18+ topic here!)

²in our example - uniformly

Properties of the Probability Measure

1. $\mathbb{P}(\emptyset) = 0$;

Properties of the Probability Measure

1. $\mathbb{P}(\emptyset) = 0$;
2. if $A, B \in \mathcal{F}$ are mutually exclusive events, i.e., if $A \cap B = \emptyset$, then

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B);$$

Properties of the Probability Measure

1. $\mathbb{P}(\emptyset) = 0$;
2. if $A, B \in \mathcal{F}$ are mutually exclusive events, i.e., if $A \cap B = \emptyset$, then

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B);$$

3. for any event $A \in \mathcal{F}$,

$$\mathbb{P}(\overline{A}) = 1 - \mathbb{P}(A);$$

Here $\overline{A} = A^c = \Omega \setminus A$.

Properties of the Probability Measure

4. If $A_1, A_2, \dots, A_n \in \mathcal{F}$ are pairwise disjoint (mutually exclusive), i.e., if $A_i \cap A_j = \emptyset$ for $i \neq j$, then

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n \mathbb{P}(A_i);$$

Properties of the Probability Measure

4. If $A_1, A_2, \dots, A_n \in \mathcal{F}$ are pairwise disjoint (mutually exclusive), i.e., if $A_i \cap A_j = \emptyset$ for $i \neq j$, then

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n \mathbb{P}(A_i);$$

5. for any events $A, B \in \mathcal{F}$ (not necessarily disjoint),

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B);$$

Intermezzo

Recall the problem from the last lecture:

Assume our mobile phone Weather App says that there is a 50% chance of snow for this Saturday, and also 50% chance of snow this Sunday.

Is it true that it will snow for sure (i.e., with probability 1) this weekend?

What is your answer now?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- How to define the Probability Measure in the Discrete Case?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- How to define the Probability Measure in the Discrete Case?
- For the coin toss experiment, what is $\mathbb{P}(H)$?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- How to define the Probability Measure in the Discrete Case?
- For the coin toss experiment, what is $\mathbb{P}(H)$?
- Is it true that $\mathbb{P}(\{H\}) = \mathbb{P}(\{T\})$ in the coin toss experiment?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- How to define the Probability Measure in the Discrete Case?
- For the coin toss experiment, what is $\mathbb{P}(H)$?
- Is it true that $\mathbb{P}(\{H\}) = \mathbb{P}(\{T\})$ in the coin toss experiment?
- What is $\mathbb{P}(\bar{A})$?

Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- How to define the Probability Measure in the Discrete Case?
- For the coin toss experiment, what is $\mathbb{P}(H)$?
- Is it true that $\mathbb{P}(\{H\}) = \mathbb{P}(\{T\})$ in the coin toss experiment?
- What is $\mathbb{P}(\bar{A})$?
- What is $\mathbb{P}(A \cup B)$?